

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Talmanco 20 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 20 mg tadalafil.

Excipient with known effect

Each film-coated tablet contains 237.9 mg lactose (234.5 mg as anhydrous and 3.4 mg as monohydrate).

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet.

A white, film-coated, round, biconvex, bevelled edge tablet (approximately 10.7 mm diameter) debossed with 'M' on one side of the tablet and 'TA20' on the other side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Talmanco is indicated in adults for the treatment of pulmonary arterial hypertension (PAH) classified as WHO functional class II and III, to improve exercise capacity (see section 5.1).

Efficacy has been shown in idiopathic PAH (IPAH) and in PAH related to collagen vascular disease.

4.2 Posology and method of administration

Treatment should only be initiated and monitored by a physician experienced in the treatment of PAH.

Posology

The recommended dose is 40 mg (2 x 20 mg) taken once daily with or without food.

Elderly

Dose adjustments are not required in elderly patients.

Renal impairment

In patients with mild to moderate renal impairment a starting dose of 20 mg once per day is recommended. The dose may be increased to 40 mg once per day, based on individual efficacy and tolerability. In patients with severe renal impairment the use of tadalafil is not recommended (see sections 4.4 and 5.2).

Hepatic impairment

Due to limited clinical experience in patients with mild to moderate hepatic cirrhosis (Child-Pugh Class A and B), following single doses of 10 mg, a starting dose of 20 mg once per day may be considered. If tadalafil is prescribed, a careful individual benefit/risk evaluation should be undertaken by the prescribing physician. Patients with severe hepatic cirrhosis (Child-Pugh Class C) have not been studied and therefore dosing of tadalafil is not recommended (see sections 4.4 and 5.2).

Paediatric population

The safety and efficacy of tadalafil in the paediatric population has not yet been established. Currently available data are described in section 5.1.

Method of administration

Talmanco is for oral use.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Acute myocardial infarction within the last 90 days.

Severe hypotension (< 90/50 mm Hg).

- In clinical studies, tadalafil was shown to augment the hypotensive effects of nitrates. This is thought to result from the combined effects of nitrates and tadalafil on the nitric oxide/cGMP pathway. Therefore, administration of tadalafil to patients who are using any form of organic nitrate is contraindicated (see section 4.5).

The co-administration of PDE5 inhibitors, including tadalafil, with guanylate cyclase stimulators, such as riociguat, is contraindicated as it may potentially lead to symptomatic hypotension (see section 4.5).

Patients who have loss of vision in one eye because of non-arteritic anterior ischemic optic neuropathy (NAION), regardless of whether this episode was in connection or not with previous PDE5 inhibitor exposure (see section 4.4).

4.4 Special warnings and precautions for use

Cardiovascular

The following groups of patients with cardiovascular disease were not included in PAH clinical studies:

- Patients with clinically significant aortic and mitral valve disease
- Patients with pericardial constriction
- Patients with restrictive or congestive cardiomyopathy
- Patients with significant left ventricular dysfunction
- Patients with life-threatening arrhythmias
- Patients with symptomatic coronary artery disease
- Patients with uncontrolled hypertension.

Since there are no clinical data on the safety of tadalafil in these patients, the use of tadalafil is not recommended.

Pulmonary vasodilators may significantly worsen the cardiovascular status of patients with pulmonary veno-occlusive disease (PVOD). Since there are no clinical data on administration of tadalafil to patients with veno-occlusive disease, administration of tadalafil to such patients is not recommended. Should signs of pulmonary oedema occur when tadalafil is administered, the possibility of associated PVOD should be considered.

Tadalafil has systemic vasodilatory properties that may result in transient decreases in blood pressure. Physicians should carefully consider whether their patients with certain underlying conditions, such as severe left ventricular outflow obstruction, fluid depletion, autonomic hypotension or patients with resting hypotension, could be adversely affected by such vasodilatory effects.

In patients who are taking α_1 blockers concomitant administration of tadalafil may lead to symptomatic hypotension in some patients (see section 4.5). Therefore, the combination of tadalafil and doxazosin is not recommended.

Vision

Visual defects and cases of NAION have been reported in connection with the intake of tadalafil and other PDE5 inhibitors. Analyses of observational data suggest an increased risk of acute NAION in men with erectile dysfunction following exposure to tadalafil or other PDE5 inhibitors. As this may be relevant for all patients exposed to tadalafil, the patient should be advised that in case of sudden visual defect, he should stop taking tadalafil and consult a physician immediately (see section 4.3). Patients with known hereditary degenerative retinal disorders, including retinitis pigmentosa, were not included in the clinical studies, and use in these patients is not recommended.

Decreased or sudden hearing loss

Cases of sudden hearing loss have been reported after the use of tadalafil. Although other risk factors were present in some cases (such as age, diabetes, hypertension, previous hearing loss history and associated connective tissue diseases) patients should be advised to seek prompt medical attention in the event of sudden decrease or loss of hearing.

Renal and hepatic impairment

Due to increased tadalafil exposure (AUC), limited clinical experience, and the lack of ability to influence clearance by dialysis, tadalafil is not recommended in patients with severe renal impairment.

Patients with severe hepatic cirrhosis (Child-Pugh Class C) have not been studied and, therefore, dosing of tadalafil is not recommended.

Priapism and anatomical deformation of the penis

Priapism has been reported in men treated with PDE5 inhibitors. Patients who experience erections lasting 4 hours or more should be instructed to seek immediate medical assistance. If priapism is not treated immediately, penile tissue damage and permanent loss of potency may result.

Tadalafil should be used with caution in patients with anatomical deformation of the penis (such as angulation, cavernosal fibrosis or Peyronie's disease), or in patients who have conditions which may predispose them to priapism (such as sickle cell anaemia, multiple myeloma or leukaemia).

Use with CYP3A4 inducers or inhibitors

For patients chronically taking potent inducers of CYP3A4, such as rifampicin, the use of tadalafil is not recommended (see section 4.5).

For patients taking concomitant potent inhibitors of CYP3A4, such as ketoconazole or ritonavir, the use of tadalafil is not recommended (see section 4.5).

Treatments for erectile dysfunction

The safety and efficacy of combinations of tadalafil and other PDE5 inhibitors or other treatments for erectile dysfunction have not been studied. Patients should be informed not to take tadalafil with these medicinal products.

Prostacyclin and its analogues

The efficacy and safety of tadalafil co-administered with prostacyclin or its analogues has not been studied in controlled clinical studies. Therefore, caution is recommended in case of co-administration.

Bosentan

The efficacy of tadalafil in patients already on bosentan therapy has not been conclusively demonstrated (see sections 4.5 and 5.1).

Excipients with known effects

Talmanco tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

This medicinal product contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Effects of other substances on tadalafil

Cytochrome P450 Inhibitors

Azole Antifungals (e.g. ketoconazole)

Ketoconazole (200 mg daily), increased tadalafil (10 mg) single dose exposure (AUC) 2-fold and C_{max} by 15%, relative to the AUC and C_{max} values for tadalafil alone. Ketoconazole (400 mg daily) increased tadalafil (20 mg) single dose exposure (AUC) 4-fold and C_{max} by 22%.

Protease inhibitors (e.g. ritonavir)

Ritonavir (200 mg twice daily), which is an inhibitor of CYP3A4, CYP2C9, CYP2C19, and CYP2D6, increased tadalafil (20 mg) single dose exposure (AUC) 2-fold with no change in C_{max} . Ritonavir (500 mg or 600 mg twice daily) increased tadalafil (20 mg) single-dose exposure (AUC) by 32% and decreased C_{max} by 30%.

Cytochrome P450 Inducers

Endothelin-1 receptor antagonists (e.g. bosentan)

Bosentan (125 mg twice daily), a substrate of CYP2C9 and CYP3A4 and a moderate inducer of CYP3A4, CYP2C9 and possibly CYP2C19, reduced tadalafil (40 mg once per day) systemic exposure by 42% and C_{max} by 27% following multiple dose co-administration. The efficacy of tadalafil in patients already on bosentan therapy has not been conclusively demonstrated (see sections 4.4 and 5.1). Tadalafil did not affect the exposure (AUC and C_{max}) of bosentan or its metabolites.

The safety and efficacy of combinations of tadalafil and other endothelin-1 receptor antagonists have not been studied.

Antimicrobial medicinal products (e.g. rifampicin)

A CYP3A4 inducer, rifampicin (600 mg daily), reduced tadalafil AUC by 88% and C_{max} by 46%, relative to the AUC and C_{max} values for tadalafil alone (10 mg).

Effects of tadalafil on other medicinal products

Nitrates

In clinical studies, tadalafil (5, 10 and 20 mg) was shown to augment the hypotensive effects of nitrates. This interaction lasted for more than 24 hours and was no longer detectable when 48 hours had elapsed after the last tadalafil dose. Therefore, administration of tadalafil to patients who are using any form of organic nitrate is contraindicated (see section 4.3).

Anti-hypertensives (including Calcium channel blockers)

The co-administration of doxazosin (4 and 8 mg daily) and tadalafil (5 mg daily dose and 20 mg as a single dose) increases the blood pressure-lowering effect of this alpha-blocker in a significant manner. This effect lasts at least twelve hours and may be symptomatic, including syncope. Therefore, this combination is not recommended (see section 4.4).

In interaction studies performed in a limited number of healthy volunteers, these effects were not reported with alfuzosin or tamsulosin.

In clinical pharmacology studies, the potential for tadalafil (10 and 20 mg) to augment the hypotensive effects of antihypertensive medicinal products was examined. Major classes of antihypertensive medicinal products were studied either as monotherapy or as part of combination therapy. In patients taking multiple antihypertensive medicinal products whose hypertension was not well controlled, greater reductions in blood pressure were observed compared to patients whose blood pressure was well controlled, where the reduction was minimal and similar to that in healthy subjects. In patients receiving concomitant antihypertensive medicinal products, tadalafil 20 mg may induce a blood pressure decrease, which (with the exception of doxazosin -see above) is, in general, minor and not likely to be clinically relevant.

Riociguat

Preclinical studies showed an additive systemic blood pressure lowering effect when PDE5 inhibitors were combined with riociguat. In clinical studies, riociguat has been shown to augment the hypotensive effects of PDE5 inhibitors. There was no evidence of favourable clinical effect of the combination in the population studied. Concomitant use of riociguat with PDE5 inhibitors, including tadalafil, is contraindicated (see section 4.3).

Alcohol

Alcohol concentrations were not affected by co-administration with tadalafil (10 mg or 20 mg). In addition, no changes in tadalafil concentrations were seen after co-administration with alcohol. Tadalafil (20 mg) did not augment the mean blood pressure decrease produced by alcohol (0.7 g/kg or approximately 180 ml of 40% alcohol [vodka] in an 80 kg male), but in some subjects, postural dizziness and orthostatic hypotension were observed. The effect of alcohol on cognitive function was not augmented by tadalafil (10 mg).

CYP1A2 substrates (e.g. theophylline)

When tadalafil 10 mg was administered with theophylline (a non-selective phosphodiesterase inhibitor) there was no pharmacokinetic interaction. The only pharmacodynamic effect was a small (3.5 bpm) increase in heart rate.

CYP2C9 substrates (e.g. R-warfarin)

Tadalafil (10 mg and 20 mg) had no clinically significant effect on exposure (AUC) to S-warfarin or R-warfarin (CYP2C9 substrate), nor did tadalafil affect changes in prothrombin time induced by warfarin.

Acetylsalicylic acid

Tadalafil (10 mg and 20 mg) did not potentiate the increase in bleeding time caused by acetyl salicylic acid.

P-glycoprotein substrates (e.g. digoxin)

Tadalafil (40 mg once per day) had no clinically significant effect on the pharmacokinetics of digoxin.

Oral contraceptive pill

At steady-state, tadalafil (40 mg once per day) increased ethinylestradiol exposure (AUC) by 26% and C_{max} by 70% relative to oral contraceptive administered with placebo. There was no statistically significant effect of tadalafil on levonorgestrel which suggests the effect of ethinylestradiol is due to inhibition of gut sulphation by tadalafil. The clinical relevance of this finding is uncertain.

Terbutaline

A similar increase in AUC and C_{max} seen with ethinylestradiol may be expected with oral administration of terbutaline, probably due to inhibition of gut sulphation by tadalafil. The clinical relevance of this finding is uncertain.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are limited data from the use of tadalafil in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development (see section 5.3). As a precautionary measure, it is preferable to avoid the use of tadalafil during pregnancy.

Breast-feeding

Available pharmacodynamic/toxicological data in animals have shown excretion of tadalafil in milk. A risk to the breast-fed child cannot be excluded. Tadalafil should not be used during breast-feeding.

Fertility

Effects were seen in dogs that might indicate impairment of fertility. Two subsequent clinical studies suggest that this effect is unlikely in humans, although a decrease in sperm concentration was seen in some men (see sections 5.1 and 5.3).

4.7 Effects on ability to drive and use machines

Tadalafil has negligible influence on the ability to drive or use machines. Although the frequency of reports of dizziness in placebo and tadalafil arms in clinical studies was similar, patients should be aware of how they react to tadalafil, before driving or operating machinery.

4.8 Undesirable effects

Summary of the safety profile

The most commonly reported adverse reactions, occurring in $\geq 10\%$ of patients in the tadalafil 40 mg treatment arm, were headache, nausea, back pain, dyspepsia, flushing, myalgia, nasopharyngitis and pain in extremity. The adverse reactions reported were transient, and generally mild or moderate. Adverse reaction data are limited in patients over 75 years of age.

In the pivotal placebo-controlled study of tadalafil for the treatment of PAH, a total of 323 patients were treated with tadalafil at doses ranging from 2.5 mg to 40 mg once daily and 82 patients were treated with placebo. The duration of treatment was 16 weeks. The overall frequency of discontinuation due to adverse events was low (tadalafil 11%, placebo 16%). Three hundred and fifty seven (357) patients who completed the pivotal study entered a long-term extension study. Doses studied were 20 mg and 40 mg once daily.

Tabulated list of adverse reactions

The table below lists the adverse reactions reported during the placebo-controlled clinical study in patients with PAH treated with tadalafil. Also included in the table are some adverse reactions which have been reported in clinical studies and/or post marketing with tadalafil in the treatment of male erectile dysfunction. These events have either been assigned a frequency of "Not known," as the frequency in PAH patients cannot be estimated from the available data or assigned a frequency based on the clinical study data from the pivotal placebo-controlled study of tadalafil.

Frequency estimate: Very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1\,000$ to $< 1/100$), rare ($\geq 1/10\,000$ to $< 1/1\,000$), very rare ($< 1/10\,000$) and not known (cannot be estimated from the available data).

Very common	Common	Uncommon	Rare	Not known¹
<i>Immune system disorders</i>				
	Hypersensitivity reactions ⁵			Angioedema
<i>Nervous system disorders</i>				
Headache ⁶	Syncope, Migraine ⁵	Seizures ⁵ , Transient amnesia ⁵		Stroke ² (including haemorrhagic events)

Very common	Common	Uncommon	Rare	Not known ¹
<i>Eye disorders</i>				
	Blurred vision			Non-arteritic anterior ischemic optic neuropathy (NAION), Retinal vascular occlusion, Visual field defect
<i>Ear and labyrinth disorders</i>				
		Tinnitus		Sudden hearing loss
<i>Cardiac disorders</i>				
	Palpitations ^{2, 5}	Sudden cardiac death ^{2, 5} , Tachycardia ^{2, 5}		Unstable angina pectoris, Ventricular arrhythmia, Myocardial Infarction ²
<i>Vascular disorders</i>				
Flushing	Hypotension	Hypertension		
<i>Respiratory, thoracic and mediastinal disorders</i>				
Nasopharyngitis (including nasal congestion, sinus congestion and rhinitis)	Epistaxis			
<i>Gastrointestinal disorders</i>				
Nausea, Dyspepsia (including abdominal pain/discomfort ³)	Vomiting, Gastroesophageal reflux			
<i>Skin and subcutaneous tissue disorders</i>				
	Rash	Urticaria ⁵ , Hyperhidrosis (sweating) ⁵		Stevens-Johnson Syndrome, Exfoliative dermatitis
<i>Musculoskeletal and connective tissue disorders</i>				
Myalgia, Back pain, Pain in extremity (including limb discomfort)				
<i>Renal and urinary disorders</i>				
		Haematuria		
<i>Reproductive system and breast disorders</i>				
	Increased uterine bleeding ⁴	Priapism ⁵ , Penile haemorrhage, Haemospermia		Prolonged erections
<i>General disorders and administration site conditions</i>				
	Facial oedema, Chest pain ²			

¹ Events not reported in registration studies and cannot be estimated from the available data. The adverse reactions have been included in the table as a result of postmarketing or clinical study data from the use of tadalafil in the treatment of erectile dysfunction.

- ² Most of the patients in whom these events have been reported had pre-existing cardiovascular risk factors.
- ³ Actual MedDRA terms included are abdominal discomfort, abdominal pain, abdominal pain lower, abdominal pain upper, and stomach discomfort.
- ⁴ Clinical non-MedDRA term to include reports of abnormal/excessive menstrual bleeding conditions such as menorrhagia, metrorrhagia, menometrorrhagia, or vaginal hemorrhage.
- ⁵ The adverse reactions have been included in the table as a result of postmarketing or clinical study data from the use of tadalafil in the treatment of erectile dysfunction; and in addition, the frequency estimates are based on only 1 or 2 patients experiencing the adverse reaction in the pivotal placebo-controlled study of tadalafil.
- ⁶ Headache was the most commonly reported adverse reaction. Headache may occur at the beginning of therapy; and decreases over time even if treatment is continued.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [the national reporting system](#) listed in [Appendix V](#).

4.9 Overdose

Single doses of up to 500 mg have been given to healthy subjects, and multiple daily doses up to 100 mg have been given to patients with erectile dysfunction. Adverse reactions were similar to those seen at lower doses.

In cases of overdose, standard supportive measures should be adopted as required. Haemodialysis contributes negligibly to tadalafil elimination.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Urologicals, drugs used in erectile dysfunction, ATC code: G04BE08.

Mechanism of action

Tadalafil is a potent and selective inhibitor of phosphodiesterase type 5 (PDE5), the enzyme responsible for the degradation of cyclic guanosine monophosphate (cGMP). Pulmonary arterial hypertension is associated with impaired release of nitric oxide by the vascular endothelium and consequent reduction of cGMP concentrations within the pulmonary vascular smooth muscle. PDE5 is the predominant phosphodiesterase in the pulmonary vasculature. Inhibition of PDE5 by tadalafil increases the concentrations of cGMP resulting in relaxation of the pulmonary vascular smooth muscle cell and vasodilation of the pulmonary vascular bed.

Pharmacodynamic effects

Studies *in vitro* have shown that tadalafil is a selective inhibitor of PDE5. PDE5 is an enzyme found in corpus cavernosum smooth muscle, vascular and visceral smooth muscle, skeletal muscle, platelets, kidney, lung, and cerebellum. The effect of tadalafil is more potent on PDE5 than on other phosphodiesterases. Tadalafil is > 10,000-fold more potent for PDE5 than for PDE1, PDE2, and PDE4, enzymes which are found in the heart, brain, blood vessels, liver, and other organs. Tadalafil is > 10,000-fold more potent for PDE5 than for PDE3, an enzyme found in the heart and blood vessels. This selectivity for PDE5 over PDE3 is important because PDE3 is an enzyme involved in cardiac contractility. Additionally, tadalafil is approximately 700-fold more potent for PDE5 than for PDE6, an enzyme which is found in the retina and is responsible for phototransduction. Tadalafil is also > 10,000-fold more potent for PDE5 than for PDE7 through PDE10.

Clinical efficacy and safety

Efficacy in patients with pulmonary arterial hypertension (PAH)

A randomised, double-blind, placebo-controlled study was conducted in 405 patients with pulmonary arterial hypertension. Allowed background therapy included bosentan (stable maintenance dose up to 125 mg twice daily) and chronic anticoagulation, digoxin, diuretics and oxygen. More than half (53.3%) of the patients in the study were receiving concomitant bosentan therapy.

Patients were randomised to one of five treatment groups (tadalafil 2.5 mg, 10 mg, 20 mg, 40 mg, or placebo). Patients were at least 12 years of age and had a diagnosis of PAH that was idiopathic, related to collagen disease, related to anorexigen use, related to human immunodeficiency virus (HIV) infection, associated with an atrial-septal defect, or associated with surgical repair of at least 1 year in duration of a congenital systemic-to-pulmonary shunt (for example, ventricular septal defect, patent ductus arteriosus). The mean age of all patients was 54 years (range 14 to 90 years) with the majority of patients being Caucasian (80.5%) and female (78.3%). Pulmonary arterial hypertension (PAH) etiologies were predominantly idiopathic PAH (61.0%) and related to collagen vascular disease (23.5%). The majority of patients had a World Health Organization (WHO) Functional Class III (65.2%) or II (32.1%). The mean baseline 6-minute-walk-distance (6MWD) was 343.6 meters.

The primary efficacy endpoint was the change from baseline at week 16 in 6-minute walk distance (6MWD). Only tadalafil 40 mg achieved the protocol defined level of significance with a placebo-adjusted median increase in 6MWD of 26 metres ($p = 0.0004$; 95% CI: 9.5, 44.0; Pre-specified Hodges-Lehman method) (mean 33 metres, 95% CI: 15.2, 50.3). The improvement in walk distance was apparent from 8 weeks of treatment. Significant improvement ($p < 0.01$) in the 6MWD was demonstrated at week 12 when the patients were asked to delay taking study medicinal product in order to reflect trough active substance concentration. Results were generally consistent in subgroups according to age, gender, PAH aetiology and baseline WHO functional class and 6MWD. The placebo-adjusted median increase in 6MWD was 17 metres ($p = 0.09$; 95% CI: -7.1, 43.0; Pre-specified Hodges-Lehman method) (mean 23 metres, 95% CI: -2.4, 47.8) in those patients who received tadalafil 40 mg in addition to their concomitant bosentan ($n = 39$), and was 39 metres ($p < 0.01$, 95% CI: 13.0, 66.0; Pre-specified Hodges-Lehman method) (mean 44 metres, 95% CI: 19.7, 69.0) in those patients who received tadalafil 40 mg alone ($n = 37$).

The proportion of patients with improvement in WHO functional class by week 16 was similar in the tadalafil 40 mg and placebo groups (23% vs. 21%). The incidence of clinical worsening by week 16 in patients treated with tadalafil 40 mg (5%; 4 of 79 patients) was less than placebo (16%; 13 of 82 patients). Changes in the Borg dyspnoea score were small and non-significant with both placebo and tadalafil 40 mg.

Additionally, improvements compared to placebo were observed with tadalafil 40 mg in the physical functioning, role-physical, bodily pain, general health, vitality and social functioning domains of the SF-36. No improvements were observed in the role emotional and mental health domains of the SF-36. Improvements compared to placebo were observed with tadalafil 40 mg in the EuroQol (EQ-5D) US and UK index scores comprising mobility, self-care, usual activities, pain/discomfort, anxiety/depression components, and in the visual analogue scale (VAS).

Cardiopulmonary hemodynamics was performed in 93 patients. Tadalafil 40 mg increased cardiac output (0.6 L/min) and reduced pulmonary artery pressures (-4.3 mm Hg) and pulmonary vascular resistance (-209 dyn.s/cm⁵) compared to baseline ($p < 0.05$). However, post hoc analyses demonstrated that changes from baseline in cardiopulmonary hemodynamic parameters for the tadalafil 40 mg treatment group were not significantly different compared to placebo.

Long-term treatment

357 patients from the placebo-controlled study entered a long-term extension study. Of these, 311 patients had been treated with tadalafil for at least 6 months and 293 for 1 year (median exposure 365 days; range 2 days to 415 days). For those patients for which there are data, the survival rate at 1

year is 96.4%. Additionally, 6 minute walk distance and WHO functional class status appeared to be stable in those treated with tadalafil for 1 year.

Tadalafil 20 mg administered to healthy subjects produced no significant difference compared to placebo in supine systolic and diastolic blood pressure (mean maximal decrease of 1.6/0.8 mm Hg, respectively), in standing systolic and diastolic blood pressure (mean maximal decrease of 0.2/4.6 mm Hg, respectively), and no significant change in heart rate.

In a study to assess the effects of tadalafil on vision, no impairment of colour discrimination (blue/green) was detected using the Farnsworth-Munsell 100-hue test. This finding is consistent with the low affinity of tadalafil for PDE6 compared to PDE5. Across all clinical studies, reports of changes in colour vision were rare (< 0.1%).

Three studies were conducted in men to assess the potential effect on spermatogenesis of tadalafil 10 mg (one 6-month study) and 20 mg (one 6-month and one 9-month study) administered daily. In two of these studies decreases were observed in sperm count and concentration related to tadalafil treatment of unlikely clinical relevance. These effects were not associated with changes in other parameters such as motility, morphology and FSH.

Paediatric population

A single study has been performed in paediatric patients with Duchenne Muscular Dystrophy (DMD) in which no evidence of efficacy was seen. The randomised, double-blind, placebo-controlled, parallel, 3-arm study of tadalafil was conducted in 331 boys aged 7-14 years with DMD receiving concurrent corticosteroid therapy. The study included a 48-week double-blind period where patients were randomised to tadalafil 0.3 mg/kg, tadalafil 0.6 mg/kg, or placebo daily. Tadalafil did not show efficacy in slowing the decline in ambulation as measured by the primary 6 minute walk distance (6MWD) endpoint: least squares (LS) mean change in 6MWD at 48 weeks was -51.0 meters (m) in the placebo group, compared with -64.7 m in the tadalafil 0.3 mg/kg group ($p = 0.307$) and -59.1 m in the tadalafil 0.6 mg/kg group ($p = 0.538$). In addition, there was no evidence of efficacy from any of the secondary analyses performed in this study. The overall safety results from this study were generally consistent with the known safety profile of tadalafil and with adverse events (AEs) expected in a paediatric DMD population receiving corticosteroids.

The European Medicines Agency has deferred the obligation to submit the results of studies with tadalafil in one or more subsets of the paediatric population in the treatment of pulmonary arterial hypertension (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

Absorption

Tadalafil is readily absorbed after oral administration and the mean maximum observed plasma concentration (C_{max}) is achieved at a median time of 4 hours after dosing. Absolute bioavailability of tadalafil following oral dosing has not been determined.

The rate and extent of absorption of tadalafil are not influenced by food, thus tadalafil may be taken with or without food. The time of dosing (morning versus evening after a single 10 mg administration) had no clinically relevant effects on the rate and extent of absorption.

Distribution

The mean volume of distribution is approximately 77 l at steady state, indicating that tadalafil is distributed into tissues. At therapeutic concentrations, 94% of tadalafil in plasma is bound to proteins. Protein binding is not affected by impaired renal function.

Less than 0.0005% of the administered dose appeared in the semen of healthy subjects.

Biotransformation

Tadalafil is predominantly metabolised by the cytochrome P450 (CYP) 3A4 isoform. The major circulating metabolite is the methylcatechol glucuronide. This metabolite is at least 13,000-fold less

potent than tadalafil for PDE5. Consequently, it is not expected to be clinically active at observed metabolite concentrations.

Elimination

The mean oral clearance for tadalafil is 3.4 l/h at steady state and the mean terminal half-life is 16 hours in healthy subjects. Tadalafil is excreted predominantly as inactive metabolites, mainly in the faeces (approximately 61% of the dose) and to a lesser extent in the urine (approximately 36% of the dose).

Linearity/non-linearity

Over a dose range of 2.5 to 20 mg, tadalafil exposure (AUC) increases proportionally with dose in healthy subjects. Between 20 mg to 40 mg, a less than proportional increase in exposure is observed.

During tadalafil 20 mg and 40 mg once daily dosing, steady-state plasma concentrations are attained within 5 days, and exposure is approximately 1.5-fold of that after a single dose.

Population pharmacokinetics

In patients with pulmonary hypertension not receiving concomitant bosentan, the average tadalafil exposure at steady-state following 40 mg was 26% higher when compared to those of healthy volunteers. There were no clinically relevant differences in C_{max} compared to healthy volunteers. The results suggest a lower clearance of tadalafil in patients with pulmonary hypertension compared to healthy volunteers.

Other special populations

Elderly

Healthy elderly subjects (65 years or over), had a lower oral clearance of tadalafil, resulting in 25% higher exposure (AUC) relative to healthy subjects aged 19 to 45 years after a 10 mg dose. This effect of age is not clinically significant and does not warrant a dose adjustment.

Renal impairment

In clinical pharmacology studies using single-dose tadalafil (5 to 20 mg), tadalafil exposure (AUC) approximately doubled in subjects with mild (creatinine clearance 51 to 80 ml/min) or moderate (creatinine clearance 31 to 50 ml/min) renal impairment and in subjects with end-stage renal disease on dialysis. In haemodialysis patients, C_{max} was 41% higher than that observed in healthy subjects.

Haemodialysis contributes negligibly to tadalafil elimination.

Due to increased tadalafil exposure (AUC), limited clinical experience, and the lack of ability to influence clearance by dialysis, tadalafil is not recommended in patients with severe renal impairment.

Hepatic impairment

Tadalafil exposure (AUC) in subjects with mild and moderate hepatic impairment (Child-Pugh Class A and B) is comparable to exposure in healthy subjects when a dose of 10 mg is administered. If tadalafil is prescribed, a careful individual benefit/risk evaluation should be undertaken by the prescribing physician. There are no available data about the administration of doses higher than 10 mg of tadalafil to patients with hepatic impairment.

Patients with severe hepatic cirrhosis (Child-Pugh Class C) have not been studied and therefore dosing of tadalafil in these patients is not recommended.

Patients with diabetes

Tadalafil exposure (AUC) in patients with diabetes was approximately 19% lower than the AUC value for healthy subjects after a 10 mg dose. This difference in exposure does not warrant a dose adjustment.

Race

Pharmacokinetic studies have included subjects and patients from different ethnic groups, and no differences in the typical exposure to tadalafil have been identified. No dose adjustment is warranted.

Gender

In healthy female and male subjects following single and multiple-doses of tadalafil, no clinically relevant differences in exposure were observed. No dose adjustment is warranted.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, and toxicity to reproduction.

There was no evidence of teratogenicity, embryotoxicity or foetotoxicity in rats or mice that received up to 1000 mg/kg/day tadalafil. In a rat prenatal and postnatal development study, the no observed effect dose was 30 mg/kg/day. In the pregnant rat the AUC for calculated free active substance at this dose was approximately 18 times the human AUC at a 20 mg dose.

There was no impairment of fertility in male and female rats. In dogs given tadalafil daily for 6 to 12 months at doses of 25 mg/kg/day (resulting in at least a 3-fold greater exposure [range 3.7 - 18.6] than seen in humans given a single 20 mg dose) and above, there was regression of the seminiferous tubular epithelium that resulted in a decrease in spermatogenesis in some dogs. See also section 5.1.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core:

Povidone (K-25)
Sodium laurilsulfate
Poloxamer 188
Anhydrous lactose
Microcrystalline cellulose (PH 101)
Croscarmellose sodium
Colloidal anhydrous silica
Magnesium stearate

Film-coating:

Lactose monohydrate
Hypromellose (2910/15 mPa·s) (E464)
Titanium dioxide (E171)
Triacetin

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

PVC/PE/PVdC-Alu blisters in carton containing 28 and 56 film-coated tablets.

PVC/PE/PVdC-Alu perforated unit dose blisters in carton containing 28 x 1 and 56 x 1 film-coated tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1153/001
EU/1/16/1153/002
EU/1/16/1153/003
EU/1/16/1153/004

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 09 January 2017
Date of latest renewal: 09 November 2021

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer(s) responsible for batch release

McDermott Laboratories Limited t/a Gerard Laboratories t/a Mylan Dublin
35/36 Baldoyle Industrial Estate
Grange Road, Dublin 13
Ireland

Mylan Hungary Kft
Mylan utca 1
Komárom, 2900
Hungary

Mylan Germany GmbH
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1
Bad Homburg v. d. Hoehe
Hessen, 61352,
Germany

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

• Periodic Safety Update Reports (PSURs)

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

• Risk Management Plan (RMP)

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON****1. NAME OF THE MEDICINAL PRODUCT**

Talmanco 20 mg film-coated tablets
tadalafil

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 20 mg tadalafil.

3. LIST OF EXCIPIENTS

Contains lactose. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

28 film-coated tablets

28 x 1 film-coated tablets

56 film-coated tablets

56 x 1 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER(S)
--

EU/1/16/1153/001
EU/1/16/1153/002
EU/1/16/1153/003
EU/1/16/1153/004

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY
--

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Talmanco 20 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Talmanco 20 mg film-coated tablets
tadalafil

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viatis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

B. PACKAGE LEAFLET

Package leaflet: Information for the patient

Talmanco 20 mg film-coated tablets tadalafil

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Talmanco is and what it is used for
2. What you need to know before you take Talmanco
3. How to take Talmanco
4. Possible side effects
5. How to store Talmanco
6. Contents of the pack and other information

1. What Talmanco is and what it is used for

Talmanco contains the active substance tadalafil.

Talmanco is a treatment for pulmonary arterial hypertension in adults.

It belongs to a group of medicines called phosphodiesterase type 5 (PDE5) inhibitors which work by helping the blood vessels around your lungs relax, improving the flow of blood into your lungs. The result of this is an improved ability to do physical activity.

2. What you need to know before you take Talmanco

Do not take Talmanco:

- if you are allergic to tadalafil or any of the other ingredients of this medicine (listed in section 6).
- if you are taking any form of nitrates such as amyl nitrite, used in the treatment of chest pain. Tadalafil has been shown to increase the effects of these medicines. If you are taking any form of nitrate or are unsure tell your doctor.
- if you have ever had loss of vision – a condition described as “stroke of the eye” (non-arteritic anterior ischemic optic neuropathy - NAION)
- if you have had a heart attack in the last 3 months
- if you have low blood pressure
- if you are taking riociguat. This medicine is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs secondary to blood clots). PDE5 inhibitors, such as Talmanco, have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.

Warnings and precautions

Talk to your doctor or pharmacist before taking this medicine if you have:

- any heart problems other than your pulmonary hypertension
- problems with your blood pressure
- any hereditary eye disease

- an abnormality of red blood cells (sickle cell anaemia)
- cancer of the bone marrow (multiple myeloma)
- cancer of the blood cells (leukaemia)
- any deformation of your penis, or unwanted or persistent erections lasting more than 4 hours
- a serious liver problem
- a serious kidney problem

If you experience sudden decrease or loss of vision, contact your doctor immediately.

Decreased or sudden hearing loss has been noted in some patients taking tadalafil. Although it is not known if the event is directly related to tadalafil, if you experience decreased or sudden hearing loss, contact your doctor immediately.

Children and adolescents

This medicine is not indicated for use by children and adolescents under the age of 18.

Other medicines and Talmanco

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Do NOT take these tablets if you are already taking nitrates.

Some medicines may be affected by Talmanco or they may affect how well Talmanco will work. Tell your doctor or pharmacist if you are already taking:

- bosentan (another treatment for pulmonary arterial hypertension)
- nitrates (for chest pain)
- alpha blockers used to treat high blood pressure or prostate problems
- riociguat
- rifampicin (to treat bacterial infections)
- ketoconazole tablets (to treat fungal infections)
- ritonavir (for HIV treatment)
- tablets for erectile dysfunction (PDE5 inhibitors)

Talmanco with alcohol

Drinking alcohol may temporarily lower your blood pressure. If you have taken or are planning to take Talmanco, avoid excessive drinking (over 5 units of alcohol), since this may increase the risk of dizziness when standing up.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Do not take this medicine when pregnant, unless it is strictly necessary and you have discussed this with your doctor.

Do not breast-feed while taking these tablets as it is not known if the medicine passes into human breast milk. Ask your doctor or pharmacist for advice before taking any medicine while pregnant or breast-feeding.

When dogs were treated there was reduced sperm development in the testes. A reduction in sperm was seen in some men. These effects are unlikely to lead to a lack of fertility.

Driving and using machines

Dizziness has been reported. Check carefully how you react to the medicines before driving or using any machinery.

Talmanco contains lactose and sodium

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially ‘sodium-free’.

3. How to take Talmanco

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Talmanco is supplied as a 20 mg tablet. **The usual dose** is two 20 mg tablets taken once a day. You should take both tablets at the same time, one after the other. If you have a mild or moderate liver or kidney problem, your doctor may advise you to take only one 20 mg tablet per day.

Swallow the tablets whole with a drink of water. The tablets can be taken with or without food.

If you take more Talmanco than you should

If you or anyone else takes more tablets than they should, tell your doctor or go to a hospital immediately, taking the medicine or pack with you. You may experience any of the side effects described in section 4.

If you forget to take Talmanco

Take your dose as soon as you remember if it is within 8 hours of when you should have taken your dose. Do not take a double dose to make up for a forgotten dose.

If you stop taking Talmanco

Do not stop taking your tablets, unless advised otherwise by your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild to moderate in nature.

If you experience any of the following side effects stop using the medicine and seek medical help immediately:

Common (may affect up to 1 in 10 people)

- allergic reactions including skin rashes.
- chest pain - do not use nitrates, but seek immediate medical assistance.

Uncommon (may affect up to 1 in 100 people)

- priapism, a prolonged and possibly painful erection after taking this medicine. If you have such an erection, which lasts continuously for more than 4 hours you should contact a doctor immediately.

Rare (may affect up to 1 in 1,000 people)

- sudden loss of vision.

Other side effects have been reported:

Very Common (may affect more than 1 in 10 people)

- headache.
- flushing, nasal and sinus congestion (blocked nose).
- nausea, indigestion (including abdominal pain or discomfort).
- muscle aches, back pain and pain in the extremity (including limb discomfort).

Common (may affect up to 1 in 10 people)

- blurred vision.
- low blood pressure.
- nosebleed.
- vomiting.
- increased or abnormal uterine bleeding.
- swelling of the face.
- acid reflux.
- migraine.
- irregular heartbeat.
- fainting.

Uncommon (may affect up to 1 in 100 people)

- seizures.
- passing memory loss.
- hives.
- excessive sweating.
- penile bleeding.
- presence of blood in semen and/or urine.
- high blood pressure.
- fast heart rate.
- sudden cardiac death.
- ringing in the ears.

PDE5 inhibitors are also used for the treatment of erectile dysfunction in men. Some side effects have been **rarely** (may affect up to 1 in 1,000 people) reported:

- Partial, temporary or permanent decrease or loss of vision in one or both eyes and serious allergic reaction which causes swelling of the face or throat. Sudden decrease or loss of hearing has also been reported.

Some side effects have been reported in men taking tadalafil for the treatment of erectile dysfunction. These events were not seen in clinical studies for pulmonary arterial hypertension and therefore frequency is **not known** (frequency cannot be estimated from the available data):

- swelling of the eyelids, eye pain, red eyes, heart attack and stroke.

Most but not all of those men reporting fast heart rate, irregular heartbeat, heart attack, stroke and sudden cardiac death had known heart problems before taking tadalafil. It is not possible to determine whether these events were related directly to tadalafil.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Talmanco

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after “EXP”. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Talmanco contains

- The active substance is tadalafil. Each tablet contains 20 mg of tadalafil.
- The other ingredients in the tablet core are povidone, sodium laurilsulfate, poloxamer 188, anhydrous lactose (see section 2 'Talmanco contains lactose and sodium'), microcrystalline cellulose (PH 101), croscarmellose sodium, colloidal anhydrous silica and magnesium stearate. The film-coating contains lactose monohydrate (see section 2 'Talmanco contains lactose and sodium'), hypromellose (E464), titanium dioxide (E171) and triacetin.

What Talmanco looks like and contents of the pack

This medicine is a white, film-coated, round, biconvex, bevelled edge tablet marked with 'M' on one side of the tablet and 'TA20' on the other side.

This medicine is available in blister packs containing 28 or 56 tablets and in perforated unit dose blister packs containing 28 x 1 or 56 x 1 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

Manufacturers

McDermott Laboratories Limited t/a Gerard Laboratories t/a Mylan Dublin
35/36 Baldoyle Industrial Estate
Grange Road, Dublin 13
Ireland

Mylan Hungary Kft
Mylan utca 1
Komárom, 2900
Hungary

Mylan Germany GmbH
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1
Bad Homburg v. d. Hoehe
Hessen, 61352,
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien
Mylan bvba/sprl
Tél/Tel: +32(0)2 658 61 00

България
Майлан ЕООД
Тел.: +359 2 44 55400

Česká republika
Viatis CZ s.r.o.
Tel: +420 222 004 400

Danmark
Viatis ApS
Tlf: +45 28 11 69 32

Deutschland
Viatis Healthcare GmbH
Tel: +49 800 0700 800

Eesti
BGP Products Switzerland GmbH Eesti
filiaal
Tel: +372 6363 052

Ελλάδα
Generics Pharma Hellas EΠE
Τηλ: +30 210 993 6410

España
Viatis Pharmaceuticals, S.L.U.
Tel: +34 900 102 712

France
Substipharm
Tél: +33 1 43181300

Hrvatska
Mylan Hrvatska d.o.o.
Tel: +385 1 23 50 599

Ireland
Mylan Ireland Limited
Tel: +353 1 8711600

Ísland
Icepharma hf
Sími: +354 540 8000

Italia
Mylan Italia S.r.l
Tel: +39 02 612 46921

Lietuva
Mylan Healthcare UAB
Tel: +370 5 205 1288

Luxembourg/Luxemburg
Mylan bvba/sprl
Tel: +32 (0)2 658 61 00
(Belgique/Belgien)

Magyarország
Mylan EPD Kft
Tel: +36 1 465 2100

Malta
V.J. Salomone Pharma Ltd
Tel: +356 21 22 01 74

Nederland
Mylan BV
Tel: +31 (0)20 426 3300

Norge
Viatis AS
Tlf: +47 66 75 33 00

Österreich
Arcana Arzneimittel GmbH
Tel: +43 1 416 2418

Polska
Mylan Healthcare Sp. z.o.o.
Tel: +48 22 546 64 00

Portugal
Mylan, Lda.
Tel: +351 21 412 72 56

România
BGP Products SRL
Tel: +40 372 579 000

Slovenija
Mylan Healthcare d.o.o.
Tel: +386 1 236 31 80

Slovenská republika
Viatis Slovakia s.r.o.
Tel: +421 2 32 199 100

Suomi/Finland
Viatis Oy
Puh/Tel: +358 20 720 9555

Κύπρος

Varnavas Hadjipanayis Ltd

Τηλ: +357 222 077 00

Sverige

Viatris AB

Tel: +46 (0) 8 630 19 00

Latvija

Mylan Healthcare SIA

Tel: +371 676 055 80

United Kingdom (Northern Ireland)

Mylan IRE Healthcare Limited

Tel: +353 18711600

This leaflet was last revised in {MM/YYYY}.

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>.