

Question 1

From largest to smallest, the asymptotic size of the given functions are (with lg taken to represent \log_2 and lg^{1000} taken to mean performing the lg operation one-thousand times):

$$lg(n^{1000}) \tag{1}$$

$$(lgn)^n \tag{2}$$

$$n^{lgn} \tag{3}$$

$$n^{lg lgn} \tag{4}$$

$$(lgn)^{(lgn)} \tag{5}$$

$$n^2 \tag{6}$$

$$nlgn \tag{7}$$

$$n \tag{8}$$

$$lgn \tag{9}$$

$$n^{(1/lgn)} \tag{10}$$

$$(1 + 0.001)^n \tag{11}$$

$$lg_{1000} n \tag{12}$$

$$lg^{1000} n \tag{13}$$

$$1 \tag{14}$$

Question 2

The answers to this question rely on the discussion of the master method in chapter 4 of the CLRS textbook.

2.1 $T(n) = 2T(n/3) + 1$, so $a = 2, b = 3, f(n) = 1$. Using the master method, this is case 1: $f(n) = O(n^{\log_3 2 - \epsilon})$, so $\mathbf{T(n)} = \Theta(\mathbf{n^{\log_3 2}})$.

2.2 $T(n) = 5T(n/4) + n$, so $a = 5, b = 4, f(n) = n$. Again this is case 1 of the master method: $f(n) = O(n^{\log_4 5 - \epsilon})$, so $\mathbf{T(n)} = \Theta(\mathbf{n^{\log_5 4}})$.

2.3 $T(n) = 8T(n/2) + n^3$, so $a = 8, b = 2, f(n) = n^3$. This is case 2 of the master method: $f(n) = \Theta(n^{\log_2 8})$, so $\mathbf{T(n)} = \Theta(\mathbf{n^3 lgn})$

2.4 $T(n) = T(n^{1/2}) + 1$. This recurrence relation cannot be directly translated into the form of the master method, so we use a change of variables $m = lgn$. $S(m) = S(m/2) + 1$, so $a_m = 1, b_m = 2, f(m) = 1$. This is case 3 of the master method, because $S(m) =$

$\Theta(m^{\log_2 1}) = \Theta(lgm)$. Changing back to our original variables, $T(n) = T(2^m) = S(m) = \Theta(lgm) = \Theta(\lg \lg n)$

Question 3

Question 4

Suppose we label the three pegs P_1, P_2 , and P_3 . Without loss of generality, we can assume that initially all n pegs are on P_1 and our goal is to move them to P_3 under the constraints posed in the question. For convenience we also label the $n \geq 1$ discs d_1, \dots, d_n where d_1 is the smallest and d_n is the largest disc. Let $T(n)$ denote the number of steps required for n discs.

Our algorithm must first handle the base case, which is trivial: when $n = 1$, move d_n directly from P_1 to P_3 . For any input n , this is the final step.

In general, we can solve the Towers of Hanoi problem in three steps with recursion:

1. Move all discs d_1, \dots, d_{n-1} from P_1 to P_2
2. Move disc d_n from P_1 to P_3
3. Move discs d_1, \dots, d_{n-1} from P_2 to P_3 .

Notice that by relabeling the discs that step 1 is equivalent to solving the problem at level $T(n-1)$ if our original goal was to move from P_1 to P_2 . Similarly, step 3 is equivalent to $T(n-1)$ if the original goal was to move from P_2 to P_3 . Step 2 will always take exactly 1 operation. This gives us the recurrence relation $\mathbf{T(n) = 2T(n-1) + 1}$.

We can get the total number of moves as a function of n using this fact. To help understand the recurrence relation it is helpful to examine a few cases with small n . We can easily see that in the base case $T(n=1) = 1$. Thus, $T(n=2) = 2(1) + 1 = 3$. Further,

$$\begin{aligned} T(n=3) &= 2(3) + 1 \\ &= 2(2(1)) + 2(1) + 1 \\ &= 2^2 + 2^1 + 2^0 \\ &= 7. \end{aligned}$$

More generally, $T(n)$ can be viewed as a summation series: $T(n) = \sum_{i=0}^{n-1} 2^i = 2^n - 1$. To see this, we use the recurrence relation $T(n) = 2T(n-1) + 1$ and see that at each step we add 2^{n-1} steps (this is also apparent from the result in Method 1 above). In the base case we already have a minimum of $2^1 - 1 = 1$ steps. For $n = 2$ we add $2^{2-1} = 2$ steps for a total of $2 + 1 = 3$ steps. In general, at the n^{th} step we are adding 2^{n-1} steps to a total of $2^{n-1} - 1$ steps from the $(n-1)^{th}$ iteration. Because $2^{n-1} + 2^{n-1} = 2 \times 2^{n-1} = 2^n$, we can simplify the series into the closed form equation $2^n - 1$ steps (see also: Equation A.6 in CLRS).

That is, in general for input of size n the Towers of Hanoi algorithm above requires **$2^n - 1$ operations**.