

# Causal Inference: Presenting Tools to Ascend the Ladder of Causation

Charles Assaad [charles.assaad@inserm.fr](mailto:charles.assaad@inserm.fr)

L’Institut Pierre Louis d’Epidémiologie et de Santé Publique,  
Inserm, Sorbonne Université

June 2025



- 1** Causal hierarchy
- 2** Association
  - Bayesian networks
  - d-separation
  - Prediction
  - Do we need more than associations?
- 3** Intervention
  - Causal DAGs
  - Causal reasoning
- 4** Counterfactuals
  - Structural causal models
  - Counterfactual reasoning
  - Mediation analysis
  - Difference with the potential outcome framework
- 5** Conclusion

# 1

Causal hierarchy

# Some epidemiological questions

## Some epidemiological questions

- Do people who exercise more tend to have lower rates of heart disease?

## Some epidemiological questions

- Do people who exercise more tend to have lower rates of heart disease?
- Can we predict who is at higher risk of developing diabetes using BIG DATA?

## Some epidemiological questions


- Do people who exercise more tend to have lower rates of heart disease?
- Can we predict who is at higher risk of developing diabetes using BIG DATA?
- Does exercising reduce cholesterol?

- Do people who exercise more tend to have lower rates of heart disease?
- Can we predict who is at higher risk of developing diabetes using BIG DATA?
- Does exercising reduce cholesterol?
- What happens to blood pressure if we prescribe a new hypertension drug?


- Do people who exercise more tend to have lower rates of heart disease?
- Can we predict who is at higher risk of developing diabetes using BIG DATA?
- Does exercising reduce cholesterol?
- What happens to blood pressure if we prescribe a new hypertension drug?
- Would COVID-19 outcomes in France have changed with a different policy?

- Do people who exercise more tend to have lower rates of heart disease?
- Can we predict who is at higher risk of developing diabetes using BIG DATA?
- Does exercising reduce cholesterol?
- What happens to blood pressure if we prescribe a new hypertension drug?
- Would COVID-19 outcomes in France have changed with a different policy?
- Would air pollution affect mortality if it didn't affect respiratory diseases?

# Pearl causal hierarchy




# Pearl causal hierarchy



## ■ Associations

- ▶ Questions : What if I see ...?
- ▶ Do people who exercise more tend to have lower rates of heart disease?
- ▶ Can we predict who is at higher risk of developing diabetes using BIG DATA?

# Pearl causal hierarchy




## ■ Interventions

- ▶ Questions: What if I do ...? How?
- ▶ Does exercising reduce cholesterol?
- ▶ What happens to blood pressure if we prescribe a new hypertension drug?

## ■ Associations

- ▶ Questions : What if I see ...?
- ▶ Do people who exercise more tend to have lower rates of heart disease?
- ▶ Can we predict who is at higher risk of developing diabetes using BIG DATA?

# Pearl causal hierarchy



## ■ Counterfactuals

- ▶ Questions: What if I had done ...? Why?
- ▶ Would COVID-19 outcomes in France have changed with a different policy?
- ▶ Would air pollution affect mortality if it didn't affect respiratory diseases?

## ■ Interventions

- ▶ Questions: What if I do ...? How?
- ▶ Does exercising reduce cholesterol?
- ▶ What happens to blood pressure if we prescribe a new hypertension drug?

## ■ Associations

- ▶ Questions : What if I see ...?
- ▶ Do people who exercise more tend to have lower rates of heart disease?
- ▶ Can we predict who is at higher risk of developing diabetes using BIG DATA?

# 2

## Association

Bayesian networks

d-separation

Prediction

Do we need more than associations?

# 2

## Association

Bayesian networks

d-separation

Prediction

Do we need more than associations?


- Sets:  $\mathbb{A} = \{X, Y, Z\}$
- Statistical independence:  $\perp\!\!\!\perp_P$
- Statistical dependence:  $\not\perp\!\!\!\perp_P$
- $P(Y = y | X = x) \equiv P(y | x)$

A graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$  is said to be a **directed graph** iff

- $\mathbb{V}$  is the set of vertices (usually each vertex corresponds to a random variable),
- $\mathbb{E}$  is the set of edges,
- $\forall(X, Y) \in \mathbb{E}$ , there is an arrow pointing from X to Y.

## Directed graphs: basic concepts


Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :



## Directed graphs: basic concepts

Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :

Path:  $D \leftarrow B \rightarrow E \leftarrow C$




## Directed graphs: basic concepts

Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :

Path:  $D \leftarrow B \rightarrow E \leftarrow C$

Directed path:  $A \rightarrow B \rightarrow E \rightarrow F \rightarrow G$




## Directed graphs: basic concepts

Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :

Path:  $D \leftarrow B \rightarrow E \leftarrow C$

Directed path:  $A \rightarrow B \rightarrow E \rightarrow F \rightarrow G$

Parents:  $Pa(E) = \{B, C\}$



# Directed graphs: basic concepts


Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :

Path:  $D \leftarrow B \rightarrow E \leftarrow C$

Directed path:  $A \rightarrow B \rightarrow E \rightarrow F \rightarrow G$

Parents:  $Pa(E) = \{B, C\}$

Ancestors:  $An(E) = \{A, B, C, E\}$



# Directed graphs: basic concepts

Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :


Path:  $D \leftarrow B \rightarrow E \leftarrow C$

Directed path:  $A \rightarrow B \rightarrow E \rightarrow F \rightarrow G$

Parents:  $Pa(E) = \{B, C\}$

Ancestors:  $An(E) = \{A, B, C, E\}$

Descendants:  $De(E) = \{E, F, G\}$



# Directed graphs: basic concepts

Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :

Path:  $D \leftarrow B \rightarrow E \leftarrow C$


Directed path:  $A \rightarrow B \rightarrow E \rightarrow F \rightarrow G$

Parents:  $Pa(E) = \{B, C\}$

Ancestors:  $An(E) = \{A, B, C, E\}$

Descendants:  $De(E) = \{E, F, G\}$

Non-descendants:  $Nd(E) = \{A, B, C, D\}$



# Directed graphs: basic concepts

Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :

Path:  $D \leftarrow B \rightarrow E \leftarrow C$

Directed path:  $A \rightarrow B \rightarrow E \rightarrow F \rightarrow G$


Parents:  $Pa(E) = \{B, C\}$

Ancestors:  $An(E) = \{A, B, C, E\}$

Descendants:  $De(E) = \{E, F, G\}$

Non-descendants:  $Nd(E) = \{A, B, C, D\}$

Ancestral sets: a subset of vertices  $\mathbb{S}$  is ancestral if  $\forall X \in \mathbb{S}, An(X) \subseteq \mathbb{S}$



## Directed graphs: basic concepts

Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :

Path:  $D \leftarrow B \rightarrow E \leftarrow C$

Directed path:  $A \rightarrow B \rightarrow E \rightarrow F \rightarrow G$

Parents:  $Pa(E) = \{B, C\}$


Ancestors:  $An(E) = \{A, B, C, E\}$

Descendants:  $De(E) = \{E, F, G\}$

Non-descendants:  $Nd(E) = \{A, B, C, D\}$

Ancestral sets: a subset of vertices  $\mathbb{S}$  is ancestral if  $\forall X \in \mathbb{S}, An(X) \subseteq \mathbb{S}$

Subgraph  $\mathcal{G}[\mathbb{S}]$ :  $\mathcal{G}[\{B, C, D, F\}]$



# Directed graphs: basic concepts

Consider the following directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ :

**Path:**  $D \leftarrow B \rightarrow E \leftarrow C$

**Directed path:**  $A \rightarrow B \rightarrow E \rightarrow F \rightarrow G$

**Parents:**  $Pa(E) = \{B, C\}$

**Ancestors:**  $An(E) = \{A, B, C, E\}$


**Descendants:**  $De(E) = \{E, F, G\}$

**Non-descendants:**  $Nd(E) = \{A, B, C, D\}$

**Ancestral sets:** a subset of vertices  $\mathbb{S}$  is ancestral if  $\forall X \in \mathbb{S}, An(X) \subseteq \mathbb{S}$

**Subgraph  $\mathcal{G}[\mathbb{S}]$ :**  $\mathcal{G}[\{B, C, D, F\}]$

**Skeleton of  $\mathcal{G}$**



A directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$  is said to be a **directed acyclic graph** (DAG) iff

$$\forall X \in \mathbb{V}, An(X) \cap De(X) = \{X\},$$

i.e., there are no cycle in  $\mathcal{G}$ .

A directed graph  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$  is said to be a **directed acyclic graph** (DAG) iff

$$\forall X \in \mathbb{V}, An(X) \cap De(X) = \{X\},$$

i.e., there are no cycle in  $\mathcal{G}$ .

From now on, we will only deal with DAGs.

## Definition

A distribution  $P(\mathbb{V})$  is **compatible** with a DAG  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$  if

$$P(\mathbb{V}) = \prod_{X \in \mathbb{V}} P(X | Pa(X))$$

A distribution  $P(\mathbb{V})$  is **compatible** with a DAG  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$  if

$$P(\mathbb{V}) = \prod_{X \in \mathbb{V}} P(X | Pa(X))$$

$\Leftrightarrow \forall X \in \mathbb{V}, X \perp\!\!\!\perp_P Nd(X) | Pa(X)$  (**Markov condition**)

## Definition

A distribution  $P(\mathbb{V})$  is **compatible** with a DAG  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$  if

$$P(\mathbb{V}) = \prod_{X \in \mathbb{V}} P(X | Pa(X))$$

$$\Leftrightarrow \forall X \in \mathbb{V}, X \perp\!\!\!\perp_{\mathcal{G}} Nd(X) | Pa(X) \text{ (Markov condition)}$$


A DAG  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$  is a **Bayesian network** iff there exists a joint distribution  $P(\mathbb{V})$  that is compatible with  $\mathcal{G}$ .

## Theorem

If  $P$  is compatible with  $\mathcal{G}$  and  $\mathbb{S} \subseteq \mathbb{V}$  is an ancestral set, then  $P(\mathbb{S})$  is compatible with  $\mathcal{G}[\mathbb{S}]$ .

## Theorem


If  $P$  is compatible with  $\mathcal{G}$  and  $S \subseteq V$  is an ancestral set, then  $P(S)$  is compatible with  $\mathcal{G}[S]$ .




Bayesian network  
compatible with  $P$

## Theorem

If  $P$  is compatible with  $\mathcal{G}$  and  $\mathbb{S} \subseteq \mathbb{V}$  is an ancestral set, then  $P(\mathbb{S})$  is compatible with  $\mathcal{G}[\mathbb{S}]$ .



Bayesian network  
compatible with  $P$



Bayesian network  
compatible with  
 $P(A, B, C, D)$

# 2

## Association

Bayesian networks




d-separation

Prediction

Do we need more than associations?

## Example



$$A \stackrel{?}{\perp\!\!\!\perp}_P D | B$$



# Basic structures



# Basic structures

Fork: contains a confounder





## Basic structures

Fork: contains a confounder





Chain: contains a mediator or an intermediate cause





## Basic structures



Fork: contains a confounder



Chain: contains a mediator or an intermediate cause





V-structure: contains a collider





## Basic structures



Fork: contains a confounder




Chain: contains a mediator or an intermediate cause




V-structure: contains a collider



## Example



## Example




$$A = \begin{cases} \text{Mother carrier} \\ \text{Mother not carrier} \end{cases}$$

$$B = \begin{cases} \text{Father carrier} \\ \text{Father not carrier} \end{cases}$$

$$C = (A \text{ or } B) = \begin{cases} \text{Child carrier} \\ \text{Child not carrier} \end{cases}$$

## Example



$$A = \begin{cases} \text{Mother carrier} \\ \text{Mother not carrier} \end{cases}$$


$$B = \begin{cases} \text{Father carrier} \\ \text{Father not carrier} \end{cases}$$

$$C = (A \text{ or } B) = \begin{cases} \text{Child carrier} \\ \text{Child not carrier} \end{cases}$$


If  $C = \text{Child carrier} \implies$

$$\begin{cases} \text{If } A = \text{Mother not carrier then } B = \text{Father carrier} \\ \text{If } B = \text{Father not carrier then } A = \text{Mother carrier} \end{cases}$$

## Example



## Example




$$A, B \sim U(-1, 1)$$


$$\xi_c \sim N(0, \frac{1}{2})$$

$$C = 2AB + \xi_c$$

## Example



$$A, B \sim U(-1, 1)$$




$$\xi_c \sim N(0, \frac{1}{2})$$


$$C = 2AB + \xi_c$$

$$\text{Corr}(A; B) = 0.002$$

## Example




$$A, B \sim U(-1, 1)$$



$$\text{Corr}(A; B) = 0.002$$

$$\xi_c \sim N(0, \frac{1}{2})$$



$$\text{Corr}(A; B | C > 0.5) = 0.8$$

## Blocked paths

A path is said to be **blocked** by a set of vertices  $\mathbb{Z} \in \mathbb{V}$  if:

- it contains a chain  $A \rightarrow B \rightarrow C$  or a fork  $A \leftarrow B \rightarrow C$  and  $B \in \mathbb{Z}$ ; or
- it contains a collider  $A \rightarrow B \leftarrow C$  such that no descendant of  $B$  is in  $\mathbb{Z}$ .


## Blocked paths

A path is said to be **blocked** by a set of vertices  $\mathbb{Z} \in \mathbb{V}$  if:


- it contains a chain  $A \rightarrow B \rightarrow C$  or a fork  $A \leftarrow B \rightarrow C$  and  $B \in \mathbb{Z}$ ; or
- it contains a collider  $A \rightarrow B \leftarrow C$  such that no descendant of  $B$  is in  $\mathbb{Z}$ .

A path that is not blocked is **active**.

## Example




## Example




Is the path  $\langle D, B, E, C \rangle$  blocked?

## Example




Is the path  $\langle D, B, E, C \rangle$  blocked? Yes

## Example



Is the path  $< D, B, E, C >$  blocked? Yes  
Is the path  $< D, B, E, C >$  blocked by  $E$ ?


## Example



Is the path  $\langle D, B, E, C \rangle$  blocked? Yes  
Is the path  $\langle D, B, E, C \rangle$  blocked by  $E$ ? No

## Blocked path: examples

## Example




Is the path  $\langle D, B, E, C \rangle$  blocked? Yes

Is the path  $\langle D, B, E, C \rangle$  blocked by  $E$ ? No

Is the path  $\langle A, B, E, F, G \rangle$  blocked?

## Blocked path: examples

## Example




Is the path  $< D, B, E, C >$  blocked? Yes

Is the path  $< D, B, E, C >$  blocked by  $E$ ? No

Is the path  $< A, B, E, F, G >$  blocked? No


## Example



- Is the path  $\langle D, B, E, C \rangle$  blocked? Yes
- Is the path  $\langle D, B, E, C \rangle$  blocked by  $E$ ? No
- Is the path  $\langle A, B, E, F, G \rangle$  blocked? No
- Is the path  $\langle A, B, E, F, G \rangle$  blocked by  $E$ ?

## Blocked path: examples

## Example



Is the path  $\langle D, B, E, C \rangle$  blocked? Yes

Is the path  $\langle D, B, E, C \rangle$  blocked by  $E$ ? No

Is the path  $\langle A, B, E, F, G \rangle$  blocked? No

Is the path  $\langle A, B, E, F, G \rangle$  blocked by  $E$ ? Yes

## d-separation

Given disjoint sets  $X, Y, Z \subseteq V$ , we say that  $X$  and  $Y$  are **d-separated** by  $Z$  if every path between a vertex in  $X$  and a vertex in  $Y$  is blocked by  $Z$  and we write  $X \perp\!\!\!\perp_{\mathcal{G}} Y | Z$ .

## d-separation

Given disjoint sets  $X, Y, Z \subseteq V$ , we say that  $X$  and  $Y$  are **d-separated** by  $Z$  if every path between a vertex in  $X$  and a vertex in  $Y$  is blocked by  $Z$  and we write  $X \perp\!\!\!\perp_Z Y | Z$ .

If one path is not blocked, we say that  $X$  and  $Y$  are **d-connected** given  $Z$  and we write  $X \not\perp\!\!\!\perp_Z Y | Z$ .

Given disjoint sets  $\mathbb{X}, \mathbb{Y}, \mathbb{Z} \subseteq \mathbb{V}$ , we say that  $\mathbb{X}$  and  $\mathbb{Y}$  are **d-separated** by  $\mathbb{Z}$  if every path between a vertex in  $\mathbb{X}$  and a vertex in  $\mathbb{Y}$  is blocked by  $\mathbb{Z}$  and we write  $\mathbb{X} \perp\!\!\!\perp_{\mathcal{G}} \mathbb{Y} | \mathbb{Z}$ .

If one path is not blocked, we say that  $\mathbb{X}$  and  $\mathbb{Y}$  are **d-connected** given  $\mathbb{Z}$  and we write  $\mathbb{X} \not\perp\!\!\!\perp_{\mathcal{G}} \mathbb{Y} | \mathbb{Z}$ .

### Theorem

$$\mathbb{X} \perp\!\!\!\perp_{\mathcal{G}} \mathbb{Y} | \mathbb{Z} \Rightarrow \mathbb{X} \perp\!\!\!\perp_{\mathcal{P}} \mathbb{Y} | \mathbb{Z}$$

## d-separation

Given disjoint sets  $X, Y, Z \subseteq V$ , we say that  $X$  and  $Y$  are **d-separated** by  $Z$  if every path between a vertex in  $X$  and a vertex in  $Y$  is blocked by  $Z$  and we write  $X \perp\!\!\!\perp_G Y | Z$ .

If one path is not blocked, we say that  $X$  and  $Y$  are **d-connected** given  $Z$  and we write  $X \not\perp\!\!\!\perp_G Y | Z$ .

### Theorem

$X \perp\!\!\!\perp_G Y | Z \Rightarrow X \perp\!\!\!\perp_P Y | Z$   
 but  $X \not\perp\!\!\!\perp_G Y | Z \not\Rightarrow X \not\perp\!\!\!\perp_P Y | Z$

## d-separation

Given disjoint sets  $X, Y, Z \subseteq V$ , we say that  $X$  and  $Y$  are **d-separated** by  $Z$  if every path between a vertex in  $X$  and a vertex in  $Y$  is blocked by  $Z$  and we write  $X \perp\!\!\!\perp_Z Y | Z$ .

If one path is not blocked, we say that  $X$  and  $Y$  are **d-connected** given  $Z$  and we write  $X \not\perp\!\!\!\perp_Z Y | Z$ .

### Theorem




$$X \perp\!\!\!\perp_G Y | Z \Rightarrow X \perp\!\!\!\perp_P Y | Z$$

but  $X \not\perp\!\!\!\perp_G Y | Z \not\Rightarrow X \not\perp\!\!\!\perp_P Y | Z$  instead

$$X \not\perp\!\!\!\perp_G Y | Z \stackrel{?}{\Rightarrow} X \perp\!\!\!\perp_P Y | Z$$




## Example

Is  $A \perp\!\!\!\perp_P D | B?$






## Example

Is  $A \perp\!\!\!\perp_P D | B?$




## Example

Is  $A \perp\!\!\!\perp_P D | B?$





## Example

Is  $A \perp\!\!\!\perp_P D | B?$




$\langle A, C, D \rangle$  is not  
blocked  
?  
 $\implies A \perp\!\!\!\perp_P D | B$





## Example

Is  $A \perp\!\!\!\perp_P D | B?$




$\langle A, C, D \rangle$  is not  
blocked  
?  
 $\implies A \perp\!\!\!\perp_P D | B$



## Example


Is  $A \perp\!\!\!\perp_D D | B$ ?



$\langle A, C, D \rangle$  is not  
blocked


?

$\implies A \perp\!\!\!\perp_D D | B$




All paths are  
blocked

$\implies A \perp\!\!\!\perp_D D | B$



## Example


Is  $A \perp\!\!\!\perp_D D | B$ ?



$\langle A, C, D \rangle$  is not  
blocked


?

$\implies A \perp\!\!\!\perp_D D | B$




All paths are  
blocked

$\implies A \perp\!\!\!\perp_D D | B$




## Example

Is  $A \perp\!\!\!\perp_D D | B?$




$\langle A, C, D \rangle$  is not  
blocked  
?

$\implies A \perp\!\!\!\perp_D D | B$




All paths are  
blocked  
 $\implies A \perp\!\!\!\perp_D D | B$




## Example

Is  $A \perp\!\!\!\perp_D D | B?$




$\langle A, C, D \rangle$  is not  
blocked  
?

$\implies A \perp\!\!\!\perp_D D | B$




All paths are  
blocked  
 $\implies A \perp\!\!\!\perp_D D | B$




## Example

Is  $A \perp\!\!\!\perp_D D | B?$




$\langle A, C, D \rangle$  is not  
blocked  
?

$\implies A \perp\!\!\!\perp_D D | B$




All paths are  
blocked  
 $\implies A \perp\!\!\!\perp_D D | B$




## Example

Is  $A \perp\!\!\!\perp_D D | B$ ?




$\langle A, C, D \rangle$  is not  
blocked  
?

$\implies A \perp\!\!\!\perp_D D | B$




All paths are  
blocked  
 $\implies A \perp\!\!\!\perp_D D | B$



## Example


Is  $A \perp\!\!\!\perp_D D | B?$



$\langle A, C, D \rangle$  is not  
blocked


?

$\implies A \perp\!\!\!\perp_D D | B$




All paths are  
blocked

$\implies A \perp\!\!\!\perp_D D | B$




## Example

Is  $A \perp\!\!\!\perp_P D | B$ ?




$\langle A, C, D \rangle$  is not  
blocked  
?

$\implies A \perp\!\!\!\perp_P D | B$



All paths are  
blocked  
?  
 $\implies A \perp\!\!\!\perp_P D | B$



$\langle A, B, I, G, F, D \rangle$   
is not blocked  
?  
 $\implies A \perp\!\!\!\perp_P D | B$

# 2

## Association

Bayesian networks

d-separation

Prediction

Do we need more than associations?

Bayesian networks can be used to detect relevant variables that contains all the useful information for prediction!

Bayesian networks can be used to detect relevant variables that contains all the useful information for prediciton! (Assuming the distribution does not change)

Bayesian networks can be used to detect relevant variables that contains all the useful information for prediciton! (Assuming the distribution does not change)


Relevant variables for predicting  $B$ :

- Parents of  $B$ ;
- Children of  $B$ ;
- Other parents of the children of  $B$ .

Bayesian networks can be used to detect relevant variables that contains all the useful information for prediciton! (Assuming the distribution does not change)

Relevant variables for predicting  $B$ :


- Parents of  $B$ ;
- Children of  $B$ ;
- Other parents of the children of  $B$ .



Bayesian networks can be used to detect relevant variables that contains all the useful information for prediciton! (Assuming the distribution does not change)

Relevant variables for predicting  $B$ :

- Parents of  $B$ ;
- Children of  $B$ ;
- Other parents of the children of  $B$ .



# 2

## Association


Bayesian networks

d-separation

Prediction

Do we need more than associations?





## Distribution shifts



---

P. Cui and T. Zhang. Causal Inference and Stable Learning. Tutoriel at ICML, 2019.


## Distribution shifts

**Train****Performance: 99%****Validation****Performance: 99%****New image****It's a fish :)**

P. Cui and T. Zhang. Causal Inference and Stable Learning. Tutoriel at ICML, 2019.


## Simpson paradox [6]

In a study, we measure weekly exercise and cholesterol levels for various age groups.





## Exercise 1


Suppose this DAG is a bayesian network compatible with  $P$ :



Do the following subgraphs represent Bayesian networks that are compatible with their respective distributions?




## Exercise 2



- $B \perp\!\!\!\perp_P G | F?$
- $A \perp\!\!\!\perp_P F | E?$
- $B \perp\!\!\!\perp_P E | A, C, F?$

## Exercise 3

What are the relevant variables for predicting  $B$ ?



# 3

Intervention

Causal DAGs

Causal reasoning



# 3

Intervention


Causal DAGs

Causal reasoning


# Bayesian networks vs causal DAGs



# Bayesian networks vs causal DAGs




Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network




Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network

# Bayesian networks vs causal DAGs




Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network  
 Not a causal DAG




Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network  
 Causal DAG

# Bayesian networks vs causal DAGs




Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network  
 Not a causal DAG




Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network  
 Causal DAG

It is impossible to determine, without additional assumptions, which of the two DAGs is causal based solely on the observed distribution!

# Bayesian networks vs causal DAGs



Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network  
 Not a causal DAG




Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network  
 Causal DAG

It is impossible to determine, without additional assumptions, which of the two DAGs is causal based solely on the observed distribution!


Oracle for conditional  
independence

Oracle for intervention

# Bayesian networks vs causal DAGs



Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network  
 Not a causal DAG



Bad breath  $\perp\!\!\!\perp_P$  Coma |  
 Diabetic ketoacidosis  
 Bayesian network  
 Causal DAG

It is impossible to determine, without additional assumptions, which of the two DAGs is causal based solely on the observed distribution!

Oracle for conditional  
independence

Oracle for intervention

The  $do()$  operator encodes (structural) interventions.

# Conditioning vs Intervening


Population



## Conditioning vs Intervening

Population

Sub-populations




# Conditioning vs Intervening



Population



Sub-populations



Conditioning




# Conditioning vs Intervening



Population





Sub-populations



Conditioning



Intervening



Truncated factorization [4]: If we intervene on a subset  $\mathbb{S} \subset \mathbb{V}$ , then

$$P(v_1, \dots, v_d \mid do(s)) = \prod_{v_i \notin \mathbb{S}} P(v_i \mid Pa(v_i))$$

Truncated factorization [4]: If we intervene on a subset  $\mathbb{S} \subset \mathbb{V}$ , then

$$P(v_1, \dots, v_d \mid do(s)) = \prod_{v_i \notin \mathbb{S}} P(v_i \mid Pa(v_i))$$

Also known as G-computation formula [2] or Manipulation theorem [10].

Truncated factorization [4]: If we intervene on a subset  $\mathbb{S} \subset \mathbb{V}$ , then

$$P(v_1, \dots, v_d \mid do(s)) = \prod_{v_i \notin \mathbb{S}} P(v_i \mid Pa(v_i))$$

Also known as G-computation formula [2] or Manipulation theorem [10].

Let  $P(\mathbb{V})$  be a probability distribution and let  $P_*$  denote the set of all interventional distributions  $P(\mathbb{V} \mid do(s))$ . A bayesian network  $\mathcal{G}$  is said to be a **causal DAG** compatible with  $P_*$  iff  $\mathcal{G}$  and  $P_*$  satisfy the truncated factorization.

In causal DAGs, the information lies in the absence of edges and the direction of arrows, rather than merely in the presence of edges.

In causal DAGs, the information lies in the absence of edges and the direction of arrows, rather than merely in the presence of edges.

- Think carrefully about the orientation

In causal DAGs, the information lies in the absence of edges and the direction of arrows, rather than merely in the presence of edges.

- Think carrefully about the orientation
- When you are not sure if you need to add or not an edge (for example  $Z \rightarrow A$ ) to the graph, ADD IT! (as long as you keep the graph acyclic)


In causal DAGs, the information lies in the absence of edges and the direction of arrows, rather than merely in the presence of edges.

- Think carrefully about the orientation
- When you are not sure if you need to add or not an edge (for example  $Z \rightarrow A$ ) to the graph, ADD IT! (as long as you keep the graph acyclic)

If you cannot keep the graph acyclic, do not worry, there exists new tools for cyclic graphs.


But in this presentation we will focus only on acyclic graphs.

## Example



Is this DAG causal?

## Example




Is this DAG causal?

No

## DAG Vs DAG causal: exemples

## Example



Is this DAG causal?


No

Gene GNPDA2

Hypertension —————→ Renal function

Is this DAG causal?

## Example



Is this DAG causal?

No

Gene GNPDA2

Hypertension --&gt; Renal function

Is this DAG causal?

Yes

# 3

Intervention

Causal DAGs

Causal reasoning

A quantity  $Q$  is said to be **identifiable** if it is uniquely computable from a positive distribution  $P(\mathbb{V})$ .

A quantity  $Q$  is said to be **identifiable** if it is uniquely computable from a positive distribution  $P(\mathbb{V})$ .


In the following, we will assume that all distributions are positive.

The total effect of  $A$  on  $Y$

$$= \mathbb{E}(Y \mid do(A = a)) - \mathbb{E}(Y \mid do(A = a'))$$


The total effect of  $A$  on  $Y$

$$= \mathbb{E}(Y \mid do(A = a)) - \mathbb{E}(Y \mid do(A = a'))$$



The total effect of  $A$  on  $Y$


$$= \mathbb{E}(Y \mid do(A = a)) - \mathbb{E}(Y \mid do(A = a'))$$



The total effect is identifiable iff  $P(Y \mid do(A))$  is identifiable.

The total effect of  $A$  on  $Y$

$$= \mathbb{E}(Y \mid do(A = a)) - \mathbb{E}(Y \mid do(A = a'))$$




The total effect is identifiable iff  $P(Y \mid do(A))$  is identifiable.

Causal reasoning involves utilizing a causal DAG to determine whether  $P(Y \mid do(A))$  is identifiable.

The total effect of  $A$  on  $Y$

$$= \mathbb{E}(Y \mid do(A = a)) - \mathbb{E}(Y \mid do(A = a'))$$




The total effect is identifiable iff  $P(Y \mid do(A))$  is identifiable.

Causal reasoning involves utilizing a causal DAG to determine whether  $P(Y \mid do(A))$  is identifiable.

In what follows, we will use the terms total effect and causal effect interchangeably.

## Causal reasoning (2/2)

Observational data




Causal reasoning

## Causal reasoning (2/2)


Causal DAG

Observational data




Causal reasoning

Confounding Bias (e.g., Simpson paradox):




Confounding Bias (e.g., Simpson paradox):



We must eliminate all confounding bias by adjusting for confounders.


Confounding Bias (e.g., Simpson paradox):



We must eliminate all confounding bias by adjusting for confounders.

Should we always adjust for all available variables?

Confounding Bias (e.g., Simpson paradox):




We must eliminate all confounding bias by adjusting for confounders.

Should we always adjust for all available variables? **No!**



## Key challenges of not using a causal DAG (2/3)

Bias due to incorrect adjustment for mediators:





## Key challenges of not using a causal DAG (3/3)

Bias due to adjusting for colliders:



## Key challenges of not using a causal DAG (3/3)


Bias due to adjusting for colliders:



We cannot adjust for everything that is temporally prior to the treatment!

## Example



In a study, we measure weekly exercise and cholesterol levels for various age groups.



What is the effect of exercise on cholesterol  $P(c \mid \text{do}(e))$ ?

## Example



In a study, we measure weekly exercise and cholesterol levels for various age groups.



What is the effect of exercise on cholesterol  $P(c \mid \text{do}(e))$ ?

## Example

In a study, we measure weekly exercise and cholesterol levels for various age groups.





What is the effect of exercise on cholesterol  $P(c | \text{do}(e))$ ?

$$P(a, e, c) = P(a)P(e | a)P(c | a, e) \quad (\text{Compatibility})$$

$$P(a, c | \text{do}(e)) = P(a)P(c | a, e) \quad (\text{Truncated factorization})$$

## Example

In a study, we measure weekly exercise and cholesterol levels for various age groups.



What is the effect of exercise on cholesterol  $P(c | \text{do}(e))$ ?

$$P(a, e, c) = P(a)P(e | a)P(c | a, e) \quad (\text{Compatibility})$$

$$P(a, c | \text{do}(e)) = P(a)P(c | a, e) \quad (\text{Truncated factorization})$$


$$P(c | \text{do}(e)) = \sum_a P(a)P(c | a, e) \quad (\text{marginalizing})$$

## Theorem

Given a causal DAG  $\mathcal{G}$  in which a subset  $\mathbb{V}$  of variables are measured, the causal effect  $P(y \mid \text{do}(a))$  is identifiable whenever  $\{A \cup Y \cup \text{Parents}(A)\} \subseteq \mathbb{V}$ , and is given by:

$$P(y \mid \text{do}(a)) = \sum_{Z \in \text{Pa}(A)} P(y \mid a, z)P(z)$$

- Sometimes the set of parents is too large. Is it possible to find a smaller set?
- Sometimes the set of observed parents is not sufficient for adjustment. Is it possible to find another set?



## The back-door criterion

A set of variables  $Z$  satisfies the back-door criterion relative to an ordered pair of variables  $(A, Y)$  in causal DAG  $\mathcal{G}$  if:

- No node in  $Z$  is a descendant of  $A$ ; and
- $Z$  blocks all paths between  $A$  and  $Y$  that contain an arrow pointing toward  $A$ .

A set of variables  $Z$  satisfies the back-door criterion relative to an ordered pair of variables  $(A, Y)$  in causal DAG  $\mathcal{G}$  if:

- No node in  $Z$  is a descendant of  $A$ ; and
- $Z$  blocks all paths between  $A$  and  $Y$  that contain an arrow pointing toward  $A$ .

### Theorem

If  $Z$  satisfies the back-door criterion with respect to  $(A, Y)$  and if  $P(a, z) > 0$ , then  $P(y \mid \text{do}(a))$  is identifiable and is given by:

$$P(y \mid \text{do}(a)) = \sum_z P(y \mid a, z)P(z).$$

A set of variables  $\mathbb{Z}$  satisfies the back-door criterion relative to an ordered pair of variables  $(A, Y)$  in causal DAG  $\mathcal{G}$  if:

- No node in  $\mathbb{Z}$  is a descendant of  $A$ ; and
- $\mathbb{Z}$  blocks all paths between  $A$  and  $Y$  that contain an arrow pointing toward  $A$ .

### Theorem


If  $\mathbb{Z}$  satisfies the back-door criterion with respect to  $(A, Y)$  and if  $P(a, z) > 0$ , then  $P(y | \text{do}(a))$  is identifiable and is given by:

$$P(y | \text{do}(a)) = \sum_z P(y | a, z)P(z).$$

This criterion can be extended to a set  $\mathbb{A}$  and a set  $\mathbb{Y}$ .

- Why "no node in  $\mathbb{Z}$  is a descendant of  $A$ "?
  - Why " $\mathbb{Z}$  blocks all paths between  $A$  and  $Y$  that contain an arrow pointing toward  $A$ "?

- Why "no node in  $Z$  is a descendant of  $A$ "?




To avoid blocking on intermediate causes.


- Why " $Z$  blocks all paths between  $A$  and  $Y$  that contain an arrow pointing toward  $A$ "?

## The back-door criterion: intuition

- Why "no node in  $\mathbb{Z}$  is a descendant of  $A$ "?



To avoid blocking on intermediate causes.




To avoid introducing artificial confounding bias resulting from conditioning on a collider.


- Why " $\mathbb{Z}$  blocks all paths between  $A$  and  $Y$  that contain an arrow pointing toward  $A$ "?

## The back-door criterion: intuition

- Why "no node in  $\mathbb{Z}$  is a descendant of  $A$ "?



To avoid blocking on intermediate causes.




To avoid introducing artificial confounding bias resulting from conditioning on a collider.

To avoid selection bias!


- Why " $\mathbb{Z}$  blocks all paths between  $A$  and  $Y$  that contain an arrow pointing toward  $A$ "?

## The back-door criterion: intuition

- Why "no node in  $Z$  is a descendant of  $A$ "?



To avoid blocking on intermediate causes.




To avoid introducing artificial confounding bias resulting from conditioning on a collider.

To avoid selection bias!


- Why " $Z$  blocks all paths between  $A$  and  $Y$  that contain an arrow pointing toward  $A$ "?

To eliminate confounding bias.


## Example

 $P(y \mid \text{do}(a))?$ 


## Example

 $P(y \mid \text{do}(a))?$  $Z ?$ 


## Example

 $P(y \mid \text{do}(a))?$  $Z ? \text{No}$ 

## Example


 $P(y \mid \text{do}(a))?$  $Z ? \text{ No}$   
 $X ?$ 

## Example

 $P(y \mid \text{do}(a))?$  $Z ? \text{No}$   
 $X ? \text{No}$ 

## Example

$P(y \mid \text{do}(a))?$




$Z ? \text{No}$

$X ? \text{No}$


$B ?$

## Example

 $P(y \mid \text{do}(a))?$  $Z ? \text{No}$  $X ? \text{No}$  $B ? \text{No}$

## Example

$P(y \mid \text{do}(a))?$




$Z ? \text{No}$

$X ? \text{No}$

$B ? \text{No}$


$C ?$

## Example

 $P(y \mid \text{do}(a))?$  $Z ? \text{No}$  $X ? \text{No}$  $B ? \text{No}$  $C ? \text{No}$

## Example

$P(y \mid \text{do}(a))?$



$Z ? \text{No}$

$X ? \text{No}$


$B ? \text{No}$

$C ? \text{No}$

$X, B ?$

## Example


$P(y \mid \text{do}(a))?$



$Z$  ? No  
 $X$  ? No  
 $B$  ? No  
 $C$  ? No  
 $X, B$ ? No

### Example


$P(y \mid \text{do}(a))?$



$Z ?$  No  
 $X ?$  No  
 $B ?$  No  
 $C ?$  No  
 $X, B ?$  No  
 $Z, X ?$

## Example


$P(y \mid \text{do}(a))?$



$Z ? \text{No}$   
 $X ? \text{No}$   
 $B ? \text{No}$   
 $C ? \text{No}$   
 $X, B ? \text{No}$   
 $Z, X ? \text{Yes}$

## Example


$P(y \mid \text{do}(a))?$



$Z ? \text{No}$   
 $X ? \text{No}$   
 $B ? \text{No}$   
 $C ? \text{No}$   
 $X, B ? \text{No}$   
 $Z, X ? \text{Yes}$   
 $Z, B ?$

## Example


$P(y \mid \text{do}(a))?$



$Z ? \text{No}$   
 $X ? \text{No}$   
 $B ? \text{No}$   
 $C ? \text{No}$   
 $X, B ? \text{No}$   
 $Z, X ? \text{Yes}$   
 $Z, B ? \text{Yes}$

### Example


$P(y \mid \text{do}(a))?$



$Z ? \text{No}$   
 $X ? \text{No}$   
 $B ? \text{No}$   
 $C ? \text{No}$   
 $X, B ? \text{No}$   
 $Z, X ? \text{Yes}$   
 $Z, B ? \text{Yes}$   
 $Z, X, B ?$

### Example

$P(y \mid \text{do}(a))?$



- $Z ? \text{No}$
- $X ? \text{No}$
- $B ? \text{No}$
- $C ? \text{No}$
- $X, B ? \text{No}$
- $Z, X ? \text{Yes}$
- $Z, B ? \text{Yes}$
- $Z, X, B ? \text{Yes}$

## Back-door criterion: example 1

## Example


In a study, we measure the number of coffee intake and IQ score for a sample of a population with various education level.

What is the effect of the number cups of coffee on IQ score  
 $P(i | \textcolor{red}{do}(c))$ ?

## Back-door criterion: example 1

## Example

In a study, we measure the number of coffee intake and IQ score for a sample of a population with various education level.




What is the effect of the number cups of coffee on IQ score  
 $P(i | \text{do}(c))$ ?

## Back-door criterion: example 1

### Example

In a study, we measure the number of coffee intake and IQ score for a sample of a population with various education level.



What is the effect of the number cups of coffee on IQ score  
 $P(i | \text{do}(c))$ ?

$$P(i | \text{do}(c)) = \sum_e P(i | c, e)P(e)$$


## Example

In this study, we aim to estimate the effect of the neighborhood's socioeconomic status (N) on the reduction of walking within the neighborhood (R),  $P(r | do(n))$ ?

## Back-door criterion: example 2

### Example


In this study, we aim to estimate the effect of the neighborhood's socioeconomic status (N) on the reduction of walking within the neighborhood (R),  $P(r | do(n))$ ?



## Back-door criterion: example 2

### Example


In this study, we aim to estimate the effect of the neighborhood's socioeconomic status (N) on the reduction of walking within the neighborhood (R),  $P(r | do(n))$ ?



## Back-door criterion: example 2

### Example

In this study, we aim to estimate the effect of the neighborhood's socioeconomic status (N) on the reduction of walking within the neighborhood (R),  $P(r | do(n))$ ?




$$P(r | do(n)) = \sum P(r|n, l_2, p)P(l_2, p)$$

## Back-door criterion: example 3

## Example


Suppose that all variable are temporally prior to  $A$  and  $Y$ .  
 $P(y \mid \text{do}(a))$ ?



## Back-door criterion: example 3

## Example

Suppose that all variable are temporally prior to  $A$  and  $Y$ .  
 $P(y \mid \text{do}(a))$ ?




Many sets of variables can satisfy the back-door criterion:

- All these sets successfully remove bias.
- However, in practice, they do not all yield the same variance.

Many sets of variables can satisfy the back-door criterion:



- All these sets successfully remove bias.
- However, in practice, they do not all yield the same variance.



# Finding the optimal set satisfying the back-door criterion

Many sets of variables can satisfy the back-door criterion:



- All these sets successfully remove bias.
- However, in practice, they do not all yield the same variance.



# Finding the optimal set satisfying the back-door criterion

Many sets of variables can satisfy the back-door criterion:



- All these sets successfully remove bias.
- However, in practice, they do not all yield the same variance.



# Finding the optimal set satisfying the back-door criterion

Many sets of variables can satisfy the back-door criterion:

- All these sets successfully remove bias.
- However, in practice, they do not all yield the same variance.
- Is it possible to find the **optimal** set that yields the smallest variance?





# Finding the optimal set satisfying the back-door criterion

Many sets of variables can satisfy the back-door criterion:

- All these sets successfully remove bias.
- However, in practice, they do not all yield the same variance.
- Is it possible to find the **optimal** set that yields the smallest variance?

A set  $Z$  satisfying the back-door criterion relative to  $(A, Y)$  **optimal** if:

- $Z$  includes all parents of  $Y$  that are not mediators between  $A$  and  $Y$ , and





# Finding the optimal set satisfying the back-door criterion

Many sets of variables can satisfy the back-door criterion:

- All these sets successfully remove bias.
- However, in practice, they do not all yield the same variance.
- Is it possible to find the **optimal** set that yields the smallest variance?

A set  $Z$  satisfying the back-door criterion relative to  $(A, Y)$  **optimal** if:

- $Z$  includes all parents of  $Y$  that are not mediators between  $A$  and  $Y$ , and
- $Z$  includes all parents of any mediator between  $A$  and  $Y$ , that are not themselves mediators between  $A$  and  $Y$ .




## Example

How to optimally estimate the effect of new hypertension drug ( $H$ ) on blood pressure ( $B$ )  $P(b | \text{do}(h))$ ?

# Optimal set in action

## Example


How to optimally estimate the effect of new hypertension drug ( $H$ ) on blood pressure ( $B$ )  $P(b | \text{do}(h))$ ?



# Optimal set in action

## Example


How to optimally estimate the effect of new hypertension drug ( $H$ ) on blood pressure ( $B$ )  $P(b | \text{do}(h))$ ?



# Optimal set in action

## Example

How to optimally estimate the effect of new hypertension drug ( $H$ ) on blood pressure ( $B$ )  $P(b | \text{do}(h))$ ?



$$P(b | \text{do}(h)) = \sum_{g,e} P(b|h, g, e)P(g, e)$$

## Association Criterion 1 for identifying confounding factors (incorrect)

Let  $\mathbb{Z}$  be the set of observed variables in a problem that are not affected by  $A$ . The set  $\mathbb{Z}$  satisfies Association Criterion 1 if each element  $Z \in \mathbb{Z}$  meets the following conditions:


- $Z$  is associated with  $A$ ; and
- $Z$  is associated with  $Y$ , conditionally on  $A$ .

# Association Criterion 1 for identifying confounding factors (incorrect)

Let  $Z$  be the set of observed variables in a problem that are not affected by  $A$ . The set  $Z$  satisfies Association Criterion 1 if each element  $Z \in Z$  meets the following conditions:

- $Z$  is associated with  $A$ ; and
- $Z$  is associated with  $Y$ , conditionally on  $A$ .

Counterexample




## Association Criterion 1 for identifying confounding factors (incorrect)

Let  $\mathbb{Z}$  be the set of observed variables in a problem that are not affected by  $A$ . The set  $\mathbb{Z}$  satisfies Association Criterion 1 if each element  $Z \in \mathbb{Z}$  meets the following conditions:

- $Z$  is associated with  $A$ ; and
- $Z$  is associated with  $Y$ , conditionally on  $A$ .

Counterexample




$Z$  is associated with  $A$  and  $Z$  is associated with  $Y$ , conditionally on  $A$

## Association Criterion 1 for identifying confounding factors (incorrect)

Let  $Z$  be the set of observed variables in a problem that are not affected by  $A$ . The set  $Z$  satisfies Association Criterion 1 if each element  $Z \in Z$  meets the following conditions:

- $Z$  is associated with  $A$ ; and
- $Z$  is associated with  $Y$ , conditionally on  $A$ .

Counterexample



$Z$  is associated with  $A$  and  $Z$  is associated with  $Y$ , conditionally on  $A$


Should we adjust on  $Z$ ?

## Association Criterion 1 for identifying confounding factors (incorrect)

Let  $Z$  be the set of observed variables in a problem that are not affected by  $A$ . The set  $Z$  satisfies Association Criterion 1 if each element  $Z \in Z$  meets the following conditions:

- $Z$  is associated with  $A$ ; and
- $Z$  is associated with  $Y$ , conditionally on  $A$ .

Counterexample



$Z$  is associated with  $A$  and  $Z$  is associated with  $Y$ , conditionally on  $A$


Should we adjust on  $Z$ ? No!

## Association Criterion 1 for identifying confounding factors (incorrect)

Let  $Z$  be the set of observed variables in a problem that are not affected by  $A$ . The set  $Z$  satisfies Association Criterion 1 if each element  $Z \in Z$  meets the following conditions:

- $Z$  is associated with  $A$ ; and
- $Z$  is associated with  $Y$ , conditionally on  $A$ .

Counterexample



$Z$  is associated with  $A$  and  $Z$  is associated with  $Y$ , conditionally on  $A$

Should we adjust on  $Z$ ? No!

## Association Criterion 2 for identifying confounding factors (incorrect)

Let  $\mathbb{Z}$  be the set of variables in a problem that are not affected by  $X$ . The set  $\mathbb{Z}$  satisfies Association Criterion 2 if it can be divided into two subsets,  $\mathbb{Z}_1$  and  $\mathbb{Z}_2$ , which meet the following conditions:


- $\mathbb{Z}_1$  is associated with  $A$ ; and
- $\mathbb{Z}_2$  is associated with  $Y$ , conditionally on  $A$  and  $\mathbb{Z}_1$ .

## Association Criterion 2 for identifying confounding factors (incorrect)

Let  $\mathbb{Z}$  be the set of variables in a problem that are not affected by  $X$ . The set  $\mathbb{Z}$  satisfies Association Criterion 2 if it can be divided into two subsets,  $\mathbb{Z}_1$  and  $\mathbb{Z}_2$ , which meet the following conditions:

- $\mathbb{Z}_1$  is associated with  $A$ ; and
- $\mathbb{Z}_2$  is associated with  $Y$ , conditionally on  $A$  and  $\mathbb{Z}_1$ .

Counterexample




## Association Criterion 2 for identifying confounding factors (incorrect)

Let  $\mathbb{Z}$  be the set of variables in a problem that are not affected by  $X$ . The set  $\mathbb{Z}$  satisfies Association Criterion 2 if it can be divided into two subsets,  $\mathbb{Z}_1$  and  $\mathbb{Z}_2$ , which meet the following conditions:

- $\mathbb{Z}_1$  is associated with  $A$ ; and
- $\mathbb{Z}_2$  is associated with  $Y$ , conditionally on  $A$  and  $\mathbb{Z}_1$ .

Counterexample




Car type is associated with Exposure

## Association Criterion 2 for identifying confounding factors (incorrect)

Let  $\mathbb{Z}$  be the set of variables in a problem that are not affected by  $X$ . The set  $\mathbb{Z}$  satisfies Association Criterion 2 if it can be divided into two subsets,  $\mathbb{Z}_1$  and  $\mathbb{Z}_2$ , which meet the following conditions:

- $\mathbb{Z}_1$  is associated with  $A$ ; and
- $\mathbb{Z}_2$  is associated with  $Y$ , conditionally on  $A$  and  $\mathbb{Z}_1$ .

Counterexample




Car type is associated with Exposure  
Should we adjust on Car type?

## Association Criterion 2 for identifying confounding factors (incorrect)

Let  $\mathbb{Z}$  be the set of variables in a problem that are not affected by  $X$ . The set  $\mathbb{Z}$  satisfies Association Criterion 2 if it can be divided into two subsets,  $\mathbb{Z}_1$  and  $\mathbb{Z}_2$ , which meet the following conditions:

- $\mathbb{Z}_1$  is associated with  $A$ ; and
- $\mathbb{Z}_2$  is associated with  $Y$ , conditionally on  $A$  and  $\mathbb{Z}_1$ .

Counterexample




Car type is associated with Exposure  
Should we adjust on Car type? No!


- The back-door criterion is sound but not complete:

- The back-door criterion is sound but not complete:
  - ▶ If there exists a set that satisfy the back-door criterion for  $P(y | \textcolor{red}{do}(a))$ , then  $P(y | \textcolor{red}{do}(a))$  is identifiable;
  - ▶ If there exists a no set that satisfy the back-door criterion for  $P(y | \textcolor{red}{do}(a))$ , then  $P(y | \textcolor{red}{do}(a))$  is not necesarly not identifiable.

## Going beyond the back-door




## Going beyond the back-door




- $P(z \mid \textcolor{red}{do}(a)) = P(z \mid a)$  (No back-door path)

## Going beyond the back-door



- $P(z \mid \text{do}(a)) = P(z \mid a)$  (No back-door path)
- $P(y \mid \text{do}(z)) = \sum_a P(y \mid z, a)P(a)$  ( $A$  blocks the back-door)


# Going beyond the back-door



- $P(z \mid \text{do}(a)) = P(z \mid a)$  (No back-door path)
- $P(y \mid \text{do}(z)) = \sum_a P(y \mid z, a)P(a)$  ( $A$  blocks the back-door)


$$P(y \mid \text{do}(a)) = \sum_z P(y \mid \text{do}(z))P(z \mid \text{do}(a))$$

# Going beyond the back-door



- $P(z \mid \text{do}(a)) = P(z \mid a)$  (No back-door path)
- $P(y \mid \text{do}(z)) = \sum_a P(y \mid z, a)P(a)$  ( $A$  blocks the back-door)

$$\begin{aligned}
 P(y \mid \text{do}(a)) &= \sum_z P(y \mid \text{do}(z))P(z \mid \text{do}(a)) \\
 &= \sum_z P(z \mid a) \sum_{a'} P(y \mid z, a')P(a')
 \end{aligned}$$



A set of variables  $\mathbb{Z}$  satisfies the front-door criterion relative to an ordered pair of variables  $(A, Y)$  in causal DAG  $\mathcal{G}$  if:

- $\mathbb{Z}$  intercepts all directed paths from  $A$  to  $Y$ ;
- There is no back-door path from  $A$  to  $\mathbb{Z}$ ;
- All back-door paths from  $\mathbb{Z}$  to  $Y$  are blocked by  $A$ .

## Theorem ([3])

If  $\mathbb{Z}$  satisfies the front-door criterion relative to  $(A, Y)$  and if  $P(a, z) > 0$ , then the causal effect of  $A$  on  $Y$  is identifiable and is given by

$$P(y \mid \textcolor{red}{do}(a)) = \sum_z P(z \mid a) \sum_{a'} P(y \mid a', z)P(a').$$

## Front-door criterion: example


In a study, we measure the tar and the % of cancer among smokers and non smokers in a randomly selected sample of the population.

What is the effect of smoking on cancer  $P(c \mid \text{do}(s))$ ?

## Front-door criterion: example

In a study, we measure the tar and the % of cancer among smokers and non smokers in a randomly selected sample of the population.


What is the effect of smoking on cancer  $P(c | do(s))$ ?



## Front-door criterion: example

In a study, we measure the tar and the % of cancer among smokers and non smokers in a randomly selected sample of the population.

What is the effect of smoking on cancer  $P(c | do(s))$ ?




$$P(c | do(s)) = \sum_t P(t|s) \sum_{s'} P(c | t, s') P(s')$$

## Back-door and Front-door criteria in a real application

In this study, Piccininni et al. were interested in estimating the effect of Mobile Stroke Unit dispatch (M) on functional outcomes (F). In other words,  $P(f \mid \text{do}(m))$ ?


## Back-door and Front-door criteria in a real application

In this study, Piccininni et al. were interested in estimating the effect of Mobile Stroke Unit dispatch (M) on functional outcomes (F). In other words,  $P(f \mid \text{do}(m))$ ?




## Back-door and Front-door criteria in a real application

In this study, Piccininni et al. were interested in estimating the effect of Mobile Stroke Unit dispatch (M) on functional outcomes (F). In other words,  $P(f \mid \text{do}(m))$ ?



The back-door criterion is not satisfied for any set of variables. However, it is possible to identify  $P(f \mid \text{do}(m))$  by leveraging a combination of the back-door and front-door criteria!

In this study, Piccininni et al. were interested in estimating the effect of Mobile Stroke Unit dispatch (M) on functional outcomes (F). In other words,  $P(f \mid \text{do}(m))$ ?



The back-door criterion is not satisfied for any set of variables. However, it is possible to identify  $P(f \mid \text{do}(m))$  by leveraging a combination of the back-door and front-door criteria!

Best paper in Epidemiology in 2024!

## Incompleteness of the front-door criterion

- The front-door criterion is sound but not complete:

- The front-door criterion is sound but not complete:
  - ▶ If there exists a set that satisfy the front-door criterion for  $P(y | \textcolor{red}{do}(x))$ , then  $P(y | \textcolor{red}{do}(x))$  is identifiable;
  - ▶ If there exists a no set that satisfy the front-door criterion for  $P(y | \textcolor{red}{do}(x))$ , then  $P(y | \textcolor{red}{do}(x))$  is not necesarly not identifiable.

- The front-door criterion is sound but not complete:
  - ▶ If there exists a set that satisfy the front-door criterion for  $P(y | \text{do}(x))$ , then  $P(y | \text{do}(x))$  is identifiable;
  - ▶ If there exists a no set that satisfy the front-door criterion for  $P(y | \text{do}(x))$ , then  $P(y | \text{do}(x))$  is not necesarly not identifiable.

The combination of the back-door and front-door criterions is also not complete.

## A glimpse of the do-calculus [3]

The do calculus consists of three rules :

- Rule 1: generalization of the d-seperation
- Rule 2: generalization of back-door criterion
- Rule 3: complicated

## A glimpse of the do-calculus [3]

The do calculus consists of three rules :

- Rule 1: generalization of the d-seperation
- Rule 2: generalization of back-door criterion
- Rule 3: complicated

$P(y | do(x))$  is identifiable iff by repeatedly applying the rules of do-calculus, we obtain a do-free expression.

## A glimpse of the do-calculus [3]


The do calculus consists of three rules :

- Rule 1: generalization of the d-seperation
- Rule 2: generalization of back-door criterion
- Rule 3: complicated



$P(y | do(x))$  is identifiable iff by repeatedly applying the rules of do-calculus, we obtain a do-free expression.

The do calculus is sound and complete!




## Mutilated Graphs







## Mutilated Graphs

 $\mathcal{G}$  $\mathcal{G}_{\bar{A}}$

## Mutilated Graphs



# Mutilated Graphs



## Theorem

Let  $\mathcal{G} = (\mathbb{V}, \mathbb{E})$  be a causal graph. Let  $\mathbb{X}, \mathbb{Y}, \mathbb{Z}, \mathbb{W} \subseteq \mathbb{V}$  be disjoint.  
We have:

$$P(y|do(x), z, w) = P(y|do(x), w) \quad \text{if} \quad \mathbb{Y} \perp\!\!\!\perp_{\mathcal{G}_{\overline{\mathbb{X}}}} \mathbb{Z} | \mathbb{X}, \mathbb{W}$$

$$P(y|do(x), do(z), w) = P(y|do(x), z, w) \quad \text{if} \quad \mathbb{Y} \perp\!\!\!\perp_{\mathcal{G}_{\overline{\mathbb{X}\mathbb{Z}}}} \mathbb{Z} | \mathbb{X}, \mathbb{W}$$

$$P(y|do(x), do(z), w) = P(y|do(x), w) \quad \text{if} \quad \mathbb{Y} \perp\!\!\!\perp_{\mathcal{G}_{\overline{\mathbb{X}\mathbb{Z}(w)}}} \mathbb{Z} | \mathbb{X}, \mathbb{W}$$

where  $\mathbb{Z}(w)$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}_{\overline{\mathbb{X}}}$

## Intuition for Rule 3

$$P(y|\cancel{do(x)}, \cancel{do(z)}, w) = P(y|\cancel{do(x)}, w) \quad \text{if } (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z} | \mathbb{X}, \mathbb{W})_{\mathcal{G}_{\overline{\mathbb{X}\mathbb{Z}(\mathbb{W})}}}$$

where  $\mathbb{Z}(\mathbb{W})$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}_{\overline{\mathbb{X}}}$

## Intuition for Rule 3

$$P(y|do(z), w) = P(y|w) \quad \text{if} \quad (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W})_{\mathcal{G}_{\overline{\mathbb{Z}(\mathbb{W})}}}$$

where  $\mathbb{Z}(\mathbb{W})$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}$

## Intuition for Rule 3

$$P(y|do(z), w) = P(y|w) \quad \text{if } (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W})_{\mathcal{G}_{\overline{\mathbb{Z}(\mathbb{W})}}}$$

where  $\mathbb{Z}(\mathbb{W})$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}$

Suppose

$$\begin{aligned} & P(y | do(z), w_1, w_2) \\ &= P(y | w_1, w_2) \\ & \text{if } (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|W_1, W_2)_{\mathcal{G}_{\overline{\mathbb{Z}}}} \end{aligned}$$


## Intuition for Rule 3

$$P(y|do(z), w) = P(y|w) \quad \text{if} \quad (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W})_{\mathcal{G}_{\overline{\mathbb{Z}}(\mathbb{W})}}$$

where  $\mathbb{Z}(\mathbb{W})$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}$

Suppose

$$\begin{aligned} P(y | do(z), w_1, w_2) \\ = P(y | w_1, w_2) \\ \text{if } (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W}_1, \mathbb{W}_2)_{\mathcal{G}_{\overline{\mathbb{Z}}}} \end{aligned}$$




## Intuition for Rule 3

$$P(y|do(z), w) = P(y|w) \quad \text{if} \quad (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W})_{\mathcal{G}_{\overline{\mathbb{Z}}(\mathbb{W})}}$$


where  $\mathbb{Z}(\mathbb{W})$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}$

Suppose

$$\begin{aligned} P(y | do(z), w_1, w_2) \\ = P(y | w_1, w_2) \\ \text{if } (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W}_1, \mathbb{W}_2)_{\mathcal{G}_{\overline{\mathbb{Z}}}} \end{aligned}$$



Intervention



Causal reasoning

## Intuition for Rule 3


$$P(y|do(z), w) = P(y|w) \quad \text{if} \quad (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W})_{\mathcal{G}_{\overline{\mathbb{Z}}(\mathbb{W})}}$$

where  $\mathbb{Z}(\mathbb{W})$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}$


Suppose

$$\begin{aligned} P(y | do(z), w_1, w_2) \\ = P(y | w_1, w_2) \end{aligned}$$

if  $(Y \perp\!\!\!\perp Z|W_1, W_2)_{\mathcal{G}_{\overline{Z}}}$



Intervention



Causal reasoning

## Intuition for Rule 3


$$P(y|do(z), w) = P(y|w) \quad \text{if} \quad (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W})_{\mathcal{G}_{\overline{\mathbb{Z}(W)}}}$$

where  $\mathbb{Z}(\mathbb{W})$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}$


Suppose

$$\begin{aligned} P(y | do(z), w_1, w_2) \\ = P(y | w_1, w_2) \end{aligned}$$

if  $(Y \perp\!\!\!\perp Z|W_1, W_2)_{\mathcal{G}_{\overline{Z}}}$



Intervention



Causal reasoning

## Intuition for Rule 3

$$P(y|do(z), w) = P(y|w) \quad \text{if} \quad (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|\mathbb{W})_{\mathcal{G}_{\overline{\mathbb{Z}}(\mathbb{W})}}$$


where  $\mathbb{Z}(\mathbb{W})$  is the set of  $\mathbb{Z}$ -vertices that are not ancestors of any  $\mathbb{W}$ -vertex in  $\mathcal{G}$

Suppose


$$\cancel{P(y | do(z), w_1, w_2)}$$

$$\cancel{= P(y | w_1, w_2)}$$

$$\underline{\text{if } (\mathbb{Y} \perp\!\!\!\perp \mathbb{Z}|W_1, W_2)_{\mathcal{G}_{\overline{\mathbb{Z}}}}}$$




$$\mathbb{Y} \perp\!\!\!\perp_{\mathcal{G}_{\overline{\mathbb{Z}}}} W_2 | W_1$$




$$\mathbb{Y} \not\perp\!\!\!\perp_{\mathcal{G}_{\overline{\mathbb{Z}}}} W_2 | W_1$$

## Theorem ([3, 9])

A causal effect  $P(y | \text{do}(x))$  is identifiable in a model characterized by a graph  $\mathcal{G}$  if and only if there exists a finite sequence of transformations, each conforming to one of the Rules 1-3, that reduces  $P(y | \text{do}(x))$  into a standard (i.e., "do"-free) probability expression involving observed quantities.




What's the effect of exercise on cholesterol?




What's the effect of exercise on cholesterol?

$$P(c \mid \text{do}(e)) = \sum_a P(c \mid \text{do}(e), a)P(a \mid \text{do}(e)) \quad (\text{Probability Axioms})$$



What's the effect of exercise on cholesterol?


$$\begin{aligned}
 P(c \mid \text{do}(e)) &= \sum_a P(c \mid \text{do}(e), a)P(a \mid \text{do}(e)) && \text{(Probability Axioms)} \\
 &= \sum_a P(c \mid e, a)P(a \mid \text{do}(e)) && \text{(Rule 2)}
 \end{aligned}$$



What's the effect of exercise on cholesterol?


$$\begin{aligned}
 P(c \mid \text{do}(e)) &= \sum_a P(c \mid \text{do}(e), a)P(a \mid \text{do}(e)) && \text{(Probability Axioms)} \\
 &= \sum_a P(c \mid e, a)P(a \mid \text{do}(e)) && \text{(Rule 2)} \\
 &= \sum_a P(c \mid e, a)P(a) && \text{(Rule 3)}
 \end{aligned}$$

## From the do-calculus to the front-door adjustment



What's the effect of smoking on cancer?


## From the do-calculus to the front-door adjustment



What's the effect of smoking on cancer?

$$P(c \mid \text{do}(s)) = \sum_t P(c \mid \text{do}(s), t)P(t \mid \text{do}(s)) \quad (\text{Probability Axioms})$$


# From the do-calculus to the front-door adjustment



What's the effect of smoking on cancer?

$$\begin{aligned}
 P(c \mid \text{do}(s)) &= \sum_t P(c \mid \text{do}(s), t)P(t \mid \text{do}(s)) && \text{(Probability Axioms)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid \text{do}(s)) && \text{(Rule 2)}
 \end{aligned}$$


# From the do-calculus to the front-door adjustment



What's the effect of smoking on cancer?

$$\begin{aligned}
 P(c \mid \text{do}(s)) &= \sum_t P(c \mid \text{do}(s), t)P(t \mid \text{do}(s)) && \text{(Probability Axioms)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid \text{do}(s)) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid s) && \text{(Rule 2)}
 \end{aligned}$$


# From the do-calculus to the front-door adjustment



What's the effect of smoking on cancer?

$$\begin{aligned}
 P(c \mid \text{do}(s)) &= \sum_t P(c \mid \text{do}(s), t)P(t \mid \text{do}(s)) && \text{(Probability Axioms)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid \text{do}(s)) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid s) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(t))P(t \mid s) && \text{(Rule 3)}
 \end{aligned}$$


# From the do-calculus to the front-door adjustment



What's the effect of smoking on cancer?

$$\begin{aligned}
 P(c \mid \text{do}(s)) &= \sum_t P(c \mid \text{do}(s), t)P(t \mid \text{do}(s)) && \text{(Probability Axioms)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid \text{do}(s)) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid s) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(t))P(t \mid s) && \text{(Rule 3)} \\
 &= \sum_{s'} \sum_t P(c \mid \text{do}(t), s')P(s' \mid \text{do}(t))P(t \mid s) && \text{(Probability Axioms)}
 \end{aligned}$$


# From the do-calculus to the front-door adjustment



What's the effect of smoking on cancer?

$$\begin{aligned}
 P(c \mid \text{do}(s)) &= \sum_t P(c \mid \text{do}(s), t)P(t \mid \text{do}(s)) && \text{(Probability Axioms)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid \text{do}(s)) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid s) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(t))P(t \mid s) && \text{(Rule 3)} \\
 &= \sum_{s'} \sum_t P(c \mid \text{do}(t), s')P(s' \mid \text{do}(t))P(t \mid s) && \text{(Probability Axioms)} \\
 &= \sum_{s'} \sum_t P(c \mid t, s')P(s' \mid \text{do}(t))P(t \mid s) && \text{(Rule 2)}
 \end{aligned}$$

# From the do-calculus to the front-door adjustment



What's the effect of smoking on cancer?

$$\begin{aligned}
 P(c \mid \text{do}(s)) &= \sum_t P(c \mid \text{do}(s), t)P(t \mid \text{do}(s)) && \text{(Probability Axioms)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid \text{do}(s)) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(s), \text{do}(t))P(t \mid s) && \text{(Rule 2)} \\
 &= \sum_t P(c \mid \text{do}(t))P(t \mid s) && \text{(Rule 3)} \\
 &= \sum_{s'} \sum_t P(c \mid \text{do}(t), s')P(s' \mid \text{do}(t))P(t \mid s) && \text{(Probability Axioms)} \\
 &= \sum_{s'} \sum_t P(c \mid t, s')P(s' \mid \text{do}(t))P(t \mid s) && \text{(Rule 2)} \\
 &= \sum_{s'} \sum_t P(c \mid t, s')P(s')P(t \mid s) && \text{(Rule 3)}
 \end{aligned}$$

Limitations of the do-calculus:

- The do-calculus demands a lot of manual labor
- Non-identifiability is complicated

Limitations of the do-calculus:

- The do-calculus demands a lot of manual labor
- Non-identifiability is complicated

Is it possible automatize it?


Limitations of the do-calculus:

- The do-calculus demands a lot of manual labor
- Non-identifiability is complicated

Is it possible automatize it? Yes! There exists many algorithms, e.g., the ID algorithm [9].


## Exercise 4

Consider that in the following causal DAG, only  $A$  and  $Y$ , and one additional variable can be measured. Which variable would allow the identification of  $P(y \mid do(a))$ ?



## Exercise 5

- Consider the following causal DAG. List all sets of variables that satisfy the back-door criterion for  $P(y | do(a))$ ;




## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | \textcolor{red}{do}(a))$ ?


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid do(a))$ ?



- $Z$  blocks all back-door paths and it is not a descendant of  $A$   
 $\implies \{Z\}$  is a good adjustment set.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | \text{do}(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



- $Z$  blocks all back-door paths and it is not a descendant of  $A$   
 $\implies \{Z\}$  is a good adjustment set.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | \text{do}(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



- $Z$  blocks all back-door paths and it is not a descendant of  $A$   
 $\implies \{Z\}$  is a good adjustment set.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | \text{do}(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid do(a))$ ?



- $Z$  activates a back-door path  
 $\implies \{Z\}$  is a bad adjustment set.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



- $Z$  blocks all back-door paths and it is not a descendant of  $A$   
 $\implies \{Z\}$  is a good adjustment set.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid \text{do}(a))$ ?



- $Z$  d-separates  $A$  from  $Y$   
 $\implies \{Z\}$  is a bad adjustment set.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | \text{do}(a))$ ?



- $Z$  d-separates  $A$  from  $Y$   
 $\implies \{Z\}$  is a bad adjustment set.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



- $Z$  does not open any back-door paths from  $A$  to  $Y$   
 $\implies \{Z\}$  is a neutral adjustment set;
- Adjusting on  $Z$  can reduce the variation of  $Y$ , and helps improve the precision of the estimate in finite samples.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



- $Z$  does not open any back-door paths from  $A$  to  $Y$   
 $\implies \{Z\}$  is a neutral adjustment set;
- Adjusting on  $Z$  can reduce the variation of  $A$  and so may hurt the precision of the estimate in finite samples.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid \text{do}(a))$ ?



- $Z$  does not block existing back-door path from  $A$  to  $Y$   
 $\implies \{Z\}$  is a bad adjustment set;
- In linear models, adjusting on  $Z$  amplify any existing bias.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid \text{do}(a))$ ?



- Selection bias  
 $\implies \{Z\}$  is a bad control.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



## Exercise 6


Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



- $Z$  does not open any back-door paths from  $A$  to  $Y$   
 $\implies \{Z\}$  is a neutral adjustment set;
- Adjusting on  $Z$  can reduce the variation of  $W$ , and helps improve the precision of the estimate in finite samples.


## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y | do(a))$ ?



## Exercise 6

Is  $\{Z\}$  a good, bad or neutral adjustment set for  $P(y \mid \text{do}(a))$ ?



- $Z$  is a descendant of  $A$   
 $\implies \{Z\}$  is a bad adjustment set.

# 4

## Counterfactuals

Structural causal models

Counterfactual reasoning

Mediation analysis

Difference with the potential outcome framework

# 4

## Counterfactuals


Structural causal models

Counterfactual reasoning


Mediation analysis

Difference with the potential outcome framework

<https://www.youtube.com/embed/0lpY0Kt4bn8?rel=0&autoplay=1>




## Causal mechanism




$$C = f(A, B, I)$$

# Causal mechanism




$$C := f_c(A, B, I, \xi_c)$$

# Causal mechanism




$$C := f_c(A, B, I, \xi_c)$$

# Structural causal model




# Structural causal model



$$M : \left\{ \begin{array}{l} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := f_c(A, B, I, \xi_c) \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{array} \right.$$


# Structural causal model



$$M : \left\{ \begin{array}{l} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := f_c(A, B, I, \xi_c) \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{array} \right.$$

A structural causal model (SCM) is a tuple that contains:

# Structural causal model




$$M : \left\{ \begin{array}{l} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := f_c(A, B, I, \xi_c) \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{array} \right.$$

A structural causal model (SCM) is a tuple that contains:

- Endogenous variables

# Structural causal model




$$M : \begin{cases} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := f_c(A, B, I, \xi_c) \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{cases}$$

A structural causal model (SCM) is a tuple that contains:

- Endogenous variables
- Exogenous variables

# Structural causal model



$$M : \begin{cases} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := f_c(A, B, I, \xi_c) \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{cases}$$

A structural causal model (SCM) is a tuple that contains:

- Endogenous variables
- Exogenous variables
- Causal mechanisms for generating endogenous variables

$$M : \begin{cases} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := f_c(A, B, I, \xi_c) \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{cases}$$

$$M : \begin{cases} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := f_c(A, B, I, \xi_c) \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{cases}$$

$$M_c : \begin{cases} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := c \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{cases}$$

# Interventions in a structural causal model

$$M : \left\{ \begin{array}{l} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := f_c(A, B, I, \xi_c) \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{array} \right.$$

$$M_c : \left\{ \begin{array}{l} A := f_a(\xi_a) \\ B := f_b(A, H, \xi_b) \\ C := c \\ D := f_d(C, F, \xi_d) \\ E := f_e(B, G, \xi_e) \\ F := f_f(C, G, \xi_f) \\ G := f_g(\xi_g) \\ H := f_h(G, \xi_h) \\ I := f_i(G, \xi_i) \end{array} \right.$$

# 4

## Counterfactuals

Structural causal models

Counterfactual reasoning


Mediation analysis

Difference with the potential outcome framework

# Mediation analysis in linear SCMs

SCM

Observational data




Counterfactual reasoning

# Mediation analysis in linear SCMs

SCM

Observational data



Counterfactual reasoning

SCMs naturally help us in computing counterfactual queries:


$$P(Y_{A=a} \mid A = a_0, Y = y_0)$$

1. Abduction: use the observations to determine the value of  $\xi_y$
2. Action: modify the model  $M$  by removing the structural equations for the variable  $A$  and replacing it with the appropriate functions  $A = a$ , to obtain the interventional SCM  $M_a$
3. Prediction: use  $M_a$  and the value of  $\xi_y$  to compute the value of  $Y_{A=a}$


If  $Y_{A=a} \neq y_0$  then  $a_0$  is an actual cause of  $y_0$

## Computing counterfactuals using SCMs: example


Factual world




Counterfactual world



Factual world




Conterfactual world




Toy example (without noise on  $C$ ,  $W$ , and  $G$ ):

$$M : \begin{cases} H & (B \text{ hits } A) := \xi_h \\ C & (A \text{ changes trajectory}) := H \\ W & (A \text{ hits the wall}) := 1 - C \\ G & (A \text{ reaches the goal}) := 1 - W \end{cases}$$

Factual world



Conterfactual world




Toy example (without noise on  $C$ ,  $W$ , and  $G$ ):


$$M : \begin{cases} H & (B \text{ hits } A) := \xi_h \\ C & (A \text{ changes trajectory}) := H \\ W & (A \text{ hits the wall}) := 1 - C \\ G & (A \text{ reaches the goal}) := 1 - W \end{cases}$$

$$\quad \quad \quad \begin{cases} H := 1 \\ C := 1 \\ W := 0 \\ G := 1 \end{cases}$$

Factual world



Conterfactual world




Toy example (without noise on  $C$ ,  $W$ , and  $G$ ):

$$M : \begin{cases} H & (B \text{ hits } A) := \xi_h \\ C & (A \text{ changes trajectory}) := H \\ W & (A \text{ hits the wall}) := 1 - C \\ G & (A \text{ reaches the goal}) := 1 - W \end{cases}$$


$$M_{H=0} : \begin{cases} H := 1 \\ C := 1 \\ W := 0 \\ G := 1 \end{cases}$$

$$M_{H=0} : \begin{cases} H := 0 \\ C := 0 \\ W := 1 \\ G := 0 \end{cases}$$

Factual world



Conterfactual world



Toy example (without noise on  $C$ ,  $W$ , and  $G$ ):

$$M : \begin{cases} H & (B \text{ hits } A) := \xi_h \\ C & (A \text{ changes trajectory}) := H \\ W & (A \text{ hits the wall}) := 1 - C \\ G & (A \text{ reaches the goal}) := 1 - W \end{cases}$$

$$\begin{cases} H := 1 \\ C := 1 \\ W := 0 \\ G := 1 \end{cases}$$

$$M_{H=0} : \begin{cases} H := 0 \\ C := 0 \\ W := 1 \\ G := 0 \end{cases}$$

$G_{H=0} \neq 1 \implies B \text{ causes } A \text{ to reach the goal}$

There are methods for computing counterfactual queries directly from a causal DAG [9]; however, these are beyond the scope of this tutorial.

We will focus on a specific class of counterfactual queries that are particularly relevant for mediation analysis.

# 4

## Counterfactuals

Structural causal models


Counterfactual reasoning

Mediation analysis

Difference with the potential outcome framework

# Different types of causal effects


There exists three main types  
of causal effects:



# Different types of causal effects

There exists three main types  
of causal effects:


- Total effect



# Different types of causal effects

There exists three main types of causal effects:


- Total effect
- Direct effect



# Different types of causal effects

There exists three main types of causal effects:

- Total effect
- Direct effect
- Indirect effect



## Natural direct effect (NDE)

Natural direct effect (NDE):

$$NDE(x, x'; Y) = \mathbb{E}(y \mid do(x, pa_{Y \setminus X}^{x'})) - \mathbb{E}(y \mid do(x', pa_{Y \setminus X}^{x'}))$$

where  $pa_{Y \setminus X}^{x'}$  is the value of  $Pa(Y) \setminus \{X\}$  when  $X = x'$ .

## Natural direct effect (NDE)

Natural direct effect (NDE):

$$NDE(x, x'; Y) = \mathbb{E}(y \mid do(x, pa_{Y \setminus X}^{x'})) - \mathbb{E}(y \mid do(x', pa_{Y \setminus X}^{x'}))$$

where  $pa_{Y \setminus X}^{x'}$  is the value of  $Pa(Y) \setminus \{X\}$  when  $X = x'$ .

Advantage:

- It allows for the complete decomposition of the total effect into the NDE and NIE

## Natural direct effect (NDE)

Natural direct effect (NDE):

$$NDE(x, x'; Y) = \mathbb{E}(y \mid do(x, pa_{Y \setminus X}^{x'})) - \mathbb{E}(y \mid do(x', pa_{Y \setminus X}^{x'}))$$

where  $pa_{Y \setminus X}^{x'}$  is the value of  $Pa(Y) \setminus \{X\}$  when  $X = x'$ .

Advantage:

- It allows for the complete decomposition of the total effect into the NDE and NIE

Limitation:

- NDE cannot always be measured via experiments since it is a counterfactual expression

## Natural indirect effect (NIE)

$$TE = NDE - NIE_r$$

where  $NIE_r$  stands for the reverse transition, from  $X = 1$  to  $X = 0$ ; it becomes additive in linear systems, where reversal of transitions entails sign reversal.

## Natural indirect effect (NIE)

$$TE = NDE - NIE_r$$

where  $NIE_r$  stands for the reverse transition, from  $X = 1$  to  $X = 0$ ; it becomes additive in linear systems, where reversal of transitions entails sign reversal.

This implies that NIE is identifiable whenever NDE and the total effect are identifiable

## Natural indirect effect (NIE)

$$TE = NDE - NIE_r$$

where  $NIE_r$  stands for the reverse transition, from  $X = 1$  to  $X = 0$ ; it becomes additive in linear systems, where reversal of transitions entails sign reversal.

This implies that NIE is identifiable whenever NDE and the total effect are identifiable

Natural indirect effect (NIE):

$$NIE(x, x'; Y) = \mathbb{E}(y | do(x, pa_{Y \setminus X}^x)) - \mathbb{E}(y | do(x, pa_{Y \setminus X}^{x'}))$$


where  $pa_{Y \setminus X}^x$  is the value of  $Pa(Y) \setminus \{X\}$  when  $X = x$ .

# Mediation analysis in linear SCMs

Linearity

Causal DAG

Observational data



Mediation analysis

A linear structural causal model consists of a set of structural equations of the form:

$$y := \sum_{x \in Pa(y)} \alpha_{xy} x + \xi_y$$

where  $Pa(y)$  are direct causes of  $y$ ,  $\xi_y$  represent errors due to omitted factors and  $\alpha_{xy}$  which are known as a structural coefficient represents the strength of the causal relation.

A linear structural causal model consists of a set of structural equations of the form:

$$y := \sum_{x \in Pa(y)} \alpha_{xy} x + \xi_y$$

where  $Pa(y)$  are direct causes of  $y$ ,  $\xi_y$  represent errors due to omitted factors and  $\alpha_{xy}$  which are known as a structural coefficient represents the strength of the causal relation.

$\alpha_{xy}$  is also known as the direct effect of  $X$  on  $Y$ .

A linear structural causal model consists of a set of structural equations of the form:

$$y := \sum_{x \in Pa(y)} \alpha_{xy} x + \xi_y$$

where  $Pa(y)$  are direct causes of  $y$ ,  $\xi_y$  represent errors due to omitted factors and  $\alpha_{xy}$  which are known as a structural coefficient represents the strength of the causal relation.

$\alpha_{xy}$  is also known as the direct effect of  $X$  on  $Y$ .

The direct effect is not necessarily equivalent to a regression coefficient.

Consider a causal DAG  $\mathcal{G}$  and a direct effect  $\alpha_{xy}$  associated with link  $X \rightarrow Y$ . Let  $\mathcal{G}_{\alpha_{xy}}$  denote the graph that results when  $X \rightarrow Y$  is deleted from  $\mathcal{G}$ . A set of variables  $Z$  satisfies the **single-door criterion** iff:

- $Z$  contains no descendant of  $Y$ ;
- $Z$  d-separates  $X$  from  $Y$  in  $\mathcal{G}_{\alpha_{xy}}$ .


## Theorem

If  $\mathbb{Z}$  satisfies the single-door criterion relative to  $(X, Y)$ , then the direct effect  $\alpha_{xy}$  is identifiable and is given by

$$\alpha_{xy} = r_{XY.Z}$$


where  $r_{XY.Z}$  represents the residual correlation between  $Y$  and  $X$  after  $\mathbb{Z}$  is "partialled out".

## Single-door in action



Does the following set satisfy the single-door criterion?


## Single-door in action



Does the following set satisfy the single-door criterion?

$$\{Z\} ?$$


## Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

## Single-door in action




Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ?

## Single-door in action




Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No

## Single-door in action




Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No

$\{B\}$  ?

## Single-door in action




Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No

$\{B\}$  ? No

# Single-door in action



Does the following set satisfy the single-door criterion?


$\{Z\}$  ? No

$\{A\}$  ? No

$\{B\}$  ? No

$\{M\}$  ?

## Single-door in action



Does the following set satisfy the single-door criterion?


$\{Z\}$  ? No

$\{A\}$  ? No

$\{B\}$  ? No

$\{M\}$  ? No

# Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No


$\{A\}$  ? No

$\{B\}$  ? No

$\{M\}$  ? No

$\{A, B\}$ ?

# Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No


$\{A\}$  ? No

$\{B\}$  ? No

$\{M\}$  ? No

$\{A, B\}$ ? No

# Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No


$\{B\}$  ? No

$\{M\}$  ? No

$\{A, B\}$ ? No

$\{Z, M\}$ ?

# Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No


$\{B\}$  ? No

$\{M\}$  ? No

$\{A, B\}$ ? No

$\{Z, M\}$ ? Yes

# Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No

$\{B\}$  ? No


$\{M\}$  ? No

$\{A, B\}$ ? No

$\{Z, M\}$ ? Yes

$\{Z, M, B\}$  ?

# Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No

$\{B\}$  ? No


$\{M\}$  ? No

$\{A, B\}$ ? No

$\{Z, M\}$ ? Yes

$\{Z, M, B\}$  ? No

# Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No

$\{B\}$  ? No

$\{M\}$  ? No


$\{A, B\}$ ? No

$\{Z, M\}$ ? Yes

$\{Z, M, B\}$  ? No

$\{Z, M, A\}$ ?

# Single-door in action



Does the following set satisfy the single-door criterion?

$\{Z\}$  ? No

$\{A\}$  ? No

$\{B\}$  ? No

$\{M\}$  ? No

$\{A, B\}$ ? No

$\{Z, M\}$ ? Yes

$\{Z, M, B\}$  ? No

$\{Z, M, A\}$ ? Yes

# 4

## Counterfactuals


Structural causal models

Counterfactual reasoning


Mediation analysis

Difference with the potential outcome framework


## Structural causal models




## Structural causal models




## Potential outcome



## Structural causal models



## Potential outcome



In the structural causal models framework, the primitive assumptions becomes theorems!

5

Conclusion

# Conclusion

Counterfactual  
data


---

Experimental  
data

---

Observational  
data

---



SCMs  
counterfactual reasoning  
mediation analysis

---

Causal DAG  
causal reasoning

---

Bayesian networks  
prediction

# 6

References

## References I

- [1] Harri Helajärvi et al. “Exploring Causality between TV Viewing and Weight Change in Young and Middle-Aged Adults. The Cardiovascular Risk in Young Finns Study”. In: PloS one 9 (July 2014), e101860. doi: 10.1371/journal.pone.0101860.
- [2] M.A. Hernan and J.M. Robins. Causal Inference: What If. Chapman & Hall/CRC Monographs on Statistics & Applied Probab. CRC Press, 2023. isbn: 9781420076165.
- [3] Judea Pearl. “Causal diagrams for empirical research”. In: Biometrika 82.4 (Dec. 1995), pp. 669–688. issn: 0006-3444. doi: 10.1093/biomet/82.4.669.
- [4] Judea Pearl. Causality: Models, Reasoning, and Inference. Cambridge University Press, 2000.
- [5] Judea Pearl. “Direct and Indirect Effects”. In: Probabilistic and Causal Inference (2001).

## References II

- [6] Judea Pearl and Dana Mackenzie.  
The Book of Why: The New Science of Cause and Effect. Basic Books, 2018.
- [7] Richard Scheines. “A Woefully Incomplete History of Empirical Applications of Algorithmic Causal Discovery”. The History and Development of Search Methods for Causal Structure Workshop at the 39th Conference on Uncertainty in Artificial Intelligence. 2023.
- [8] Shohei Shimizu et al. “A Linear Non-Gaussian Acyclic Model for Causal Discovery”. In:  
Journal of Machine Learning Research 7.72 (2006),  
pp. 2003–2030.

## References III

- [9] Ilya Shpitser and Judea Pearl. “Complete Identification Methods for the Causal Hierarchy”. In: Journal of Machine Learning Research 9.64 (2008), pp. 1941–1979. url: <http://jmlr.org/papers/v9/shpitser08a.html>.
- [10] Peter Spirtes, Clark Glymour, and Richard Scheines. Causation, Prediction, and Search. 2nd. MIT press, 2000.
- [11] PETER SPIRTES et al. “Using Path Diagrams as a Structural Equation Modeling Tool”. In: Sociological Methods & Research 27.2 (1998), pp. 182–225. doi: 10.1177/0049124198027002003.
- [12] Thomas Verma and Judea Pearl. “Equivalence and synthesis of causal models”. In: UAI '90: Proceedings of the Sixth Annual Conference on Uncertainty in Artificial Intelligence. Ed. by Piero P. Bonissone et al. Elsevier, 1990, pp. 255–270.

MERCI POUR VOTRE ATTENTION

