Tight PAC-Bayes Bounds for Deep Neural Networks

Chris Kang
University of Calgary

March 20, 2026

AI Assistance Disclosure. The substantial majority of this work — including proof construction, verification, error correction, and empirical experiments — was produced by AI systems (Claude Opus 4.6, Anthropic; GPT-5.3-Codex and GPT-5.4 Pro, OpenAI; Gemini 3.1 Pro Preview and Gemini 3.1 Pro High, Google) under human direction, using multi-LLM adversarial verification across 17 review phases. Full methodology documented in the companion paper [9].

Reader's Guide. This document presents a self-contained, linearly structured proof that resolves the open problem of simultaneously achieving non-vacuous bounds, polynomial-time computability, and predictive tightness in a single PAC-Bayes framework for deep neural networks. The logical flow is as follows. We first establish monotonicity properties of the kl-inverse function (Theorems 1–3), which are foundational tools used throughout. We then state the PAC-Bayes-kl inequality (Theorem 4, cited), and build upon it to prove certificate correctness (Theorem 5), polynomial-time computability (Theorem 7), and the Pinsker-based comparison inequality for non-vacuity (Lemma 8 and Theorem 10). We then establish the rank-consistency guarantees: a Kendall τ lower bound (Theorem 12) and a direct monotonicity result (Theorem 13). Finally, the Master Theorem (Theorem 14) assembles all preceding results to simultaneously discharge all three criteria. A brief empirical summary follows.

1 Open Problem

Open Problem (solveall.org). Does there exist a single PAC-Bayes framework that *simultaneously* achieves:

1. Non-vacuous bounds ($B < 1$) at practical scale (ImageNet-scale networks),
2. Polynomial-time certified computation of the bound,
3. Strong predictive tightness: the bound ranking across hyperparameters agrees with true risk ranking?

Prior work achieved these criteria in isolation or pairwise, but no unified framework was known to satisfy all three simultaneously with rigorous theoretical guarantees and empirical verification at practical scale.

Our answer is yes. The framework uses:

1. The **PAC-Bayes-kl inequality** with Maurer's constant $C(n, \delta) = \ln(2\sqrt{n}/\delta)$,
2. **kl-inverse certificate computation** via certified bisection with explicit numerical error tracking,
3. **Universality** over all admissible prior-posterior pairs (P, Q) with $\mathrm{KL}(Q\|P) < \infty$.

Validity is universal over all such pairs; polynomial-time computability (Theorem 7) additionally requires that a certified upper bound $\mathrm{KL}_{ub}(Q) \geq \mathrm{KL}(Q\|P)$ be polynomial-time computable from the representation of (P, Q).

2 Setup and Notation

Let $\mathcal{X}$ be an input space, $\mathcal{Y} = \{1, \ldots, C\}$ a finite label set, and $\mathcal{D}$ an unknown distribution on $\mathcal{X} \times \mathcal{Y}$. A training sample $S = ((x_1, y_1), \ldots, (x_n, y_n)) \sim \mathcal{D}^n$ is split as $S = S_0 \cup S_1$ with $|S_1| = n_1$; the prior and posterior are constructed from S_0, while S_1 serves as the independent evaluation sample for PAC-Bayes bounds. Let $h_\theta : \mathcal{X} \to \mathcal{Y}$ denote the predictor parametrized by $\theta \in \Theta$, where $\Theta \subseteq \mathbb{R}^d$ is a d-dimensional parameter space. We consider the Gibbs predictor induced by posterior Q on Θ with bounded 0-1 loss $\ell : \mathcal{Y} \times \mathcal{Y} \to [0, 1]$.

Key quantities:

$$L_{\mathcal{D}}(Q) := \mathbb{E}_{\theta\sim Q}[\mathbb{E}_{(x,y)\sim\mathcal{D}}[\ell(h_\theta(x), y)]], \tag{1}$$

$$\hat{L}_S(Q) := \mathbb{E}_{\theta\sim Q}\left[\frac{1}{n}\sum_{i=1}^{n} \ell(h_\theta(x_i), y_i)\right]. \tag{2}$$

When evaluated on the evaluation subset S_1 of size n_1, we write $\hat{L}_{S_1}(Q)$ with the sum taken over S_1 and denominator n_1. For a single parameter θ, we write $\hat{L}_{S_1}(h_\theta) := \frac{1}{n_1}\sum_{i\in S_1} \ell(h_\theta(x_i), y_i)$.

The binary KL divergence is $\mathrm{kl}(p\|q) = p\ln(p/q) + (1-p)\ln((1-p)/(1-q))$ (with the standard convention $0\ln 0 := 0$), and the kl-inverse is

$$\mathrm{kl}_+^{-1}(p, \varepsilon) := \max\{q \in [p, 1] : \mathrm{kl}(p\|q) \leq \varepsilon\}.$$

Certificate parameters. The certified bound (Theorem 5) requires several error-tolerance and approximation parameters beyond the standard PAC-Bayes setup:

- $\mathrm{KL}_{ub}(Q) \geq \mathrm{KL}(Q\|P)$: a certified upper bound on the KL divergence, assumed polynomial-time computable from the representation of (P, Q).
- $\hat{L}_{S_1}^{(m)}(Q) := \frac{1}{m}\sum_{j=1}^{m} \hat{L}_{S_1}(h_{\theta_j})$ with $\theta_j \overset{\text{iid}}{\sim} Q$: Monte Carlo estimate of $\hat{L}_{S_1}(Q)$.
- $\varepsilon_{mc} \geq 0$: Monte Carlo tolerance; by Hoeffding's inequality, $|\hat{L}_{S_1}^{(m)}(Q) - \hat{L}_{S_1}(Q)| \leq \varepsilon_{mc}$ with probability $\geq 1 - \delta_{mc}$ when $m \geq \ln(2/\delta_{mc})/(2\varepsilon_{mc}^2)$.
- $\eta_{num} \geq 0$: bisection bracket tolerance for numerical computation of kl_+^{-1}.
- $C(n, \delta) := \ln(2\sqrt{n}/\delta)$: the Maurer constant [1].
- $\delta = \delta_{pb} + \delta_{mc} + \delta_{num}$: total confidence parameter, decomposed into PAC-Bayes (δ_{pb}), Monte Carlo (δ_{mc}), and numerical (δ_{num}) failure budgets.

3 Monotonicity Properties of kl_+^{-1}

These monotonicity lemmas are foundational to the proofs of the certificate correctness, comparison, and direct monotonicity theorems.

Theorem 1 (Monotonicity of $\mathrm{kl}(p\|q)$ in q). *For fixed $p \in [0, 1)$, the map $q \mapsto \mathrm{kl}(p\|q)$ is continuous and strictly increasing on $[p, 1)$, with $\mathrm{kl}(p\|p) = 0$ and $\lim_{q\uparrow 1} \mathrm{kl}(p\|q) = +\infty$.*

Proof. For $p \in (0,1)$ and $q \in (p,1)$:

$$\frac{\partial}{\partial q}\mathrm{kl}(p\|q) = -\frac{p}{q} + \frac{1-p}{1-q}.$$

Since $q > p$, we have $p/q < 1$ and $(1-p)/(1-q) > 1$, so the derivative is positive. Hence $q \mapsto \mathrm{kl}(p\|q)$ is strictly increasing on $(p,1)$. Continuity is clear from the definition. For the boundary: $\mathrm{kl}(p\|p) = p\ln 1 + (1-p)\ln 1 = 0$, and $\lim_{q\uparrow 1}(1-p)\ln\frac{1-p}{1-q} = +\infty$ since $p < 1$. The case $p = 0$ gives $\mathrm{kl}(0\|q) = -\ln(1-q)$, which is strictly increasing on $[0,1)$ with limit $+\infty$. □

Theorem 2 (Monotonicity of kl_+^{-1} in ε)**.** *For fixed $p \in [0,1]$, if $0 \le \varepsilon \le \varepsilon'$, then $\mathrm{kl}_+^{-1}(p,\varepsilon) \le \mathrm{kl}_+^{-1}(p,\varepsilon')$. The inequality is strict when $p < 1$ and $\varepsilon < \varepsilon'$.*

Proof. Weak monotonicity. The feasible set $\{q \in [p,1] : \mathrm{kl}(p\|q) \le \varepsilon\}$ is contained in $\{q \in [p,1] : \mathrm{kl}(p\|q) \le \varepsilon'\}$, so the supremum over the larger set is at least as large.

Strict monotonicity. Suppose $p < 1$ and $\varepsilon < \varepsilon'$. Let $q^* = \mathrm{kl}_+^{-1}(p,\varepsilon)$. By Theorem 1, $\mathrm{kl}(p\|\cdot)$ is continuous and strictly increasing on $[p,1)$ with $\lim_{q\uparrow 1}\mathrm{kl}(p\|q) = +\infty$. Since $\mathrm{kl}(p\|\cdot)$ is continuous and strictly increasing on $[p,1)$ with value 0 at $q = p$ and $\mathrm{kl}(p\|q^*) \le \varepsilon$, the maximum in the definition of kl_+^{-1} is attained: $\mathrm{kl}(p\|q^*) = \varepsilon$. Thus $\mathrm{kl}(p\|q^*) = \varepsilon < \varepsilon'$, and there exists $q' > q^*$ with $\mathrm{kl}(p\|q') = \varepsilon'$ (by the intermediate value theorem). Hence $\mathrm{kl}_+^{-1}(p,\varepsilon') \ge q' > q^* = \mathrm{kl}_+^{-1}(p,\varepsilon)$. □

Theorem 3 (Monotonicity of kl_+^{-1} in p)**.** *For fixed $\varepsilon \ge 0$, if $0 \le p \le p' \le 1$, then $\mathrm{kl}_+^{-1}(p,\varepsilon) \le \mathrm{kl}_+^{-1}(p',\varepsilon)$. The inequality is strict when $\varepsilon > 0$ and $p < p'$.*

Proof. Weak monotonicity. Let $q^* = \mathrm{kl}_+^{-1}(p,\varepsilon)$. We consider two cases.

Case 1: $q^* < p'$. Then $q^* \le p' \le \mathrm{kl}_+^{-1}(p',\varepsilon)$, since $\mathrm{kl}_+^{-1}(p',\varepsilon) \ge p'$ by definition.

Case 2: $q^* \ge p'$. For $q \ge p' \ge p$, we claim $\mathrm{kl}(p'\|q) \le \mathrm{kl}(p\|q)$. Indeed, the partial derivative $\frac{\partial}{\partial p}\mathrm{kl}(p\|q) = \ln\frac{p(1-q)}{q(1-p)}$, which is negative when $p < q$ (since $p(1-q) < q(1-p)$ iff $p < q$). Since $p \le p' \le q^*$, we have $\mathrm{kl}(p'\|q^*) \le \mathrm{kl}(p\|q^*) \le \varepsilon$. So q^* is feasible for (p',ε), giving $\mathrm{kl}_+^{-1}(p',\varepsilon) \ge q^*$.

Strict monotonicity. Suppose $\varepsilon > 0$ and $p < p'$. If $p' = 1$, then $\mathrm{kl}_+^{-1}(1,\varepsilon) = 1$ while $\mathrm{kl}_+^{-1}(p,\varepsilon) < 1$ (since $\mathrm{kl}(p\|q) \to +\infty$ as $q \uparrow 1$ for $p < 1$), so the inequality is strict. Now suppose $p' < 1$. In Case 1, $\mathrm{kl}_+^{-1}(p',\varepsilon) \ge p' > q^*$, which is already strict. In Case 2, the derivative $\frac{\partial}{\partial p}\mathrm{kl}(p\|q) = \ln\frac{p(1-q)}{q(1-p)} < 0$ strictly for $p < q$, so $\mathrm{kl}(p'\|q^*) < \mathrm{kl}(p\|q^*) \le \varepsilon$. Thus q^* lies in the interior of the feasible set for (p',ε), and $\mathrm{kl}_+^{-1}(p',\varepsilon) > q^*$. □

4 PAC-Bayes-kl Inequality

Theorem 4 (PAC-Bayes-kl, Maurer variant [1])**.** *Let P be a prior on Θ independent of $S_1 \sim \mathcal{D}^{n_1}$, with $n_1 \ge 8$. For any $\delta \in (0,1)$:*

$$\mathbb{P}_{S_1}\left[\forall Q : \mathrm{kl}\Big(\hat{L}_{S_1}(Q) \,\|\, L_{\mathcal{D}}(Q)\Big) \le \frac{\mathrm{KL}(Q\|P) + \ln(2\sqrt{n_1}/\delta)}{n_1}\right] \ge 1-\delta.$$

Inverting the kl *constraint via* kl_+^{-1} *yields an explicit upper bound on* $L_{\mathcal{D}}(Q)$.

This is a known result; we cite Maurer [1] and do not reproduce the proof.

5 Certificate Correctness

Theorem 5 (Certified kl-inverse bound)**.** *Assume $n_1 \ge 8$ (as required by Theorem 4). Fix a posterior Q. Define the clamped empirical loss and certified PAC-Bayes slack:*

$$p_{\mathrm{cert}} := \min\Big\{1,\, \hat{L}_{S_1}^{(m)}(Q) + \varepsilon_{mc}\Big\}, \qquad \varepsilon_{pb}^{\mathrm{cert}}(Q) := \frac{\mathrm{KL}_{ub}(Q) + C(n_1,\delta_{pb})}{n_1}.$$

The certified bound is:

$$B_{\text{cert}}(Q) := \text{kl}_+^{-1}\big(p_{\text{cert}},\ \varepsilon_{pb}^{\text{cert}}(Q)\big) + \eta_{num}.$$

Then with probability $\geq 1-\delta$ (over S_1 and any auxiliary randomness, conditional on S_0):

$$L_{\mathcal{D}}(Q) \leq B_{\text{cert}}(Q).$$

Proof. We work conditional on the realized S_0 (and hence on $P = P(S_0)$), with probability over $S_1 \sim \mathcal{D}^{n_1}$, Monte Carlo draws $\theta_{1:m} \sim Q^m$, and any numerical randomness. Let

$$\varepsilon_{pb}(Q) := \frac{\text{KL}(Q\|P) + C(n_1, \delta_{pb})}{n_1}$$

denote the true PAC-Bayes slack (using the exact KL, as opposed to $\varepsilon_{pb}^{\text{cert}}(Q)$ which uses the upper bound KL_{ub}). Let $B_{\text{cert}}^{num}(Q)$ denote the numerical output of the bisection routine for $\text{kl}_+^{-1}(p_{\text{cert}}, \varepsilon_{pb}^{\text{cert}}(Q))$, satisfying

$$\text{kl}_+^{-1}(p_{\text{cert}}, \varepsilon_{pb}^{\text{cert}}(Q)) \leq B_{\text{cert}}^{num}(Q) \leq \text{kl}_+^{-1}(p_{\text{cert}}, \varepsilon_{pb}^{\text{cert}}(Q)) + \eta_{num} = B_{\text{cert}}(Q).$$

Define three events:

- E_{pb}: the PAC-Bayes-kl inequality (Theorem 4) holds. Failure probability $\leq \delta_{pb}$.
- E_{mc}: the Monte Carlo concentration bound $|\hat{L}_{S_1}^{(m)}(Q) - \hat{L}_{S_1}(Q)| \leq \varepsilon_{mc}$ holds. Failure probability $\leq \delta_{mc}$ (Hoeffding, since $\hat{L}_{S_1}(h_{\theta_j}) \in [0,1]$).
- E_{num}: $\text{KL}(Q\|P) \leq \text{KL}_{ub}(Q)$ and the bisection returns $B_{\text{cert}}^{num}(Q)$ as above. Failure probability $\leq \delta_{num}$.

Step 1 (PAC-Bayes bound). On E_{pb}:

$$L_{\mathcal{D}}(Q) \leq \text{kl}_+^{-1}\Big(\hat{L}_{S_1}(Q),\ \varepsilon_{pb}(Q)\Big).$$

Step 2 (KL monotonicity). On E_{num}, $\text{KL}(Q\|P) \leq \text{KL}_{ub}(Q)$, so $\varepsilon_{pb}(Q) \leq \varepsilon_{pb}^{\text{cert}}(Q)$. By Theorem 2 (monotonicity in ε):

$$\text{kl}_+^{-1}\Big(\hat{L}_{S_1}(Q),\ \varepsilon_{pb}(Q)\Big) \leq \text{kl}_+^{-1}\Big(\hat{L}_{S_1}(Q),\ \varepsilon_{pb}^{\text{cert}}(Q)\Big).$$

Step 3 (MC monotonicity). On E_{mc}, $\hat{L}_{S_1}(Q) \leq \hat{L}_{S_1}^{(m)}(Q) + \varepsilon_{mc}$. Since $\hat{L}_{S_1}(Q) \in [0,1]$, we have $\hat{L}_{S_1}(Q) \leq \min\{1, \hat{L}_{S_1}^{(m)}(Q) + \varepsilon_{mc}\} = p_{\text{cert}}$. By Theorem 3 (monotonicity in p):

$$\text{kl}_+^{-1}\Big(\hat{L}_{S_1}(Q),\ \varepsilon_{pb}^{\text{cert}}(Q)\Big) \leq \text{kl}_+^{-1}\big(p_{\text{cert}},\ \varepsilon_{pb}^{\text{cert}}(Q)\big).$$

Step 4 (Numerical bracket). On E_{num}:

$$\text{kl}_+^{-1}\big(p_{\text{cert}},\ \varepsilon_{pb}^{\text{cert}}(Q)\big) \leq B_{\text{cert}}^{num}(Q) \leq B_{\text{cert}}(Q),$$

where the second inequality holds because bisection returns an upper bracket endpoint satisfying $B_{\text{cert}}^{num}(Q) \leq \text{kl}_+^{-1}(p_{\text{cert}}, \varepsilon_{pb}^{\text{cert}}(Q)) + \eta_{num} = B_{\text{cert}}(Q)$.

Combining Steps 1–4: $L_{\mathcal{D}}(Q) \leq B_{\text{cert}}(Q)$ on $E_{pb} \cap E_{mc} \cap E_{num}$. By the union bound:

$$\mathbb{P}(E_{pb}^c \cup E_{mc}^c \cup E_{num}^c \mid S_0) \leq \delta_{pb} + \delta_{mc} + \delta_{num} = \delta,$$

where the E_{mc} bound uses the tower property: $\mathbb{P}(E_{mc}^c \mid S_0) = \mathbb{E}[\mathbb{P}(E_{mc}^c \mid S_1, Q) \mid S_0] \leq \delta_{mc}$, since $\mathbb{P}(E_{mc}^c \mid S_1, Q) \leq 2e^{-2m\varepsilon_{mc}^2} \leq \delta_{mc}$ by Hoeffding's inequality and the choice $m \geq \ln(2/\delta_{mc})/(2\varepsilon_{mc}^2)$. □

Remark 6 (Analytic envelope vs. numerical output)**.** *The certificate $B_{\text{cert}}(Q)$ as defined above is an analytic quantity involving the exact kl_+^{-1}. In practice, the bisection routine returns a numerical value $B_{\text{cert}}^{num}(Q)$ satisfying $\text{kl}_+^{-1}(p_{\text{cert}}, \varepsilon_{pb}^{\text{cert}}(Q)) \leq B_{\text{cert}}^{num}(Q) \leq B_{\text{cert}}(Q)$, so $L_{\mathcal{D}}(Q) \leq B_{\text{cert}}^{num}(Q) \leq B_{\text{cert}}(Q)$ on the same event. All ranking statements (Theorems 12 and 13) are stated for B_{cert} and transfer to B_{cert}^{num} whenever the ordering margin exceeds η_{num}.*

6 Polynomial-Time Computability

Theorem 7 (Polynomial-time certificate)**.** *Let T_{fwd} denote the cost of evaluating one weight sample h_θ on the full evaluation dataset S_1 (i.e., n_1 forward passes), and let T_{KL} denote the cost of computing $\text{KL}_{ub}(Q)$. For fixed posterior Q, a value $B^{num}_{\text{cert}}(Q) \le B_{\text{cert}}(Q)$ satisfying $L_{\mathcal{D}}(Q) \le B^{num}_{\text{cert}}(Q)$ on the high-probability event is computable in time*

$$O(T_{\text{KL}} + m \cdot T_{\text{fwd}} + \log_2(1/\eta_{num}))\,,$$

where m is the Monte Carlo sample count. For a finite hyperparameter grid $|\Lambda| = M$ with union-bound confidence:

$$O(M \cdot T_{\text{KL}} + M \cdot m \cdot T_{\text{fwd}} + M \cdot \log_2(1/\eta_{num}))\,.$$

Both are polynomial in $(n_1, d, M, \log(1/\delta), 1/\varepsilon_{mc}, \log(1/\eta_{num}))$ when T_{KL} and T_{fwd} are polynomial in (d, n_1) (as assumed in (A5)–(A6) of the Master Theorem).

Proof. The numerical certificate $B^{num}_{\text{cert}}(Q) \le B_{\text{cert}}(Q)$ for a single posterior Q is computed by the following three-phase procedure:

Phase 1: KL computation. Compute $\text{KL}_{ub}(Q) \ge \text{KL}(Q\|P)$. For the isotropic Gaussian construction $P = \mathcal{N}(w_{\text{early}}, \sigma^2 I)$, $Q = \mathcal{N}(w_{\text{final}}, \sigma^2 I)$, this is $\|w_{\text{final}} - w_{\text{early}}\|^2/(2\sigma^2)$: a single $O(d)$ operation.

Phase 2: Monte Carlo estimation. Draw m i.i.d. samples $\theta_1, \ldots, \theta_m \sim Q$ and compute $\hat{L}^{(m)}_{S_1}(Q) = \frac{1}{m}\sum_{j=1}^{m} \hat{L}_{S_1}(h_{\theta_j})$. Each $\hat{L}_{S_1}(h_{\theta_j})$ requires a full pass over S_1 at cost T_{fwd}. Total: $O(m{\cdot}T_{\text{fwd}})$. The Hoeffding requirement $m \ge \ln(2/\delta_{mc})/(2\varepsilon^2_{mc})$ is polynomial in $(1/\varepsilon_{mc}, \log(1/\delta_{mc}))$.

Phase 3: kl-inverse bisection. Compute $\text{kl}^{-1}_{+}(p_{\text{cert}}, \varepsilon^{\text{cert}}_{pb})$ by bisection on $[p_{\text{cert}}, 1]$. Each bisection step evaluates $\text{kl}(p_{\text{cert}}\|q_{\text{mid}})$, which is $O(1)$. To achieve bracket width η_{num}, bisection requires $\lceil \log_2(1/\eta_{num})\rceil$ steps. Total: $O(\log_2(1/\eta_{num}))$.

Combining: $O(T_{\text{KL}} + m \cdot T_{\text{fwd}} + \log_2(1/\eta_{num}))$, polynomial in all parameters.

Extension to finite grids. For $|\Lambda| = M$ candidates, apply the above to each $\lambda \in \Lambda$ with per-candidate confidence δ_{mc}/M and δ_{num}/M (union bound; the PAC-Bayes event E_{pb} is shared across all λ). Total runtime: $O(M \cdot T_{\text{KL}} + M \cdot m \cdot T_{\text{fwd}} + M \cdot \log_2(1/\eta_{num}))$.

Remark (extension to compact continuous Λ). The finite-grid result extends to compact $\Lambda \subset \mathbb{R}^d$ via standard covering-number arguments, under additional regularity. Assuming $\lambda \mapsto \text{KL}(Q_\lambda\|P)$ is L_K-Lipschitz, $\lambda \mapsto \hat{L}_{S_1}(Q_\lambda)$ (the exact Gibbs loss, i.e., $\varepsilon_{mc} = 0$) is L_L-Lipschitz, and $p_{\text{cert}}(\lambda) < 1$ for all $\lambda \in \Lambda$, the certificate map $\lambda \mapsto B_{\text{cert}}(\lambda)$ is Lipschitz. The derivative bounds follow from $\varepsilon_{\text{cert}}(\lambda) \ge C(n_1, \delta_{pb})/n_1 > 0$ (ensuring kl^{-1}_{+} is evaluated away from its singularity at $\varepsilon = 0$) and $p_{\text{cert}} < 1$ (ensuring $q^* < 1$). Denoting $R := \text{diam}(\Lambda)$ and L_B the Lipschitz constant of $\lambda \mapsto B_{\text{cert}}(\lambda)$, an ε_{net}-net of size $M \le (2RL_B/\varepsilon_{\text{approx}})^d$ then covers Λ to accuracy $\varepsilon_{\text{approx}}$, yielding polynomial total runtime for fixed d. A full proof of this extension is omitted; the main results (Theorems 5–14) use only finite grids. □

7 Comparison Theorem and Non-Vacuity

We first establish Pinsker's inequality, then the Maurer constant dominance lemma, and finally the comparison chain.

Lemma 8 (Pinsker's inequality [5])**.** *For all $p, q \in [0, 1]$:* $\text{kl}(p\|q) \ge 2(p - q)^2$.

Proof. For fixed $q \in (0, 1)$, define $f(p) = \text{kl}(p\|q)$ for $p \in (0, 1)$. Then $f(q) = 0$, $f'(q) = [\ln(p/q) - \ln((1-p)/(1-q))]_{p=q} = 0$, and

$$f''(p) = \frac{1}{p} + \frac{1}{1-p} = \frac{1}{p(1-p)}.$$

By AM-GM, $p(1-p) \le 1/4$, so $f''(p) \ge 4$ everywhere on $(0,1)$. Since f is convex with $f(q) = f'(q) = 0$ and $f''(p) \ge 4$:

$$f(p) \ge \tfrac{1}{2} \cdot 4 \cdot (p-q)^2 = 2(p-q)^2.$$

Extension to boundaries by continuity. □

Lemma 9 (Maurer constant dominance)**.** *For all $n \ge 1$ and $\delta \in (0,1)$:*

$$C_{\mathrm{Ma}}(n,\delta) := \ln \frac{2\sqrt{n}}{\delta} \le \ln \frac{2n}{\delta} =: C_{\mathrm{Mc}}(n,\delta).$$

Equality holds iff $n = 1$.

Proof. $C_{\mathrm{Mc}} - C_{\mathrm{Ma}} = \ln \frac{2n/\delta}{2\sqrt{n}/\delta} = \ln \sqrt{n} = \frac{1}{2} \ln n \ge 0$, with equality iff $n = 1$. □

Bound variants. We compare four certificate variants arising from two choices of constant and two inversion methods. With C_{Ma} and C_{Mc} as above, define

$$\varepsilon_{\mathrm{Ma}} := \frac{\mathrm{KL}(Q\|P) + C_{\mathrm{Ma}}(n_1,\delta)}{n_1}, \qquad \varepsilon_{\mathrm{Mc}} := \frac{\mathrm{KL}(Q\|P) + C_{\mathrm{Mc}}(n_1,\delta)}{n_1}.$$

The four bounds are:

$$B_{\mathrm{kl}}^{\mathrm{Ma}} := \mathrm{kl}_+^{-1}(\hat{L}_{S_1}(Q),\ \varepsilon_{\mathrm{Ma}}), \qquad B_{\surd}^{\mathrm{Ma}} := \hat{L}_{S_1}(Q) + \sqrt{\varepsilon_{\mathrm{Ma}}/2},$$
$$B_{\mathrm{kl}}^{\mathrm{Mc}} := \mathrm{kl}_+^{-1}(\hat{L}_{S_1}(Q),\ \varepsilon_{\mathrm{Mc}}), \qquad B_{\surd}^{\mathrm{Mc}} := \hat{L}_{S_1}(Q) + \sqrt{\varepsilon_{\mathrm{Mc}}/2}.$$

Theorem 10 (Comparison theorem and non-vacuity)**.** *For any $p \in [0,1]$ and $\varepsilon \ge 0$:*

$$\mathrm{kl}_+^{-1}(p,\varepsilon) \le p + \sqrt{\varepsilon/2},$$

with equality iff $\varepsilon = 0$. Consequently, for $\varepsilon > 0$ the kl-inversion bound is strictly tighter than the square-root relaxation:

$$B_{\mathrm{kl}}^{\mathrm{Ma}} \le B_{\surd}^{\mathrm{Ma}} \le B_{\surd}^{\mathrm{Mc}},$$

with strict inequalities for $n_1 \ge 2$ (the first by kl-inversion vs. Pinsker, the second by Maurer vs. McAllester constants; see Lemma 9).

Remark 11 (Empirical non-vacuity)**.** *Non-vacuity ($B_{\mathrm{kl}}^{\mathrm{Ma}} < 1$) is established empirically in Section 11: all 32 ImageNet configurations and all 10 MNIST configurations yield $B_{\mathrm{kl}}^{\mathrm{Ma}} < 1$, with the tightest certificate $B = 0.239$ (EfficientNet-B0).*

Proof. Step 1 (kl-inversion Pinsker bound). We show $\mathrm{kl}_+^{-1}(p,\varepsilon) \le p + \sqrt{\varepsilon/2}$ for all $p \in [0,1]$ and $\varepsilon \ge 0$.

Let $q^* = \mathrm{kl}_+^{-1}(p,\varepsilon)$, so $\mathrm{kl}(p\|q^*) \le \varepsilon$. By Pinsker's inequality (Lemma 8), $\mathrm{kl}(p\|q^*) \ge 2(p-q^*)^2$. Since $q^* \ge p$:

$$\varepsilon \ge \mathrm{kl}(p\|q^*) \ge 2(q^*-p)^2, \quad \text{so} \quad q^* - p \le \sqrt{\varepsilon/2}, \quad \text{giving} \quad q^* \le p + \sqrt{\varepsilon/2}.$$

Equality analysis. If $\varepsilon = 0$, then $\mathrm{kl}(p\|q^*) = 0$ forces $q^* = p$, giving equality. For $\varepsilon > 0$ and $p < 1$: Pinsker's inequality is strict when $q^* > p$ (since $\mathrm{kl}(p\|q) > 2(p-q)^2$ for $q \ne p$ with $p \in (0,1)$), so the bound is strict. The boundary case $p = 1$: $\mathrm{kl}_+^{-1}(1,\varepsilon) = 1 < 1 + \sqrt{\varepsilon/2}$ for $\varepsilon > 0$.

Step 2 (Comparison chain). Define:

$$\varepsilon_{\mathrm{Ma}} = \frac{\mathrm{KL}(Q\|P) + C_{\mathrm{Ma}}(n_1,\delta)}{n_1}, \qquad \varepsilon_{\mathrm{Mc}} = \frac{\mathrm{KL}(Q\|P) + C_{\mathrm{Mc}}(n_1,\delta)}{n_1}.$$

By Lemma 9, $C_{\mathrm{Ma}} \le C_{\mathrm{Mc}}$, so $\varepsilon_{\mathrm{Ma}} \le \varepsilon_{\mathrm{Mc}}$. By Theorem 2:

$$B_{\mathrm{kl}}^{\mathrm{Ma}} = \mathrm{kl}_+^{-1}(\hat{L}, \varepsilon_{\mathrm{Ma}}) \le \mathrm{kl}_+^{-1}(\hat{L}, \varepsilon_{\mathrm{Mc}}) = B_{\mathrm{kl}}^{\mathrm{Mc}}.$$

By Step 1 (Pinsker bound applied to $B_{\mathrm{kl}}^{\mathrm{Mc}}$):

$$B_{\mathrm{kl}}^{\mathrm{Mc}} = \mathrm{kl}_+^{-1}(\hat{L}, \varepsilon_{\mathrm{Mc}}) \le \hat{L} + \sqrt{\varepsilon_{\mathrm{Mc}}/2} = B_{\surd}^{\mathrm{Mc}}.$$

Applying Step 1 with Maurer's constant: $B_{\mathrm{kl}}^{\mathrm{Ma}} = \mathrm{kl}_+^{-1}(\hat{L}, \varepsilon_{\mathrm{Ma}}) \le \hat{L} + \sqrt{\varepsilon_{\mathrm{Ma}}/2} = B_{\surd}^{\mathrm{Ma}}$. Since $\varepsilon_{\mathrm{Ma}} \le \varepsilon_{\mathrm{Mc}}$: $B_{\surd}^{\mathrm{Ma}} = \hat{L} + \sqrt{\varepsilon_{\mathrm{Ma}}/2} \le \hat{L} + \sqrt{\varepsilon_{\mathrm{Mc}}/2} = B_{\surd}^{\mathrm{Mc}}$, with strict inequality for $n_1 \ge 2$. Hence the full chain:

$$B_{\mathrm{kl}}^{\mathrm{Ma}} \le B_{\surd}^{\mathrm{Ma}} \le B_{\surd}^{\mathrm{Mc}}.$$

Step 3 (Non-vacuity). The comparison chain is a purely analytic inequality; non-vacuity ($B_{\mathrm{kl}}^{\mathrm{Ma}} < 1$) requires a concrete witness (P, Q, S, δ) for which the right-hand side is < 1. Our own experiments (Section 11) provide this witness: using data-dependent Gaussian priors with sample splitting on ImageNet, all 32 configurations yield $B_{\mathrm{kl}}^{\mathrm{Ma}} < 1$, with the tightest certificate $B = 0.239$ for EfficientNet-B0 (gap of 1.6 percentage points above true risk). These results confirm non-vacuity at ImageNet scale within the present framework. □

8 Kendall τ Lower Bound

Theorem 12 (Kendall τ lower bound)**.** *Let $\Lambda = \{\lambda_1, \dots, \lambda_M\}$ be a finite hyperparameter grid with $M \ge 2$, and let $\delta = \delta_{pb} + \delta_{mc} + \delta_{num}$. Define the high-probability event*

$$\mathcal{E} := \big\{\forall \lambda \in \Lambda : \; L_{\mathcal{D}}(Q_\lambda) \le B_{\mathrm{cert}}(Q_\lambda)\big\},$$

which satisfies $\Pr[\mathcal{E} \mid S_0] \ge 1 - M\delta$ by Theorem 5 and the union bound (probability over S_1 and any internal Monte Carlo randomness, conditional on S_0). Suppose the true risks $\{L_{\mathcal{D}}(Q_\lambda)\}_{\lambda \in \Lambda}$ are distinct and sorted with minimum gap δ_L. Define the maximum certification gap

$$\mathrm{gap}_{\max} := \max_{\lambda \in \Lambda} \big[B_{\mathrm{cert}}(Q_\lambda) - L_{\mathcal{D}}(Q_\lambda)\big],$$

$k_{\max} = \lfloor \mathrm{gap}_{\max}/\delta_L \rfloor$, and $\tilde{k} = \min(k_{\max}, M-1)$. On $\mathcal{E}$, Kendall's $\tau_a :=$ (concordant − discordant)$/\binom{M}{2}$ satisfies:

$$\tau_a \ge 1 - \frac{2\tilde{k}(2M - \tilde{k} - 1)}{M(M-1)}.$$

The domain guard $\tilde{k}$ ensures the bound remains in $[-1, 1]$ when $k_{\max} \ge M$. Full rank consistency ($\tau_a = 1$) holds when $\mathrm{gap}_{\max} < \delta_L$.

Proof. Setup. Suppose the true risks are distinct and sorted: $L_{\mathcal{D}}(Q_{\lambda_1}) < \cdots < L_{\mathcal{D}}(Q_{\lambda_M})$, with minimum consecutive gap $\delta_L := \min_{i=1,\dots,M-1}(L_{\mathcal{D}}(Q_{\lambda_{i+1}}) - L_{\mathcal{D}}(Q_{\lambda_i}))$. Define the certification gap $\mathrm{gap}_i := B_{\mathrm{cert}}(Q_{\lambda_i}) - L_{\mathcal{D}}(Q_{\lambda_i}) \ge 0$ (non-negative on $\mathcal{E}$), and $\mathrm{gap}_{\max} := \max_i \mathrm{gap}_i$.

Step 1 (Concordance for well-separated pairs). Consider a pair (i, j) with $j > i$ and $j - i \ge k_{\max} + 1$, where $k_{\max} = \lfloor \mathrm{gap}_{\max}/\delta_L \rfloor$. The true-risk gap satisfies:

$$L_{\mathcal{D}}(Q_{\lambda_j}) - L_{\mathcal{D}}(Q_{\lambda_i}) \ge (j - i)\delta_L \ge (k_{\max} + 1)\delta_L > \mathrm{gap}_{\max},$$

where the last inequality uses $k_{\max} = \lfloor \mathrm{gap}_{\max}/\delta_L \rfloor$, so $(k_{\max} + 1)\delta_L > \mathrm{gap}_{\max}$.

We need to show that $B_{\mathrm{cert}}(Q_{\lambda_j}) > B_{\mathrm{cert}}(Q_{\lambda_i})$, i.e., the pair is concordant. Write:

$$B_{\mathrm{cert}}(Q_{\lambda_j}) - B_{\mathrm{cert}}(Q_{\lambda_i}) = \underbrace{\big(L_{\mathcal{D}}(Q_{\lambda_j}) - L_{\mathcal{D}}(Q_{\lambda_i})\big)}_{>\mathrm{gap}_{\max}} + \underbrace{\big(\mathrm{gap}_j - \mathrm{gap}_i\big)}_{\ge -\mathrm{gap}_{\max}}.$$

The first term exceeds $\text{gap}_{\max}$. For the second: $\text{gap}_j \geq 0$ and $\text{gap}_i \leq \text{gap}_{\max}$, so $\text{gap}_j - \text{gap}_i \geq -\text{gap}_{\max}$. Therefore:

$$B_{\text{cert}}(Q_{\lambda_j}) - B_{\text{cert}}(Q_{\lambda_i}) > \text{gap}_{\max} - \text{gap}_{\max} = 0.$$

The pair (i, j) is concordant.

Step 2 (Counting at-risk pairs). Pairs with $|i - j| \leq k_{\max}$ are not guaranteed concordant. Apply the domain guard: set $\tilde{k} = \min(k_{\max}, M - 1)$ to ensure the count stays within the valid range. The number of at-risk pairs (pessimistically all discordant) is:

$$\sum_{k=1}^{\tilde{k}} (M - k) = \tilde{k}M - \sum_{k=1}^{\tilde{k}} k = \tilde{k}M - \frac{\tilde{k}(\tilde{k}+1)}{2} = \frac{\tilde{k}(2M - \tilde{k} - 1)}{2}.$$

Step 3 (Kendall τ bound). The total number of pairs is $\binom{M}{2} = M(M-1)/2$. Kendall's τ is defined as $\tau = (\text{concordant} - \text{discordant})/\binom{M}{2}$. With at most $\tilde{k}(2M - \tilde{k} - 1)/2$ discordant pairs and at least $M(M-1)/2 - \tilde{k}(2M - \tilde{k} - 1)/2$ concordant pairs:

$$\tau \geq \frac{\left[M(M-1)/2 - \tilde{k}(2M - \tilde{k} - 1)/2\right] - \tilde{k}(2M - \tilde{k} - 1)/2}{M(M-1)/2} = 1 - \frac{2\tilde{k}(2M - \tilde{k} - 1)}{M(M-1)}.$$

Full rank consistency. When $\text{gap}_{\max} < \delta_L$, we have $k_{\max} = \lfloor \text{gap}_{\max}/\delta_L \rfloor = 0$, so $\tilde{k} = 0$ and $\tau \geq 1$. Since $\tau \leq 1$ always, this gives $\tau = 1$.

Domain guard. When $k_{\max} \geq M$, the clamp $\tilde{k} = M - 1$ ensures all $\binom{M}{2}$ pairs are at risk, giving $\tau \geq 1 - \frac{2(M-1)(2M-M)}{M(M-1)} = 1 - 2 = -1$, which is the trivially valid lower bound. □

9 Direct Monotonicity

Theorem 13 (Direct monotonicity)**.** *For any λ, λ' sharing the same ε_{mc}: if $\hat{L}_{S_1}^{(m)}(Q_\lambda) < \hat{L}_{S_1}^{(m)}(Q_{\lambda'})$, $\hat{L}_{S_1}^{(m)}(Q_{\lambda'}) + \varepsilon_{mc} < 1$, and $\text{KL}_{ub}(\lambda) \leq \text{KL}_{ub}(\lambda')$, then $B_{\text{cert}}(\lambda) < B_{\text{cert}}(\lambda')$.*

Proof. Let $\lambda, \lambda' \in \Lambda$ share the same ε_{mc}. Suppose $\hat{L}_{S_1}^{(m)}(Q_\lambda) < \hat{L}_{S_1}^{(m)}(Q_{\lambda'})$, $\hat{L}_{S_1}^{(m)}(Q_{\lambda'}) + \varepsilon_{mc} < 1$, and $\text{KL}_{ub}(\lambda) \leq \text{KL}_{ub}(\lambda')$.

Step 1 (First-argument ordering). Since $\hat{L}_{S_1}^{(m)}(Q_{\lambda'}) + \varepsilon_{mc} < 1$ by hypothesis, the $\min\{1, \cdot\}$ clamp in p_{cert} is inactive for both configurations. Hence:

$$p_{\text{cert}}(\lambda) = \hat{L}_{S_1}^{(m)}(Q_\lambda) + \varepsilon_{mc} < \hat{L}_{S_1}^{(m)}(Q_{\lambda'}) + \varepsilon_{mc} = p_{\text{cert}}(\lambda') < 1.$$

Step 2 (Second-argument ordering). Since $\text{KL}_{ub}(\lambda) \leq \text{KL}_{ub}(\lambda')$:

$$\varepsilon_{pb}^{\text{cert}}(Q_\lambda) = \frac{\text{KL}_{ub}(\lambda) + C(n_1, \delta_{pb})}{n_1} \leq \frac{\text{KL}_{ub}(\lambda') + C(n_1, \delta_{pb})}{n_1} = \varepsilon_{pb}^{\text{cert}}(Q_{\lambda'}).$$

Step 3 (Strict monotonicity of kl_+^{-1}). By Theorem 3 (strict monotonicity in p, valid since $\varepsilon_{pb}^{\text{cert}}(Q_\lambda) > 0$):

$$\text{kl}_+^{-1}(p_{\text{cert}}(\lambda),\ \varepsilon_{pb}^{\text{cert}}(Q_\lambda)) < \text{kl}_+^{-1}(p_{\text{cert}}(\lambda'),\ \varepsilon_{pb}^{\text{cert}}(Q_\lambda)).$$

By Theorem 2 (weak monotonicity in ε):

$$\text{kl}_+^{-1}(p_{\text{cert}}(\lambda'),\ \varepsilon_{pb}^{\text{cert}}(Q_\lambda)) \leq \text{kl}_+^{-1}(p_{\text{cert}}(\lambda'),\ \varepsilon_{pb}^{\text{cert}}(Q_{\lambda'})).$$

Combining: $\text{kl}_+^{-1}(p_{\text{cert}}(\lambda), \varepsilon_{pb}^{\text{cert}}(Q_\lambda)) < \text{kl}_+^{-1}(p_{\text{cert}}(\lambda'), \varepsilon_{pb}^{\text{cert}}(Q_{\lambda'}))$. Adding the common term η_{num} to both sides preserves the strict inequality, giving $B_{\text{cert}}(\lambda) < B_{\text{cert}}(\lambda')$. □

10 Master Theorem

Theorem 14 (Certified framework for all three criteria). *Let $\Lambda = \{\lambda_1, \ldots, \lambda_M\}$ be a finite hyperparameter grid with $M \geq 2$, with posterior family $\{Q_\lambda\}_{\lambda \in \Lambda}$ on parameter space $\Theta \subseteq \mathbb{R}^d$, and $n_1 \geq 8$. Let the following assumptions hold:*

(A1) ***Prior independence.*** *The prior P on Θ is $\sigma(S_0)$-measurable and independent of the evaluation sample $S_1 \sim \mathcal{D}^{n_1}$.*

(A2) ***Finite KL.*** *For each $\lambda \in \Lambda$, $\mathrm{KL}(Q_\lambda \| P) \leq \mathrm{KL}_{ub}(\lambda) < \infty$.*

(A3) ***Monte Carlo access.*** *For each λ, m i.i.d. weight samples from Q_λ are available, with Hoeffding tolerance $\varepsilon_{mc} = \sqrt{\ln(2/\delta_{mc})/(2m)}$.*

(A4) ***Distinct risks.*** *The true risks $\{L_{\mathcal{D}}(Q_\lambda)\}_{\lambda \in \Lambda}$ are distinct with minimum gap $\delta_L > 0$.*

(A5) ***KL computability.*** *For each $\lambda \in \Lambda$, $\mathrm{KL}_{ub}(\lambda)$ is computable in time polynomial in $(d, n_1, \log(1/\delta))$.*

(A6) ***Efficient sampling and evaluation.*** *Drawing $\theta \sim Q_\lambda$ and evaluating h_θ on one example are each computable in time polynomial in d.*

Set $\delta = \delta_{pb} + \delta_{mc} + \delta_{num}$. Then with probability $\geq 1 - M\delta$ over S_1 and any auxiliary randomness (conditional on S_0), the certified PAC-Bayes framework provides the following simultaneous guarantees:

(I) Non-vacuity (sufficient condition). *For every $\lambda \in \Lambda$:*

$$L_{\mathcal{D}}(Q_\lambda) \leq B_{\mathrm{cert}}(Q_\lambda) \leq \hat{L}^{(m)}_{S_1}(Q_\lambda) + \varepsilon_{mc} + \sqrt{\frac{\mathrm{KL}_{ub}(\lambda) + C(n_1, \delta_{pb})}{2n_1}} + \eta_{num}.$$

In particular, $B_{\mathrm{cert}}(Q_\lambda) < 1$ whenever the right-hand side is < 1, and $B^{\mathrm{Ma}}_{\mathrm{kl}} \leq B^{\mathrm{Ma}}_{\surd} < B^{\mathrm{Mc}}_{\surd}$ strictly (Theorem 10).

(II) Polynomial-time computation. *The total computation for all M certificates requires*

$$O(M \cdot T_{\mathrm{KL}} + M \cdot m \cdot T_{\mathrm{fwd}} + M \cdot \log_2(1/\eta_{num}))$$

time, which is polynomial in $(n_1, d, M, \log(1/\delta), 1/\varepsilon_{mc})$.

(III) Predictive tightness. *With $\mathrm{gap}_{\max} := \max_\lambda[B_{\mathrm{cert}}(Q_\lambda) - L_{\mathcal{D}}(Q_\lambda)]$, $k_{\max} = \lfloor \mathrm{gap}_{\max}/\delta_L \rfloor$, and $\tilde{k} = \min(k_{\max}, M-1)$:*

$$\tau_a \geq 1 - \frac{2\tilde{k}(2M - \tilde{k} - 1)}{M(M-1)}.$$

Full rank consistency ($\tau_a = 1$) holds when $\mathrm{gap}_{\max} < \delta_L$. Additionally, for any λ, λ' sharing the same ε_{mc}: if $\hat{L}^{(m)}_{S_1}(Q_\lambda) < \hat{L}^{(m)}_{S_1}(Q_{\lambda'})$, $\hat{L}^{(m)}_{S_1}(Q_{\lambda'}) + \varepsilon_{mc} < 1$, and $\mathrm{KL}_{ub}(\lambda) \leq \mathrm{KL}_{ub}(\lambda')$, then $B_{\mathrm{cert}}(\lambda) < B_{\mathrm{cert}}(\lambda')$ (direct monotonicity).

Proof. The proof assembles Theorems 5–13 with no new derivations.

Probability event. Theorem 5 gives $\Pr[L_{\mathcal{D}}(Q_\lambda) \leq B_{\mathrm{cert}}(Q_\lambda) \mid S_0] \geq 1 - \delta$ for each fixed λ. By the union bound over $|\Lambda| = M$ configurations, $\Pr[\mathcal{E} \mid S_0] \geq 1 - M\delta$ where $\mathcal{E} := \{\forall \lambda \in \Lambda : L_{\mathcal{D}}(Q_\lambda) \leq B_{\mathrm{cert}}(Q_\lambda)\}$ is the event from Theorem 12. All statements below condition on $\mathcal{E}$.

Criterion (I). On $\mathcal{E}$, each $B_{\mathrm{cert}}(Q_\lambda)$ is a valid upper bound on $L_{\mathcal{D}}(Q_\lambda)$. The Pinsker relaxation upper bound follows from Theorem 10, and the strict dominance chain $B^{\mathrm{Ma}}_{\mathrm{kl}} \leq B^{\mathrm{Ma}}_{\surd} < B^{\mathrm{Mc}}_{\surd}$ follows from the comparison inequality and Lemma 8.

Criterion (II). Immediate from Theorem 7: KL computation contributes $O(T_{\mathrm{KL}})$ per configuration, Monte Carlo sampling contributes $O(m \cdot T_{\mathrm{fwd}})$ per configuration, and bisection contributes $O(\log_2(1/\eta_{num}))$ per configuration, yielding $O(M \cdot T_{\mathrm{KL}} + M \cdot m \cdot T_{\mathrm{fwd}} + M \cdot \log_2(1/\eta_{num}))$ in total.

Criterion (III). The Kendall τ lower bound is Theorem 12, with the domain guard $\tilde{k} = \min(k_{\max}, M-1)$ ensuring the bound lies in $[-1, 1]$. Direct monotonicity is Theorem 13, providing a complementary pairwise sufficient condition for certificate ordering. □

Remark 15 (Tighter probability allocation)**.** *The probability guarantee in Theorem 14 can be sharpened from $1-M\delta$ to $1-\delta_{\mathrm{global}}$, where $\delta_{\mathrm{global}} = \delta_{pb}+\delta_{mc}+\delta_{num}$, by exploiting the uniformity of the PAC-Bayes-kl event. Since Theorem 4 holds simultaneously for* all *posteriors Q on a single event of probability $\geq 1-\delta_{pb}$, the PAC-Bayes failure budget need only be paid once rather than M times. The union bound is then applied only over the M per-configuration Monte Carlo events (each allocated δ_{mc}/M) and numerical events (each allocated δ_{num}/M), yielding total failure probability $\delta_{pb} + \delta_{mc} + \delta_{num}$. The per-configuration Monte Carlo sample requirement increases from $m \geq \ln(2/\delta_{mc})/(2\varepsilon_{mc}^2)$ to $m \geq \ln(2M/\delta_{mc})/(2\varepsilon_{mc}^2)$ — a logarithmic cost that preserves polynomial-time computability. This observation is already used in Theorem 7 (which notes "the PAC-Bayes event E_{pb} is shared across all λ"); the simpler $1 - M\delta$ form in Theorems 12 and 14 is a valid conservative bound chosen for presentational clarity.*

11 Empirical Verification (Summary)

All three criteria are confirmed empirically on real neural networks at two scales. In experiments, ε_{mc} and η_{num} are set to 0, treating Monte Carlo and numerical error as negligible; the reported bounds are therefore plug-in approximations to the formal certificates of Theorem 5. True risk is estimated from a held-out test set.

MNIST (784-600-600-10, d = 837,610 parameters): 10/10 non-vacuous kl-inverse certificates. Best certificate $B = 0.323$ (true risk $L_{\mathcal{D}} = 0.066$). Within-group Kendall $\tau = +1.0$ at $\sigma = 0.05$.

ImageNet (4 architectures: ResNet-18, ResNet-50, MobileNetV2, EfficientNet-B0): 32/32 non-vacuous certificates across all configurations. Key results:

Architecture	d	$B_{\mathrm{kl}}^{\mathrm{Ma}}$	Gap
EfficientNet-B0	5.3M	**0.239**	1.6%
MobileNetV2	3.5M	**0.287**	1.8%
ResNet-18	11.7M	**0.304**	1.8%
ResNet-50	25.6M	0.999	(degenerate)

The tightest certificates occur at KL = 0 (no fine-tuning), where the posterior equals the prior and is independent of S_1; in this regime, a direct Hoeffding bound gives comparable results. The PAC-Bayes framework's value lies in providing a unified treatment across all KL values, including the fine-tuned (KL > 0) regime where direct concentration bounds are unavailable.

Overall Kendall $\tau = 0.692$ ($p = 2.65 \times 10^{-8}$), Spearman $\rho = 0.834$ ($p = 3.01 \times 10^{-9}$). No-fine-tune subset: $\tau = 0.983$, $\rho = 0.997$. Excluding degenerate ResNet-50: $\tau = 1.000$.

Combined: 42/42 non-vacuous certificates (10 MNIST + 32 ImageNet). All certificates computed in polynomial time (< 217s each). Certificate computation < 30s per configuration on MNIST, < 217s on ImageNet (NVIDIA RTX 3090).

Criterion	Metric	MNIST	ImageNet
1. Non-vacuity	$B_{\mathrm{kl}}^{\mathrm{Ma}} < 1$	10/10	32/32
2. Poly-time	Certificate time	< 30s each	< 217s each
3. Pred. tightness	Kendall τ	+1.0 (within-group)	+0.692 (overall)

Full experimental details, figures, and pairwise concordance analysis are in the companion paper [9].

Remark 16 (Formal non-vacuity witness with tracked error tolerances)**.** *The plug-in convention $\varepsilon_{mc} = 0$, $\eta_{num} = 0$ follows standard practice in the PAC-Bayes empirical literature [3, 4]. To confirm that non-vacuity survives formal error tracking, we exhibit one explicit witness. For EfficientNet-B0 at $\sigma = 0.001$ (no fine-tuning), with $m = 10$, $\delta_{pb} = 0.025$, $\delta_{mc} = 0.025$, $\eta_{num} = 10^{-10}$: Hoeffding gives $\varepsilon_{mc} = \sqrt{\ln(2/\delta_{mc})/(2m)} \approx 0.468$, yielding $p_{\mathrm{cert}} = \hat{L}_{S_1}^{(m)}(Q) + \varepsilon_{mc} \approx 0.693$. With $\mathrm{KL} = 0$ and $C(n_1, \delta_{pb}) \approx 9.19$, the PAC-Bayes slack is $\varepsilon_{pb}^{\mathrm{cert}} \approx 6.13 \times 10^{-4}$, giving $B_{\mathrm{cert}} = \mathrm{kl}_+^{-1}(p_{\mathrm{cert}}, \varepsilon_{pb}^{\mathrm{cert}}) + \eta_{num} \approx 0.709 < 1$. The formal certificate is looser than the plug-in value (0.709 vs 0.239) because ε_{mc} dominates with only $m = 10$ samples, but it remains strictly non-vacuous. Increasing m to 100 yields $\varepsilon_{mc} \approx 0.148$ and $B_{\mathrm{cert}} \approx 0.390$; to 1,000, $\varepsilon_{mc} \approx 0.047$ and $B_{\mathrm{cert}} \approx 0.287$, converging toward the plug-in value of 0.239.*

12 Conclusion

We have resolved the open problem: there exists a single PAC-Bayes framework — the PAC-Bayes-kl inequality with Maurer's constant, kl-inverse certificate computation via certified bisection, and universality over all admissible prior-posterior pairs — that simultaneously achieves non-vacuous bounds at practical scale, polynomial-time computability, and predictive tightness across hyperparameters. The affirmative answer is established through the theorems above, with the Master Theorem (Theorem 14) assembling all components. Empirical verification at two scales confirms all three criteria: 42/42 non-vacuous certificates, polynomial-time computation, and strong rank correlation ($\tau = 0.692$ at ImageNet scale, $\tau = 1.000$ for well-separated configurations), with the tightest certificate $B = 0.239$ lying within 1.6 percentage points of the true risk.

References

[1] A. Maurer. A note on the PAC-Bayesian theorem. *arXiv:0411099*, 2004.

[2] D. McAllester. Some PAC-Bayesian theorems. *Machine Learning*, 37(3):355–363, 1999.

[3] G. K. Dziugaite and D. M. Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. In *Proc. UAI*, 2017.

[4] W. Zhou, V. Veitch, M. Austern, R. P. Adams, and P. Orbanz. Non-vacuous generalization bounds at the ImageNet scale: a PAC-Bayesian compression approach. In *Proc. ICLR*, 2019.

[5] M. S. Pinsker. *Information and Information Stability of Random Variables and Processes.* Holden-Day, 1964.

[6] O. Catoni. *PAC-Bayesian Supervised Classification: The Thermodynamics of Statistical Learning.* IMS Lecture Notes, 2007.

[7] B. Guedj. A primer on PAC-Bayesian learning. *arXiv:1901.05353*, 2019.

[8] S. Arora, R. Ge, B. Neyshabur, and Y. Zhang. Stronger generalization bounds for deep nets via a compression approach. In *Proc. ICML*, 2018.

[9] C. Kang. Companion repository: Tight PAC-Bayes bounds for deep neural networks. `https://github.com/ckchmod/solveall-tightPAC`, 2026.