#### Beta diffusion trees

# Creighton Heaukulani

University of Cambridge

Joint work with

David A. Knowles (Stanford)
Zoubin Ghahramani (Cambridge)

ICML 2014, Beijing June 2014

We are interested in random tree structures on overlapping clusters (aka feature allocations).

1. Briefly review feature allocations.

- 1. Briefly review feature allocations.
- 2. Describe a stochastic process (the beta diffusion tree) that leads to trees over feature allocations.

- 1. Briefly review feature allocations.
- 2. Describe a stochastic process (the beta diffusion tree) that leads to trees over feature allocations.
- 3. Describe a hierarchically-clustered factor analysis model.

- 1. Briefly review feature allocations.
- 2. Describe a stochastic process (the beta diffusion tree) that leads to trees over feature allocations.
- 3. Describe a hierarchically-clustered factor analysis model.
- 4. Present some inference techniques over the tree structure and some experimental results.

 $Feature\ allocations = Overlapping\ subsets = Overlapping\ clusters$ 

 $Feature\ allocations = Overlapping\ subsets = Overlapping\ clusters$ 


Consider an image labelling application:



Feature allocations = Overlapping subsets = Overlapping clusters

Consider an image labelling application:






Feature allocations = Overlapping subsets = Overlapping clusters

Consider an image labelling application:






Feature allocations = Overlapping subsets = Overlapping clusters

Consider an image labelling application:





Feature allocations = Overlapping subsets = Overlapping clusters

Consider an image labelling application:








"image is in a kitchen" {1}

 $\label{eq:continuous} Feature \ allocations = Overlapping \ subsets = Overlapping \ clusters$ 

Consider an image labelling application:

Feature allocation:




 $Feature \ allocations = Overlapping \ subsets = Overlapping \ clusters$ 

Consider an image labelling application:

Feature allocation:

The clusters/subsets are called features






#### Hierarchies on features

Hierarchies of features are a natural assumption:

"image is in a kitchen" "image contains a chair" "image contains a table"

#### Hierarchies on features


Hierarchies of features are a natural assumption:



Consider a sequence of particles, labeled with one of N objects, diffusing in a continuous space  $\mathcal{X}$  over the unit time interval [0,1].

Consider a sequence of particles, labeled with one of N objects, diffusing in a continuous space  $\mathcal{X}$  over the unit time interval [0,1].

#### For example:



- $\mathcal{X} = \mathbb{R}$ .
- ▶ Diffusion is Brownian motion.


Consider a sequence of particles, labeled with one of N objects, diffusing in a continuous space  $\mathcal{X}$  over the unit time interval [0,1].

Consider a sequence of particles, labeled with one of N objects, diffusing in a continuous space  $\mathcal{X}$  over the unit time interval [0,1].


At random times:

- ▶ Replicate: Copies of particles may be created.
- ▶ Stop: Particles are destroyed.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.




Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



Consider the first object.


Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



- 1. The first particle starts at the origin.
- 2. May replicate or stop after  $\exp(\lambda_r)$  or  $\exp(\lambda_s)$  time.

Consider the first object.


Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



- 1. The first particle starts at the origin.
- 2. May replicate or stop after  $\exp(\lambda_r)$  or  $\exp(\lambda_s)$  time.
- 3. Diffuses as Brownian motion until t = 1.

Consider the first object.


Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



1. Follow previous particles.

Consider the second object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.




- 1. Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$
 amount of time.

Consider the second object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.




- 1. Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$
 amount of time.

Consider the second object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.




- 1. Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$
 amount of time.

Consider the second object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.




- 1. Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$
 amount of time.

Consider the second object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.




- 1. Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$
 amount of time.

Consider the third object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



Consider the third object.

- 1. Follow previous particles.
- 2. May replicate or stop after


$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$

amount of time.

3. Replicate and stop w.p.

$$rac{n_r}{ heta_r+m}$$
 and  $rac{n_s}{ heta_s+m}$ 

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



Consider the third object.

- 1. Follow previous particles.
- 2. May replicate or stop after


$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$

amount of time.

3. Replicate and stop w.p.

$$rac{n_r}{ heta_r+m}$$
 and  $rac{n_s}{ heta_s+m}$ 

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



- 1. Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$


amount of time.

3. Replicate and stop w.p.

and 
$$\frac{n_r}{\theta_r+m}$$
  $\frac{n_s}{\theta_r+m}$ 

Consider the third object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



- 1. Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$


amount of time.

3. Replicate and stop w.p.

$$rac{n_r}{ heta_r+m}$$
 and  $rac{n_s}{ heta_s+m}$ 

Consider the third object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



- Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$


amount of time.

Replicate and stop w.p.

and 
$$\frac{n_r}{\theta_r+m}$$
  $\frac{n_s}{\theta_r+m}$ 

Consider the third object.

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



Consider the third object.

- 1. Follow previous particles.
- 2. May replicate or stop after


$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$

amount of time.

Replicate and stop w.p.

$$rac{n_r}{ heta_r+m}$$
 and  $rac{n_s}{ heta_s+m}$ 

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



Consider the third object.

- 1. Follow previous particles.
- 2. May replicate or stop after


$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$

amount of time.

Replicate and stop w.p.

$$rac{n_r}{ heta_r+m}$$
 and  $rac{n_s}{ heta_s+m}$ 

Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



Consider the third object.

- 1. Follow previous particles.
- 2. May replicate or stop after

$$\exp\Bigl(\frac{\theta_r}{\theta_r+m}\lambda_r\Bigr)$$
 or 
$$\exp\Bigl(\frac{\theta_s}{\theta_s+m}\lambda_s\Bigr)$$


amount of time.

Replicate and stop w.p.


$$rac{n_r}{ heta_r+m}$$
 and  $rac{n_s}{ heta_s+m}$ 

8


Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.




The corresponding tree structure:



Let  $\lambda_s, \lambda_r, \theta_s, \theta_r > 0$  and  $\mathcal{X} = \mathbb{R}$ . Diffusion is Brownian motion.



The corresponding tree structure:



The leaves form a feature allocation of [3]:

$$\{\{1,3\},\{2,3\}\}$$


#### Related work

#### As opposed to...

Random tree structures over non-overlapping clusters (aka partitions):


- ▶ Dirichlet diffusion tree (Neal, 2003 [6])
- ▶ Pitman–Yor diffusion tree (Knowles & Ghahramani, 2011 [5])
- ► Kingman's coalescent (Kingman, 1982 [4])


Consider the following larger example:




11

Consider the following larger example:






Consider the following larger example:




A tree structure over the columns of a binary matrix.

Connections to the <u>Indian buffet process</u> (Ghahramani, Griffiths, & Sollich, 2007 [3]).

1



We defined the tree in terms of the ordered sequence of objects, however:



We defined the tree in terms of the ordered sequence of objects, however:

### Theorem (exchangeability)


The beta diffusion tree is <u>exchangeable</u> with respect to the ordering of the objects.

# Other properties


- ► The beta diffusion tree is the continuum limit of a <u>nested</u> feature allocation scheme
- ▶ The beta diffusion tree is a continuous-time branching process

Example: A hierarchical factor analysis model

# Example: A hierarchical factor analysis model



# Example: A hierarchical factor analysis model



Both the factors and factor loadings are hierarchically structured

We perform a series of Metropolis–Hastings proposals to integrate over the random tree structures.

1. Resample the paths of particles down the tree.

- 1. Resample the paths of particles down the tree.
- 2. Propose removing an internal (either replicate or stop) node. Conversely, propose a node to a branch.


- 1. Resample the paths of particles down the tree.
- 2. Propose removing an internal (either replicate or stop) node. Conversely, propose a node to a branch.
- 3. Propose changing the decisions that particles take at internal nodes (i.e., the decisions to either replicate or stop).


- 1. Resample the paths of particles down the tree.
- 2. Propose removing an internal (either replicate or stop) node. Conversely, propose a node to a branch.
- 3. Propose changing the decisions that particles take at internal nodes (i.e., the decisions to either replicate or stop).


Each is accepted or rejected with a Metropolis–Hastings step.

## Some numerical results

Test log-likelihoods:








- (a) A gene expression dataset
- (b) International development statistics
- (c) Household socioeconomic indicators

# An analysis of UN human development statistics

We can extend the analysis by Doshi–Velez & Ghahramani (2009) [1]:



- F. Doshi-Velez and Z. Ghahramani. Correlated non-parametric latent feature models. In <u>Proc. UAI</u>, 2009.
   J. Geweke. Getting it right: Joint distribution tests of posterior simulators. Journal of the American
- Statistical Association, 99(467):799-804, 2004.

  Z. Ghahramani, T. L. Griffiths, and P. Sollich, Bayesian nonparametric latent feature models.
- Bayesian Statistics, 8:201–226, 2007. See also the discussion and rejoinder.
- [4] J. F. C. Kingman. The coalescent. Stochastic Processes and Their Applications, 13(3):235–248, 1982.
- [5] D. Knowles and Z. Ghahramani. Pitman-Yor diffusion trees. In Proc. UAI, 2011.
- [6] R. M. Neal. Density modeling and clustering using Dirichlet diffusion trees. <u>Bayesian Statistics</u>, 7, 2003.