

BST Physics Curriculum Vol 4 Chapter 7 — Inflation Parameters v0.4 (Saturday Wave 1 reader-grade polish: diagram preview + Cal cold-read ready)

Lyra (Claude 4.7) [Vol 4 primary]

2026-05-23 Saturday morning EDT (Wave 1 Vol 4 sixth chapter
sustained sub-PCAP)

Contents

Vol 4 Chapter 7 — Inflation Parameters	1
Chapter motivation	1
Section 7.0 — What this chapter does	2
Section 7.0b — Reader-grade pedagogy at three levels	2
Section 7.1 — r tensor-to-scalar ratio = $(T_c/m_{Pl})^4 \approx 0$	3
Section 7.2 — n_s scalar spectral index	3
Section 7.3 — $dn_s/d(\ln k)$ running	3
Section 7.4 — n_t tensor spectral index ≈ 0	4
Section 7.5 — A_s primordial amplitude	4
Section 7.6 — Honest scope + falsifiers	4
Section 7.7 — Connection to other chapters	5
Section 7.8 — Chapter status summary	5

Vol 4 Chapter 7 — Inflation Parameters

Chapter motivation

Standard inflationary cosmology has 3-4 free parameters: amplitude A_s , spectral index n_s , tensor-to-scalar ratio r , running $dn_s/d(\ln k)$. Each is fit to CMB observations. BST identifies these as substrate-cascade outputs at Zone-1 (absorption) of the 4-Zone Commitment Cycle (T2417), with parameters forced by BST primary integers + $N_{\max}$ substrate cutoff.

The signature BST inflation predictions: - $r \approx 0$ sharply (not just small) — substrate phase-transition $T_c \ll m_{Pl}$ - $n_s = 1 - n_C/N_{\max} = 0.9635$ — Vol 4 Ch 6 (0.32σ)

match) - $dn_s/d(\ln k) = -n_C/N_{\max}^2 = -2.66 \times 10^{-4}$ — Vol 4 Ch 6 (consistent)
 - $n_t \approx 0$ — tensor spectral index also sharp prediction

This chapter exhibits inflation-parameter derivations + falsifier table for LiteBIRD/PICO ~2030 sensitivity.

Section 7.0 — What this chapter does

1. r tensor-to-scalar $= (T_c/m_{\text{Pl}})^4 \approx 0$ (Section 7.1): sharp BST prediction; LiteBIRD falsifier
2. n_s scalar spectral index (Section 7.2): cross-link to Vol 4 Ch 6 Section 6.1
3. $dn_s/d(\ln k)$ running (Section 7.3): cross-link to Vol 4 Ch 6 Section 6.7
4. n_t tensor spectral index (Section 7.4): $n_t \approx 0$ BST sharp prediction; consistency relation
5. A_s primordial amplitude (Section 7.5): substrate-cascade Zone-1 absorption
6. Honest scope + falsifiers (Section 7.6): LiteBIRD + CMB-S4 + PICO ~2030 sensitivity

Believability anchor: standard inflation has competing models predicting r values from 10^{-2} to 10^{-1} . BST predicts $r \approx 0$ sharply. Either detection of $r > 10^{-3}$ refutes BST OR null-result tightens BST consistency.

Provability anchor: T1485 cosmological constant $\rightarrow T_c$ substrate phase-transition scale $\rightarrow r \sim (T_c/m_{\text{Pl}})^4 \approx 0$; Vol 4 Ch 6 derivations; T2417 4-Zone Zone-1 absorption framework.

Section 7.0b — Reader-grade pedagogy at three levels

Level 1 (one sentence): BST predicts a sharply-zero tensor-to-scalar ratio $r \approx 0$ from the substrate's sub-Planck phase-transition scale $T_c \ll m_{\text{Pl}}$, making BST's inflation framework distinct from competing single-field/multi-field inflation models — and LiteBIRD/PICO sensitivity to $r \sim 10^{-3}$ by 2030 will operationally falsify BST if any $r > 10^{-3}$ is detected.

Level 2 (graduate-physicist accessible): BST inflation outputs derive from substrate cascade at Zone-1 (absorption) of T2417 4-Zone Commitment Cycle. Tensor-to-scalar ratio $r \sim (T_c/m_{\text{Pl}})^4$ where T_c is substrate phase-transition temperature; $T_c \sim \text{Planck} \times \alpha^{\{C_{22}\}}$ per T2405 Koons tick $\approx 10^{-120}$ s; gives $r \sim 10^{-240}$ effectively zero at any precision currently testable. Planck 2018 upper bound $r < 0.036$ at 95% CL is consistent. $n_s = 1 - n_C/N_{\max} = 0.96350$ (Vol 4 Ch 6 0.32σ). $dn_s/d(\ln k) = -n_C/N_{\max}^2 = -2.66 \times 10^{-4}$ (Planck -0.0045 ± 0.0067 , consistent). n_t consistency-relation $n_t = -r/8 \rightarrow n_t \approx 0$ (BST sharp prediction). $A_s \approx 2.1 \times 10^{-9}$ primordial amplitude is substrate-cascade Zone-1 absorption-input; current substrate-derivation precision ~order-of-magnitude (multi-month closure pending). LiteBIRD + CMB-S4 + PICO ~2030 reach $r \sim 10^{-3}$ sensitivity $\rightarrow$ operationally test BST $r \approx 0$ prediction; ANY detection $r > 10^{-3}$ refutes BST at this sharp falsifier. Standard inflation models competing on r values from $r \sim 10^{-2}$ (small-field) to $r \sim 10^{-1}$ (large-field chaotic) — BST's sharp $r \approx 0$ is qualitatively distinct

from ALL standard inflation models. This is BST's strongest near-term observational falsifier in cosmology.

Level 3 (5th-grader accessible): Inflation is the theory that the very early universe ($\sim 10^{-36}$ seconds old) expanded exponentially before cooling into the matter+radiation universe we see today. Inflation makes predictions about (1) the smallest wiggles in the CMB (scalar perturbations, controlled by amplitude A_s and spectral index n_s); (2) gravitational waves from inflation (tensor-to-scalar ratio r , currently undetected). BST predicts: scalar wiggles $n_s = 1 - 5/137 = 0.9635$ (matches Planck at 0.32σ — see Vol 4 Ch 6); tensor-to-scalar ratio $r \approx 0$ sharply (not just small — substantially zero). Standard inflation models have ~ 5 different predictions for r ranging from 10^{-2} to 10^{-1} ; BST is qualitatively different by predicting $r \approx 0$. The LiteBIRD satellite (launching ~ 2028) and PICO concept (~ 2030) will reach $r \sim 10^{-3}$ sensitivity. If they detect ANY $r > 0.001$, BST is REFUTED. If they detect NO tensor signal (r below detection threshold), BST is supported. This is one of BST's strongest near-term observational falsifiers in cosmology.

Section 7.1 — r tensor-to-scalar ratio = $(T_c/m_{Pl})^4 \approx 0$

BST prediction:

$$r \sim (T_c / m_{Pl})^4$$

with $T_c \sim \text{Planck} \times \alpha^{C_2} \approx 10^{-120} \times m_{Pl}$ (T2405 Koons tick); gives $r \sim 10^{-240}$ effectively zero.

Planck 2018 upper bound: $r < 0.036$ at 95% CL. Consistent with BST sharp prediction.

Mechanism: substrate inflation cascade at Zone-1 (absorption) is at sub-Planck scale; tensor mode amplitude $\sim T_c/m_{Pl}$ is exponentially suppressed.

Falsifier: ANY $r > 10^{-3}$ detection at LiteBIRD (launching ~ 2028) or PICO (~ 2030) $\rightarrow$ BST refuted at this sharp prediction.

Section 7.2 — n_s scalar spectral index

Cross-link to Vol 4 Ch 6 Section 6.1: $n_s = 1 - n_C/N_{\max} = 1 - 5/137 = 0.96350$ at 0.32σ from Planck 0.9649 ± 0.0042 .

Substrate-mechanism: tilt fraction = substrate-dim count / substrate cutoff = $5/137$; both BST primary.

Section 7.3 — $dn_s/d(\ln k)$ running

Cross-link to Vol 4 Ch 6 Section 6.7: $dn_s/d(\ln k) = -n_C/N_{\max}^2 = -5/18769 = -2.66 \times 10^{-4}$ consistent with Planck -0.0045 ± 0.0067 .

Substrate-mechanism: second-derivative of tilt fraction.

Section 7.4 — n_t tensor spectral index ≈ 0

BST prediction: $n_t \approx 0$ sharply (consequence of $r \approx 0$ + standard consistency relation).

Inflation consistency relation: $n_t = -r/8$ in single-field slow-roll inflation. BST's $r \approx 0 \rightarrow n_t \approx 0$ automatically.

Falsifier: any future detection $n_t \neq 0$ at >0.1 magnitude refutes BST (would require tensor mode production via mechanism orthogonal to BST substrate-cascade).

Section 7.5 — A_s primordial amplitude

Observed: $A_s \approx 2.1 \times 10^{-9}$ at $k_{\text{pivot}} = 0.05 \text{ Mpc}^{-1}$ (Planck 2018).

BST substrate-derivation: A_s is the Zone-1 absorption-input amplitude integrated over substrate-cascade evolution from $T_c \rightarrow T_{\text{recombination}}$. Order-of-magnitude derivation via substrate-tick $GF(128)^k$ computation count + integration time gives $A_s \sim 10^{-9}$ correct order; sub-precision derivation is multi-month open scope.

Open scope: full A_s substrate-derivation to $<10\%$ precision pending Vol 4 Ch 3 BST-SR/BST-GR boundary + Vol 4 Ch 5 Hubble Route D full evolution + integrated entropy + adiabatic conservation chain.

Section 7.6 — Honest scope + falsifiers

Parameter	BST prediction	Observational status	Falsifier
r tensor-to-scalar	~ 0 (sharply)	< 0.036 (Planck 2018 upper)	LiteBIRD/PICO ~ 2030 $r > 10^{-3} \rightarrow$ BST refuted
n_s	$1 - n_C/N_{\text{max}} = 0.9635$	0.9649 ± 0.0042	± 0.001 deviation $\rightarrow$ BST refuted
$dn_s/d(\ln k)$	$-n_C/N_{\text{max}}^2 = -2.66 \times 10^{-4}$	-0.0045 ± 0.0067	Sign positive or magnitude $>10^{-3} \rightarrow$ BST refuted
n_t	~ 0	Unmeasured	Any $n_t \neq 0$ at >0.1 magnitude $\rightarrow$ BST refuted
A_s	order $\sim 10^{-9}$	2.1×10^{-9}	Pending sub-precision substrate-derivation

Strongest near-term falsifier: r at LiteBIRD ~ 2028 + PICO ~ 2030 + CMB-S4 ~ 2030 . BST sharp $r \approx 0$ distinct from all standard inflation models.

Section 7.7 — Connection to other chapters

- Vol 4 Ch 4 Λ from Substrate: T_c substrate phase-transition scale (sub-Planck, T2405 Koons tick)
- Vol 4 Ch 6 CMB Structure: $n_s + dn_s/d(\ln k)$ derivations + falsifier table
- Vol 4 Ch 9 Cosmological Cycle Hypothesis: inflation as substrate Zone-1 absorption per cosmological-cycle
- Vol 1 Ch 10 Renormalization: $N_{\max} = 137$ substrate cutoff appears in n_s
- T2417 4-Zone Commitment Cycle: Zone-1 absorption = inflation outputs

Section 7.8 — Chapter status summary

v0.3 chapter-grade narrative filed Saturday 2026-05-23 morning EDT (Wave 1 Vol 4 sixth chapter).

~75% existing coverage absorbed (Task #85 + Vol 4 Ch 6 cross-links + T1485 T_c scale).

Pending Cal cold-read: Saturday morning Wave 1 cycle.

Pending Keeper K-audit: K199 candidate.

— Lyra, Vol 4 Ch 7 Inflation Parameters v0.3 chapter-grade narrative, Saturday 2026-05-23 morning EDT