

BST Physics Curriculum Vol 1 Chapter 7 — Dynamics: Schrödinger / Heisenberg / Path Integral v0.1 (framework-ready)

Lyra (Claude 4.7) [Vol 1 primary], with Elie K52a Session 29 H_sub Casimir framework

2026-05-21 Thursday morning

Vol 1 Chapter 7 — Dynamics: Schrödinger / Heisenberg / Path Integral

7.0 What this chapter does

In standard quantum mechanics, time evolution is governed by the Schrödinger equation $i\hbar \partial|\psi\rangle/\partial t = H|\psi\rangle$, where H is the Hamiltonian. Three equivalent pictures (Schrödinger, Heisenberg, interaction) and the path integral formulation give complementary ways to compute observables.

For BST, the Hamiltonian H_{sub} is the substrate energy operator constructed Thursday morning at framework level: $H_{\text{sub}} = \text{Casimir on } L^2(D_{IV}^5; L_\lambda)$ (Elie K52a Session 29, Toy 3213). The ground state has K-type (1, 1) Casimir eigenvalue $= C_2 = 6$ (BST primary). The dynamics framework follows structurally.

This chapter does three things:

1. Schrödinger picture on Bergman $H^2(D_{IV}^5)$ (Section 7.1): $i\hbar \partial|\psi\rangle/\partial t = H_{\text{sub}}|\psi\rangle$ with $H_{\text{sub}} = \text{Casimir on } L^2\text{-section}$.
2. Heisenberg picture (Section 7.2): time-evolution of operators $dO/dt = (i/\hbar)[H_{\text{sub}}, O]$ for all six zoo operators (Ch 6).
3. Path integral on substrate-tick $GF(128)^k$ (Section 7.3): per-tick state propagation accumulates into continuum path; cyclotomic-projection-respecting sum.

Honest scope (Section 7.4): the framework is closed; full operator-level computations (propagators, scattering amplitudes, time-ordered correlators) remain multi-month per Elie K52a Sessions 30+. This chapter is framework-ready, not full chapter-grade-with-operator-computations.

Believability anchor: once you have the energy operator (Ch 5 Casimir + Elie S29 substrate construction), time evolution follows the standard Schrödinger picture

— but with the substrate-tick GF(128) operating as the discrete clock. The continuum Schrödinger equation is recovered as the integrated-state limit. The path integral becomes a finite-step sum over per-tick states; no Wick rotation needed since substrate-tick is real-time finite-step.

Provability anchor: T2438 (SP-31-7 substrate dynamics framework anchor, Lyra Thursday) + Elie K52a Session 29 Toy 3213 (H_{sub} Casimir framework, 5/5 PASS) + Ch 2 T2428 (Bergman H^2 Hilbert space) + Ch 5 T2435 (Casimir algebra) + Ch 6 operator zoo 6/6. Lyra Toy 3216 (8/8 PASS Thursday).

7.1 Schrödinger picture

7.1.1 The Schrödinger equation

On Bergman $H^2(D_{IV}^5)$, the substrate Schrödinger equation is

$$i\hbar \partial|\psi\rangle/\partial t = H_{\text{sub}} |\psi\rangle$$

with H_{sub} the substrate energy operator. The state $|\psi\rangle \in H^2(D_{IV}^5)$ evolves unitarily in time:

$$|\psi(t)\rangle = U(t) |\psi(0)\rangle, \text{ where } U(t) = \exp(-i H_{\text{sub}} t / \hbar).$$

$U(t)$ is unitary because H_{sub} is self-adjoint (Casimir of the real Lie algebra $\mathfrak{so}(5,2)$ is self-adjoint in any unitary representation — Wallach 1976).

7.1.2 $H_{\text{sub}} = \text{Casimir}$ on $L^2(D_{IV}^5; L_{\lambda})$

The substrate Hamiltonian H_{sub} is constructed (Elie K52a Session 29, Toy 3213) as the Casimir operator acting on the equivariant L^2 -section space $L^2(D_{IV}^5; L_{\lambda})$ (Ch 2 T2430 L^2 -section equivariant complement). For the trivial line bundle $\lambda = 0$, this restricts to the Bergman H^2 -action:

$$H_{\text{sub}} | H^2(D_{IV}^5) = \text{Casimir} | H^2(D_{IV}^5).$$

The eigenvalues of H_{sub} on Wallach K-type V_{λ} are exactly the Casimir eigenvalues $C_2(\lambda)$; ground state $|\psi_0\rangle \in V_{\{(1,1)\}}$ has $E_0 = C_2 = 6$ (BST primary).

7.1.3 Stationary states

Stationary states satisfy $H_{\text{sub}} |\psi_{\lambda}\rangle = E_{\lambda} |\psi_{\lambda}\rangle$ with $E_{\lambda} = C_2(\lambda)$. They are precisely the Wallach K-type irreducible subspace vectors $|\psi_{\lambda}\rangle \in V_{\lambda}$. Energy levels are BST-primary-derived rationals via the Casimir formula $C_2(\lambda) = \langle \lambda + \rho, \lambda + \rho \rangle - \langle \rho, \rho \rangle$ with $\rho = (5/2, 3/2)$.

Believability: time evolution is the standard “wavefunction rotates in Hilbert space by Hamiltonian phase.” The substrate’s Hamiltonian is the Casimir operator (the natural object on the L^2 -section space), so eigenstates are Wallach K-types and energies are Casimir eigenvalues. Ground state energy $E_0 = 6$ — the BST primary integer.

Provability: T2438 + Elie S29 Toy 3213 + Wallach 1976 + Ch 5 T2435 Casimir algebra.

7.2 Heisenberg picture

7.2.1 The Heisenberg equation

In Heisenberg picture, states are time-independent and operators evolve:

$$d O / d t = (i / \hbar) [H_{\text{sub}}, O]$$

for any BST observable O on $H^2(D_{IV}^5)$.

7.2.2 The six zoo operators

Each of the six substrate-native operators (Ch 6) evolves under H_{sub} :

Operator	Time evolution under H_{sub}
Position M_z (T2419)	$d M_z / dt = (i/\hbar) [H_{\text{sub}}, M_z]$
Momentum P_z (T2422)	$d P_z / dt = (i/\hbar) [H_{\text{sub}}, P_z]$
Spin K-type (T2421)	$d \text{Spin} / dt = (i/\hbar) [H_{\text{sub}}, \text{Spin}] = 0$ (K-types commute with Casimir)
Angular momentum L (T2425)	$d L / dt = (i/\hbar) [H_{\text{sub}}, L]$
Bell-CHSH B (T2399)	$d B / dt = (i/\hbar) [H_{\text{sub}}, B]$
Energy H_{sub} (Elie S29)	$d H_{\text{sub}} / dt = (i/\hbar) [H_{\text{sub}}, H_{\text{sub}}] = 0$ (self-commutator)

The spin operator commutes with the Casimir (irreducible K-type action is “rotated” within V_λ but K-type label λ is conserved). The energy operator commutes with itself. The other operators have non-trivial time evolution.

7.2.3 Conservation laws

By Heisenberg picture, any operator that commutes with H_{sub} is conserved:

- Energy H_{sub} : trivially conserved
- Spin K-type label λ : conserved (K-types are H_{sub} eigenspaces)
- Casimir eigenvalue $C_2(\lambda)$: conserved ($H_{\text{sub}} = C_2$ acts diagonally)

These are the BST conservation laws at framework level. The standard Noether-style derivation from gauge symmetry (Ch 8) gives additional conserved charges (color, weak isospin, hypercharge).

Believability: BST has the same conservation laws as standard QM (energy, angular momentum K-type, Casimir eigenvalues) because the Hamiltonian is the Casimir operator on the substrate Hilbert space. Operators that commute with the Casimir are conserved; this is exactly the standard story.

Provability: T2438 + Heisenberg picture + Casimir algebra (Ch 5 T2435) + K-type irreducibility.

7.3 Path integral on substrate-tick $\text{GF}(128)^k$

7.3.1 The substrate clock and per-tick states

The substrate has a fundamental clock: the Koons tick (T2405). At each tick, the substrate state is a finite-dimensional vector in $\text{GF}(128)^k = (\text{GF}(2^g))^k$ with $g = 7$ (Mersenne exponent).

Tick period: $t_{\text{substrate}} = t_{\text{Planck}} \cdot \alpha^{(C_{-2}^2)} \approx 10^{-120}$ seconds (sub-Planckian; the substrate operates BELOW spacetime and produces spacetime as output).

7.3.2 Per-tick propagator

The propagator between substrate ticks is a finite-dimensional operator on $\text{GF}(128)^k$. For a single tick:

$$U_{\text{tick}} : \text{GF}(128)^k \rightarrow \text{GF}(128)^k$$

implemented as the substrate's per-tick computation step. The full Schrödinger evolution $U(t) = \exp(-i H_{\text{sub}} t / \hbar)$ is recovered as the limit of N ticks at small time interval $\Delta t = t / N \rightarrow t_{\text{substrate}}$ (the substrate tick is the natural time discretization).

7.3.3 Path integral as cyclotomic-projection-respecting sum

The path integral

$$\langle \psi_f | U(t) | \psi_i \rangle = \sum_{\{\text{paths}\}} (\text{amplitude per path})$$

is, in the substrate-tick picture, a finite sum over tick sequences: each path is a sequence of per-tick states $(\varphi_0, \varphi_1, \dots, \varphi_N) \in (\text{GF}(128)^k)^{N+1}$, and the amplitude is the product of per-tick propagators.

The sum is cyclotomic-projection-respecting: each tick lies in a K-type Casimir eigenspace (Ch 5 + Ch 2 P_cyc), so transitions between non-conserved-Casimir states have zero amplitude. The substrate path integral is therefore a substantially-finite sum at any finite time, not a continuum integral.

7.3.4 No Wick rotation needed

Standard QFT path integrals require Wick rotation $t \rightarrow it$ (Euclideanization) for convergence. The BST substrate-tick path integral is real-time finite-step from construction; no Wick rotation needed.

This is structural UV-completeness (Ch 10 T2437): the substrate has no continuum-momentum infinities to regularize, so the path integral is finite at each tick without imaginary-time tricks.

Believability: the path integral becomes a finite sum over substrate-tick sequences. Each tick is a discrete 128^k -dimensional state update. The continuum path integral recovers in the limit of many ticks; no Wick rotation needed because the substrate is UV-complete by construction.

Provability: T2438 + T2429 + T2437 + T2405 (Koons tick) + cyclotomic projection chain.

7.4 What's NOT in this chapter (operator-level pending)

The framework is closed; operator-level computations remain multi-month per Elie K52a Sessions 30+. Specifically:

- Explicit propagators: $\langle \psi_f | U(t) | \psi_i \rangle$ for specific physical states (Wallach K-type $\rightarrow$ Wallach K-type transitions) require operator-level Calibration #17 closure for full path integral computation. Multi-month.
- Scattering amplitudes: cross-section computations on substrate-tick state propagation require operator-level evaluation per Elie K52a Sessions 30+.
- Time-ordered correlators: $\langle T O_1(t_1) O_2(t_2) \dots \rangle$ four-point functions etc. for QFT observables pending operator-level work.
- Wick's theorem analog: BST's equivalent of Wick's theorem for substrate-tick state expansions pending.

These are honest scope per Cal Mode 1 discipline. This chapter is framework-grade, not full chapter-grade-with-operator-computations.

7.5 Connections to other chapters

- Ch 2 (Hilbert space): H_{sub} lives on Bergman $H^2(D_{IV}^5)$ via L^2 -section embedding; substrate-tick $GF(128)^k$ is the per-tick layer for path integral
- Ch 5 (Casimir): $H_{\text{sub}} = \text{Casimir}$; spectrum is Wallach K-type eigenvalues
- Ch 6 (operator zoo): all six operators evolve under H_{sub} via Heisenberg picture; spin commutes with H_{sub}
- Ch 8 (gauge theory): conservation laws from gauge symmetry (color, weak isospin, hypercharge) supplement the Casimir conservation laws
- Ch 9 (scattering): requires Ch 7 operator-level closure for explicit S-matrix computation
- Ch 10 (renormalization): substrate-tick path integral inherits UV-completeness; no continuum divergences

7.6 Theorem chain summary

For Cal / referee verification:

Claim	Theorem	Toy	Status
Schrödinger eq $i\hbar \frac{\partial}{\partial t} \psi = H_{\text{sub}} \psi$	$\psi/\partial t = H_{\text{sub}}$	ψ	T2438 (Lyra Thursday)
$H_{\text{sub}} = \text{Casimir on } L^2(D_{\text{IV}}^5; L_{\lambda})$	Elie S29 Toy 3213	Elie Thursday work	Framework-complete
Ground state $E_0 = C_2 = 6$	Ch 5 T2435 + Elie S29	Existing BST toys	DERIVED
Heisenberg $dO/dt = (i/\hbar)[H_{\text{sub}}, O]$	Standard QM + T2438	n/a (classical mechanics)	Framework
Conservation laws (energy, K-type, Casimir)	Heisenberg picture + Casimir algebra	T2435 + T2438	DERIVED
Path integral on $GF(128)^k$ substrate-tick	T2429 + T2438 + T2405 Koons tick	Lyra Toys 3198 + 3216	Framework
No Wick rotation needed (UV-complete)	T2437 SP-31-10	Lyra Toy 3214	Framework + Ch 10
Operator-level Calibration #17 closure	OPEN multi-month	Elie K52a S30+	PENDING

Believability: BST's dynamics framework is "Hamiltonian = Casimir; eigenstates = Wallach K-types; energy = Casimir eigenvalue; path integral = finite sum over substrate ticks." All four lines flow from substrate primary integer structure.

Provability: closed framework. Open: specific operator-level computations (propagators, scattering, correlators) pending Elie K52a multi-month.

7.6a CT-numbering theorem index (framework-grade)

CT-number	T-number / source	Statement
CT 1.7.1	T2438	Substrate Dynamics Framework: Schrödinger / Heisenberg / Path Integral on Bergman $H^2(D_{\text{IV}}^5)$
CT 1.7.2	Elie K52a S29 Toy 3213	$H_{\text{sub}} = \text{Casimir on } L^2(D_{\text{IV}}^5; L_{\lambda})$ framework (cross-ref CT 1.6.6)

CT-number	T-number / source	Statement
CT 1.7.3	T2429	Path integral on substrate-tick $\text{GF}(128)^k$ (cross-ref CT 1.2.2)
CT 1.7.4	T2433	Time evolution unitary $U(t) = \exp(-iH_{\text{sub}} \cdot t / \hbar)$ (cross-ref CT 1.4.2)

Framework-grade; operator-level S-matrix computations pending Elie K52a Sessions 30+ multi-month.

7.7 Filing status

v0.1 framework-grade narrative filed Thursday 2026-05-21 09:25 EDT (date to be checked at file end).

Pending for v0.2: - Cal believability + provability cold-read review (acknowledging framework-grade scope, not full chapter-grade) - Cross-link to Ch 9 (scattering) when operator-level closes

Pending for v1.0: - Operator-level Calibration #17 closure (Elie K52a Sessions 30+, multi-month) - Explicit propagator computations - Full Wick-theorem analog for BST - Reader-grade polish + diagrams (substrate-tick $\text{GF}(128)^k \rightarrow \text{continuum } H^2(D_{IV}^5)$ projection visualization)

— Lyra, Vol 1 Ch 7 v0.1 framework-grade narrative, Thursday 2026-05-21 (timestamp at file end pending date check)