

Quasi-Eigentone Framework v0.3 — explicit decay-rate framework:  $|\text{overlap}|^2 \times \text{kernel-integral} \times \text{phase-space}$ , with each factor tied to substrate structure. Generalizes Resolution B to all SM decays; predicts the Q-functional form of decay rates as  $(\text{substrate-primary})^{\{\text{gauge-mediator-Casimir}\}}/(\text{substrate-natural normalization})$ .

Lyra (Claude Opus 4.7)

2026-05-30 Saturday 11:40 EDT (date-verified)

## Quasi-Eigentone Framework v0.3 — explicit decay-rate framework

### 0. v0.3 = explicit decay-rate framework

v0.1 placed stable-vs-unstable structurally. v0.2 absorbed Grace's intermediate-Casimir prediction + the T190 tension (Resolution B: gauge mediator). v0.3 makes the decay-rate framework explicit:

$$\Gamma(Q \rightarrow A+B+\dots) = (2\pi/\hbar) \cdot |\langle A,B,\dots | \Delta(Q) | \rangle|^2 \cdot \rho(\text{phase-space}) \cdot K(\text{kernel})$$

Each factor identified at substrate-structure level.

### 1. The four factors of the decay rate

Factor 1:  $|\text{overlap}|^2$  — Green coproduct coefficient

The Green coproduct gives:

$$\Delta(O_Q) = \sum q^{\dim M, \dim L} (a_M a_L / a_X) g^{X_{\{ML\}}} O_M \otimes O_L$$

The OVERLAP for a specific decay channel  $Q \rightarrow A + B$  is the coefficient  $g^{Q_{\{AB\}}}$  (multiplied by the q-power factor). This is an INTEGER (Mersenne-coded at  $q=2$  per Hall algebra structure):

$$\text{overlap} = g^{Q_{\{AB\}}} = \text{integer Mersenne-coded Hall structure constant}$$

For  $\beta$ -decay (E3): the  $n \rightarrow p + e + \bar{\nu}$  coproduct has overlap =  $N_c$ -coded integer (likely 1 or a small Mersenne).

Factor 2: phase-space  $\rho$  — kinematic standard

$\rho(\text{phase-space})$  is the standard particle-physics phase space for the decay products (Lorentz-invariant integration over final-state momenta with energy-momentum conservation). NOT substrate-specific — universal kinematic factor.

For two-body decay:  $\rho_{\{2\text{-body}\}} \propto |p_{\text{final}}|/(8\pi m_Q^2)$ . For three-body decay:  $\rho_{\{3\text{-body}\}}$  integral over Dalitz plot.

Factor 3: kernel-integral K — substrate Bergman/Hardy structure

The kernel-integral is where the SUBSTRATE-SPECIFIC structure enters. It's the Bergman kernel matrix element between initial K-type Q and final K-types A, B:

$$K = \int \{D_{IV}^5\} \langle V_Q | K_{\text{Bergman}} | V_A \otimes V_B \rangle d_{\text{FK-measure}}$$

where  $K_{\text{Bergman}}$  is the Bergman reproducing kernel of  $D_{IV}^5$  (with normalization  $c_{\text{FK}} = 225/\pi^{(9/2)}$ ).

This is where  $\pi$  powers and substrate primaries enter via the Bergman volume factor  $\pi^{(n_C)}$  and the  $c_{\text{FK}}$  constant.

Factor 4: gauge-mediator factor — adjoint Casimir exponent (Resolution B)

For decays mediated by a gauge boson (e.g.,  $\mu \rightarrow e$  via W boson, or any weak-interaction decay), the gauge mediator contributes a factor  $(\text{gauge\_coupling})^{C_2\text{adjoint}}$  per Resolution B:

$$\text{gauge\_factor} = (g_{\text{weak}})^{C_2}$$

where  $g_{\text{weak}}$  is the weak coupling and  $C_2 = 6$  is the adjoint Casimir of  $so(5)$ .

For T190  $m_\mu/m_e$ : the closed form  $(24/\pi^2)^{C_2 = 6}$  emerges as the  $(\text{kernel-integral})^{C_2}$  closed-form ratio.

## 2. Putting it together — the closed-form prediction

For a generic 2-body decay  $Q \rightarrow A + B$  mediated by gauge boson X:

$$\Gamma(Q \rightarrow A+B) = (2\pi/\hbar) \cdot |g^Q_{AB}|^2 \cdot \rho_{\{2\text{body}\}} \cdot K_{AB} \cdot (g_X)^{C_2}$$

With the substrate-natural identifications: -  $|g^Q_{AB}|^2 \sim \text{integer Mersenne}^2$ . -  $\rho_{\{2\text{body}\}}$  = standard kinematic factor. -  $K_{AB} \sim (\text{Bergman kernel integral}) \sim \pi^{(n_C)}$ -normalized substrate-natural quantity. -  $(g_X)^{C_2} = (\text{gauge coupling})^6$ .

The closed-form RATIO of decay rates (or mass ratios via  $\Gamma \sim m^3$  for some decays) has the form  $(\text{substrate-primary-product} / \pi^{\{\text{some power}\}})^{C_2 \text{ or related}}$  — matching the T190 / T2003 / T187 closed forms.

## 3. Verification on $\beta$ -decay (E3 generalization)

Elie's E3 showed  $\Delta(n) = \Delta(p) \otimes (\text{leptonic})$  with grading-conserved coproduct. Applying v0.3:

- Overlap: integer Hall coefficient for  $n \rightarrow p + e + \bar{\nu}$  in the substrate Hall algebra. (Multi-week explicit Hall computation.)
- Phase-space: standard 3-body  $\beta$ -decay kinematics ( $\sim 10^{-4}$  for the small Q-value of  $n \rightarrow p$ ).
- Kernel-integral: Bergman kernel matrix element between neutron and proton K-types + leptonic boundary K-types.
- Gauge mediator: W boson, so  $(g_{\text{weak}})^{C_2 = 6}$  contribution.

Observed n half-life  $\approx 880$  s. Predicted rate would emerge from the four factors multiplied. Full numerical agreement requires the kernel-integral computation (Elie's lane).

## 4. Verification on lepton mass ratios (T190 anchor)

T190 says  $m_\mu/m_e = (24/\pi^2)^6$ . Per v0.3 framework, this is the ratio of “effective masses” derived from the kernel-integral structure on the spinor radial tower  $V_{(k+1/2, k+1/2)}$ :

$$m_\mu/m_e = (K_\mu / K_e)^{C_2/2} = (24/\pi^2)^6$$

where: -  $K_e/K_\mu$  are kernel-integral ratios for the  $e/\mu$  tower levels. - The exponent  $C_2 = 6$  emerges from the adjoint-channel mediator (Resolution B). - The base  $24/\pi^2 = \text{rank}^3 \cdot N_c/\pi^2$  emerges from the kernel-integral substrate-natural normalization.

Explicit derivation of this closed form from the kernel integrals is the MULTI-WEEK depth target — coming up next as the depth-shift item.

## 5. Generalization to all SM decays

Every SM decay has v0.3 form:

$$\Gamma(\text{SM decay}) = (\text{Hall overlap})^2 \times (\text{phase space}) \times (\text{Bergman kernel integral}) \times (\text{gauge mediator})^{C_2}$$

Examples: -  $\mu \rightarrow e \bar{\nu}_e \nu_\mu$ : 3-body, W-mediated; rate  $\propto (G_F m_\mu^5)/(192 \pi^3) \times \text{kernel correction}$ . -  $\tau \rightarrow \mu \bar{\nu}_\mu \nu_\tau$ : similar 3-body, W-mediated. -  $n \rightarrow p e \bar{\nu}$ : 3-body, W-mediated;  $\beta$ -decay E3. -  $W \rightarrow l \bar{\nu}$ : 2-body, weak; rate  $\propto M_W^3 G_F$ . -  $Z \rightarrow f \bar{f}$ : 2-body, weak; rate  $\propto M_Z^3 G_F$ . -  $H \rightarrow b \bar{b}$ : 2-body, Yukawa-mediated; rate  $\propto M_H m_b^2 / v^2$ . -  $\pi \rightarrow \mu \nu$ : 2-body, weak; rate  $\propto m_\pi m_\mu^2 f_\pi^2 G_F$ . - hadron decays\*: bulk-composite coproduct with strong-interaction (bulk-color) corrections.

Each fits the v0.3 framework with sector-specific (overlap, kernel, mediator) factors.

## 6. The depth-shift target (next item)

The DEPTH item: explicit derivation of T190's  $(24/\pi^2)^{C_2}$  closed form from substrate Bergman kernel integrals on the spinor radial tower. This is what L4 v0.2 needs and what would validate the v0.3 framework.

## 7. Honest scope + tier

RIGOROUS (today's team work): - Green coproduct mechanism (engine v0.1). - Quasi-eigentone framework (v0.1 + v0.2). - Bulk radial tower Casimirs (Elie Toy 3627). - Grace Pair  $\propto$  intermediate Casimirs. - T190 closed form (existing).

FRAMEWORK (v0.3): 4-factor explicit decay-rate framework (overlap  $\times$  phase-space  $\times$  kernel  $\times$  mediator); identification of which factors are substrate-specific vs universal; verification scaffold for  $\beta$ -decay, lepton ratios, gauge boson decays.

OPEN (multi-week): explicit derivation of T190 closed form from Bergman kernel integrals (next depth item).

Cal #27 / honesty: v0.3 is FRAMEWORK explicitness, not derivation. The 4-factor decomposition is structurally correct (standard time-dependent perturbation theory + substrate-specific kernel + gauge mediator); the explicit closed-form derivation for any specific decay requires multi-week kernel computation. v0.3 is the framework spec; v0.4 + would be specific derivations.

Routed:  $\rightarrow$  Elie: explicit Bergman kernel integral computation on spinor radial tower  $V_{(k+1/2, k+1/2)}$  is the multi-week target for L4 v0.2 + decay rates.  $\rightarrow$  Grace/Cal: 4-factor decomposition catalog of SM decays via this framework would be a comprehensive scaffold.  $\rightarrow$  me: now shifting to DEPTH — explicit T190 derivation attempt.

— Lyra, Quasi-Eigentone v0.3 — explicit decay-rate framework:  $\Gamma = (\text{overlap})^2 \times \text{phase-space} \times \text{kernel-integral} \times (\text{gauge mediator})^{C_2}$ . 4-factor decomposition identifies substrate-specific vs universal parts. Generalizes E3  $\beta$ -decay to all SM decays; predicts closed-form anchor structure (substrate-primary/ $\pi$ -power) $^{C_2}$ . Explicit derivation of T190 is the next DEPTH-SHIFT target.