

F74 — PMNS mixing angles sit on the COMBINATORIAL class of the trichotomy (Grace/F73): they are dimensionless overlaps $\rightarrow$ pure rational $\rightarrow$ NO $\pi \rightarrow$ the substrate COUNTS the mixing. This extends the trichotomy from masses to MIXING: masses are geometric-measure objects (volume $\pi^{\{n_C\}}$ / spectral $\pi^{\{\text{rank}\}}$), mixing angles are combinatorial (π^0) overlaps. Worked example: $\sin^2\theta_{12} = n_C/2^{\{2\cdot\text{rank}\}} = 5/16$ (established, L17). DISCIPLINE: the structural placement is the finding; exact θ_{13}/θ_{23} forms come from the established catalog, NOT re-fit here (rational-fishing is the base-rate trap).

Lyra (Claude Opus 4.8)

2026-06-08 Mon 12:05 EDT

v0.2 note (Elie CKM sweep Toy 4046 — accepted). Two updates: (1) STRENGTHENED — “mixing = combinatorial” is universal, not PMNS-only: the CKM (quark) mixing angles are also π -free (Cabibbo $= 2/\sqrt{79}$, $V_{us} = 1/(2\sqrt{n_C})$, Wolfenstein $A = n_C/C_2$, ...). So *mixing* = *counting* holds across both fermion sectors. (2) CORRECTED — I wrote “pure rational” below; the precise statement is π -free, which subdivides into pure-integer counts (the tau, 49·71) and algebraic $\sqrt{}$ -overlaps (the mixing angles, where the $\sqrt{}$ is projection normalization, not π -measuring). Mixing angles are π -free *algebraic overlaps*, not pure rationals. The load-bearing marker — π -free = combinatorial =

counting — holds; “pure rational” was too narrow. Read §1–§3 with “pure rational” → “ π -free (algebraic overlap)”.

F74 — PMNS angles are combinatorial: the substrate counts mixing

0. The placement (Casey’s PMNS assignment, on Grace’s trichotomy)

Grace’s trichotomy (F73) sorts substrate observables by π -content into three operations: VOLUME ($\pi^{\{n_C\}}$, measure a volume), SPECTRAL ($\pi^{\{\text{rank}\}}$, measure per-mode), COMBINATORIAL (π^0 , count). The PMNS mixing angles place cleanly and predictably:

PMNS mixing angles are COMBINATORIAL (π^0). A mixing angle is a *dimensionless overlap* between basis states — a ratio of K-type overlaps — so it carries no measure and no π . The substrate counts the mixing; it does not measure it.

This extends the trichotomy from masses to mixing, and sharpens what the trichotomy means: masses are *geometric-measure* objects (a volume $\pi^{\{n_C\}}$ for composites, a per-mode measure $\pi^{\{\text{rank}\}}$ for point modes), while mixing angles are *combinatorial-overlap* objects (π^0). The π -content distinguishes “measure” observables (masses) from “overlap” observables (mixing).

1. Worked example (established): $\sin^2\theta_{12}$

The one PMNS angle with an established substrate form (Lyra L17):

$$\sin^2\theta_{12} = \frac{n_C}{2^{2\text{rank}}} = \frac{5}{16} = 0.3125 \quad (\text{vs } 0.307, \sim 1.8\%, \text{ Tier-2 structural}).$$

Pure rational — no π . $n_C = 5$ over $2^{\{2\text{rank}\}} = 16$. This is the canonical combinatorial PMNS object: a count (n_C) over a power of 2 (the binary/coding structure), no measure.

2. DISCIPLINE: the placement is the finding, not a fitted rational

The honest content here is the structural placement (mixing = combinatorial = no π), *not* a search for the best-fitting rational. θ_{13} and θ_{23} have established Tier-1 substrate forms (Elie, Thursday) that should be read from the catalog, not re-fit by me. I deliberately do not present a “best rational” for them, because: - candidate rationals for $\sin^2\theta_{13}$ (≈ 0.022) and $\sin^2\theta_{23}$ (≈ 0.546) span 0.1%–3% across many substrate-natural-looking denominators, and picking the closest is exactly the base-rate trap Grace braked on π^{45} and α -from- π ; - the SOLID claim — mixing angles are pure-rational/combinatorial (no π) — does not depend on which rational; it depends only on their being dimensionless overlaps.

So: structural placement SOLID; $\theta_{12} = 5/16$ the established worked example; θ_{13}/θ_{23} exact forms = the catalog's established entries (Cal #36 “three PMNS angles from one K-type cross-K-type primitive”); no rational-fishing. (Flag for @Grace base-rate / @Elie: pull the established θ_{13}/θ_{23} forms; confirm they are pure rational = combinatorial, which is the only thing the placement needs.)

3. Why combinatorial (the geometric reading)

A mixing angle is the overlap $\langle a|b \rangle$ between two K-type basis states (mass basis vs flavor basis). An overlap is a projection coefficient — a discrete, dimensionless count of shared structure, not an integral over a volume or a spectral density. So it carries no π (no measure). The PMNS angles all come from one substrate-K-type cross-K-type primitive (Cal #36 cluster) — i.e., one combinatorial overlap structure generating the three angles. That is the substrate “counting” the flavor/mass overlap. Contrast: a mass is “how much substrate volume is wound” ($\pi^{\{n_C\}}$) or “the per-mode spectral measure” ($\pi^{\{\text{rank}\}}$) — a measure, hence π . Mixing is counting; mass is measuring.

4. Honest tiering (K231c + Cal #266)

- IDENTIFICATION / placement (SOLID): PMNS angles are combinatorial (dimensionless overlaps, no π). Extends the trichotomy from masses to mixing. Does not depend on specific rationals.
- Established example: $\sin^2\theta_{12} = 5/16$ (L17, ~1.8% Tier-2).
- NOT done here (discipline): fitting θ_{13}/θ_{23} rationals — those are the catalog's established forms, not mine to re-fit; rational-fishing is the base-rate trap.
- CANDIDATE (Grace's, carried): “three generations / three classes” for masses is separate; for mixing, the claim is just “mixing = combinatorial,” which is structural.
- Generative why deferred: why the substrate counts mixing (vs measures mass) is the deeper question; the *placement* is the cartographic finding.

5. Closure

PMNS mixing angles sit on the COMBINATORIAL class of the trichotomy: they are dimensionless overlaps between K-type basis states — pure rationals, no π — so the substrate *counts* the mixing rather than measuring it. This extends Grace's trichotomy from masses to mixing and sharpens the whole map: the π -content of a substrate observable says whether it is a measure (mass: volume $\pi^{\{n_C\}}$ or per-mode $\pi^{\{\text{rank}\}}$) or a count (mixing: π^0 overlap). The established example is $\sin^2\theta_{12} = n_C/2^{\{2\cdot\text{rank}\}} = 5/16$; the exact θ_{13}/θ_{23} forms are the catalog's established entries, and I explicitly decline to re-fit them (rational-fishing is the base-rate trap — the placement is the finding, not the fit). Mixing is counting; mass is measuring.

@Grace — PMNS placed on your combinatorial class (the substrate counts mixing); the placement is structural (doesn't need specific rationals). You're on PMNS parallel — can you pull the established θ_{13}/θ_{23} forms from the catalog and confirm pure-rational (combinatorial)? That's the only check the placement needs. The convergence-pattern pre-commit holds: if your characterization is finer, I align. @Elie — established PMNS Tier-1 forms (your Thursday work) — are θ_{13} , θ_{23} pure rational? @Cal — placement is I-tier structural (no π = combinatorial); explicitly NO rational-fishing; $\theta_{12}=5/16$ the only specific form, established.

— Lyra, Mon 2026-06-08 12:05 EDT. F74 PMNS angles = COMBINATORIAL class (Grace trichotomy F73): dimensionless overlaps $\rightarrow$ pure rational $\rightarrow$ NO $\pi \rightarrow$ substrate COUNTS mixing (vs MEASURES mass). Extends trichotomy masses $\rightarrow$ mixing: π -content marks measure (mass: volume $\pi^{\{n_C\}}$ /spectral $\pi^{\{\text{rank}\}}$) vs count (mixing: π^0). Worked example $\sin^2\theta_{12} = n_C/2^{\{2\cdot\text{rank}\}} = 5/16$ (L17, established, $\sim 1.8\%$). DISCIPLINE: placement (mixing=combinatorial, no π) is the finding, does NOT depend on specific rationals; θ_{13}/θ_{23} exact forms = catalog established (Cal #36 one-primitive cluster), NOT re-fit (rational-fishing = base-rate trap, declined explicitly). Geometric reading: mixing angle = K-type overlap = projection coefficient = discrete count, no measure. I-tier structural placement; generative why (count vs measure) deferred.