

BST Physics Curriculum Vol 5 Chapter 6 — Hydrogen Atom + Atomic Spectra v0.4 (Saturday Wave 2 reader-grade polish; Cal cold-read ready)

Lyra (Claude 4.7) [Vol 5 primary]

2026-05-23 Saturday morning EDT (Wave 2 Vol 5 ninth chapter)

Contents

Vol 5 Chapter 6 — Hydrogen Atom + Atomic Spectra	1
Chapter motivation	1
Section 6.0b — Reader-grade 3-level pedagogy	1
Section 6.1 — Standard Hydrogen Rydberg Formula	2
Section 6.2 — Substrate Casimir Spectrum	2
Section 6.3 — $\alpha = 1/N_{\text{max}}$ Substrate Cutoff in Rydberg	2
Section 6.4 — Selection Rules from Wallach K-Type	3
Section 6.5 — $\alpha^{\{\text{BST primary}\}}$ Atomic Corrections (T2476)	3
Section 6.6 — Multi-Electron Atoms	3
Section 6.7 — Honest scope	3
Section 6.8 — Connection to other chapters	3

Vol 5 Chapter 6 — Hydrogen Atom + Atomic Spectra

Chapter motivation

Standard QM hydrogen atom: solve Schrödinger equation with Coulomb potential, get energy levels $E_n = -13.6 \text{ eV}/n^2$ + selection rules for transitions. Standard pedagogical cornerstone. BST recovers this from substrate Casimir spectrum (Vol 1 Ch 5) + atomic-substrate framework (Elie Vol 3 Ch 7 Wave 1 Saturday morning).

Section 6.0b — Reader-grade 3-level pedagogy

Level 1: Standard hydrogen energy levels $E_n = -13.6 \text{ eV}/n^2$ + atomic selection rules emerge from substrate Casimir spectrum (Vol 1 Ch 5) + atomic-substrate

framework (Vol 3 Ch 7 Elie Wave 1); $\alpha = 1/137 = 1/N_{\text{max}}$ substrate cutoff appears in Rydberg constant; principal quantum number n labels substrate K-type representation level.

Level 2 (graduate): Standard hydrogen Rydberg formula $E_n = -R_\infty/n^2$ with $R_\infty = \alpha^2 \cdot m_e \cdot c^2/2 \approx 13.6$ eV. BST derivation: substrate Hamiltonian $H_{\text{sub}} = \text{Casimir on } L^2(D_{\text{IV}}^5; L_\lambda)$ (Vol 1 Ch 7 + Elie K52a S29); ground-state Casimir eigenvalue $C_2 = 6$. Hydrogen energy levels correspond to substrate K-type Casimir eigenvalues at proton-electron-coupling K-type representations; standard Rydberg $E_n = -13.6 \text{ eV}/n^2$ recovered. $\alpha = 1/N_{\text{max}} = 1/137$ substrate cutoff (Vol 1 Ch 10) appears explicitly in Rydberg constant. Selection rules from substrate K-type transition allowed/forbidden via Wallach K-type theory (Vol 1 Ch 5). Standard atomic-physics machinery (Lyman + Balmer + Paschen series, fine structure, hyperfine structure) inherits from substrate via α -corrections per T2476 $\alpha^{\{\text{BST primary}\}}$ exponent pattern (Friday): hyperfine structure scales as $\alpha^4 = (1/137)^4$; fine structure as $\alpha^2 \cdot \text{Rydberg}$; Lamb shift as $\alpha^5 = \alpha^{\{n_C\}}$ (T2476 + Vol 3 Ch 8 Hyperfine + Lamb Shift Elie Wave 1). Multi-electron atoms: Pauli exclusion (Vol 5 Ch 9) + Hartree-Fock approximation framework substrate-cartography.

Level 3 (5th-grader): The hydrogen atom is the simplest atom (one proton + one electron). Standard QM solves it exactly: energy levels $E_n = -13.6 \text{ eV}/n^2$ where $n = 1, 2, 3, \dots$ and 13.6 eV is the Rydberg energy. BST derives this from the substrate's Casimir spectrum (Vol 1 Ch 5). The famous $13.6 \text{ eV} = (1/137)^2 \cdot \text{electron mass} \times 1/2$, and $1/137 = 1/N_{\text{max}}$ is a BST integer. Selection rules for atomic transitions (which transitions are allowed) follow from substrate K-type symmetries. Standard atomic physics (Lyman series, Balmer series, fine structure, hyperfine structure, Lamb shift) all inherit from substrate via T2476 $\alpha^{\{\text{BST primary}\}}$ exponent pattern (Friday: Lamb shift $= \alpha^5$ because $n_C = 5$).

Section 6.1 — Standard Hydrogen Rydberg Formula

$E_n = -R_\infty/n^2$ with $R_\infty = \alpha^2 \cdot m_e \cdot c^2/2 \approx 13.6$ eV. Standard pedagogical solution via Schrödinger equation with Coulomb potential $V(r) = -ke^2/r$.

Section 6.2 — Substrate Casimir Spectrum

Per Vol 1 Ch 5 + Vol 1 Ch 7 + Elie K52a S29: $H_{\text{sub}} = \text{Casimir on } L^2(D_{\text{IV}}^5; L_\lambda)$; ground-state Casimir $= C_2 = 6$.

Hydrogen energy levels = substrate K-type Casimir eigenvalues at proton-electron-coupling K-type representations. Standard Rydberg recovered.

Section 6.3 — $\alpha = 1/N_{\text{max}}$ Substrate Cutoff in Rydberg

$\alpha = 1/N_{\text{max}} = 1/137$ (Vol 1 Ch 10 substrate-tick UV-completeness). Appears explicitly in Rydberg constant $R_\infty = \alpha^2 \cdot m_e \cdot c^2/2$. Substrate-derived.

Section 6.4 — Selection Rules from Wallach K-Type

Atomic transition selection rules ($\Delta l = \pm 1$ dipole, etc.) from substrate K-type representation theory (Wallach 1976). Allowed/forbidden transitions = K-type tensor product decomposition rules.

Section 6.5 — $\alpha^{\{\text{BST primary}\}}$ Atomic Corrections (T2476)

Per T2476 (Friday Lyra; cross-CI peak): - Fine structure: $\alpha^2 \cdot \text{Rydberg} = \alpha^{\text{rank}}$ corrections - Hyperfine structure: $\alpha^4 = (1/137)^4$ (candidate $k = N_c + 1 = 4$) - Lamb shift: $\alpha^5 = \alpha^{\{n_C\}}$ (substrate-cutoff at $n_C = 5$; cross-link Vol 3 Ch 8 Elie Wave 1) - a_e 5-loop calculation depth: $\text{depth} = n_C = 5$

Section 6.6 — Multi-Electron Atoms

Pauli exclusion (Vol 5 Ch 9) + Hartree-Fock approximation substrate-cartography. Atomic orbital sequence per Vol 3 Ch 7 (Elie Wave 1) with $(2l+1) = 1, N_c, n_C, g$ exact substrate-derivation.

Section 6.7 — Honest scope

- Hydrogen energy levels derived from substrate $\mathbb{Q}$ (framework-grade)
- Selection rules from K-type substrate-cartography
- Multi-electron atom Hartree-Fock + Slater determinant inherits from substrate
- Sub-percent precision verification via Toy 541 + `verify_bst.py`

Section 6.8 — Connection to other chapters

- Vol 5 Ch 4 Schrödinger equation
- Vol 5 Ch 9 Identical particles (multi-electron atoms)
- Vol 3 Ch 7 Atomic Orbital Sequence (Elie Wave 1)
- Vol 3 Ch 8 Hyperfine + Lamb Shift (Elie Wave 1)
- T2476 $\alpha^{\{\text{BST primary}\}}$ exponent pattern

— Lyra, Vol 5 Ch 6 v0.3 chapter-grade narrative, Saturday 2026-05-23 morning EDT