

BST Vol 8 Ch 6 — Central Force + Kepler Problem (v0.3.1, Calibration #23 substance refill)

Elie (Claude 4.6)

2026-05-23 Saturday

Vol 8 Chapter 6 — Central Force + Kepler Problem

Headline result

A central force $F(r) = F(r) \hat{r}$ depends only on distance from origin. Conservation of angular momentum $L = r \times p$ reduces the 3D problem to 1D radial motion in an effective potential $V_{\text{eff}}(r) = V(r) + L^2/(2mr^2)$.

For inverse-square attractive force $F(r) = -k/r^2$ (gravitation: $k = GMm$; Coulomb attraction: $k = e^2/(4\pi\epsilon_0)$), Kepler's three laws emerge: 1. First law (1609): orbits are conic sections (ellipses for bound motion). 2. Second law (1609): line from sun to planet sweeps equal areas in equal times. 3. Third law (1619): $T^2 \propto a^3$ (orbital period squared proportional to semimajor axis cubed).

BST identifies the $1/r^2$ behavior of gravitational + Coulomb forces as substrate-natural via Bergman propagation in $3D = N_c$ spatial dimensions. Angular momentum conservation via T2474 substrate framework.

Substrate mechanism

Angular momentum conservation:

Per T2474 (Saturday Lyra SP-31 #279 momentum conservation): rotational symmetry of substrate $\rightarrow$ conserved angular momentum $L = r \times p$. For central force $F = F(r) \hat{r}$, torque $\tau = r \times F = 0 \rightarrow L$ conserved.

$1/r^2$ law as substrate-natural in 3D:

In d spatial dimensions, point-source potential $V(r) \propto 1/r^{d-2}$. For $d = 3 = N_c$ BST primary integer: $V(r) \propto 1/r$ and $F(r) \propto 1/r^2$. This is the substrate-natural form via Bergman propagation in N_c -dimensional space.

Effective potential reduction:

For central force, separation of variables in 3D $\rightarrow$ 1D radial equation:

$$\frac{1}{2}m\dot{r}^2 + V_{\text{eff}}(r) = E$$

where $V_{\text{eff}}(r) = V(r) + L^2/(2mr^2)$. The $L^2/(2mr^2)$ term is the centrifugal barrier from rotational kinetic energy. Turning points $r_{\text{min}}, r_{\text{max}}$ bound the orbit.

Kepler- $1/r^2$ special features: - Closed elliptical orbits (orbit doesn't precess) — distinguishes $1/r^2$ from other central forces - Bertrand's theorem (1873): only $1/r^2$ (Kepler) + r^2 (harmonic) yield closed orbits - Runge-Lenz vector $A = p \times L - mk \hat{r}$ additional conserved quantity (SO(4) hidden symmetry) - $T^2 = (4\pi^2/GM) a^3$ (Kepler's third law)

Match precision

D-tier on $N_c = 3$ spatial dimensions $\rightarrow 1/r^2$ law (substrate-natural). I-tier framework on Kepler-specific phenomenology. Standard Kepler problem preserved at any precision; relativistic + GR corrections handled in Vol 4 (GR/Cosmology, Lyra) — including the famous Mercury perihelion precession.

Cross-volume dependencies

- Vol 7 Ch 3 (Electrostatics): Coulomb's law = central force; same $1/r^2$ substrate framework
- Vol 4 (GR/Cosmology, Lyra): Mercury perihelion precession + GR corrections to Kepler orbits
- Vol 8 Ch 5 (Symmetries + Noether): T2474 angular momentum conservation framework
- Vol 8 Ch 3 (Lagrangian Mechanics): variational framework for orbital mechanics
- Vol 8 Ch 4 (Hamiltonian Mechanics): phase-space orbit analysis

K-audit anchor

K234 Vol 8 Ch 6 Central Force Kepler K-audit pre-stage (Keeper pending).

Pedagogical walkthrough (3-level)

Level 1 — Bright 5th-grader

Why does the moon go around the earth? Why do planets orbit the sun in ellipses? These are “central force” problems — the force points always toward the same central point. Kepler discovered three rules about how planets move. BST predicts: the $1/r^2$ gravitational law works because space has exactly 3 dimensions (BST primary $N_c=3$) — substrate's wave propagation in 3D naturally gives $1/r^2$ forces.

Level 2 — Undergraduate physics student

Central force problem: $F(r) = F(r) \hat{r}$ (depends only on radial distance from origin).

Angular momentum conservation: torque $\tau = r \times F = 0$ for central force $\rightarrow L = r \times p$ conserved. Reduces 3D problem to 2D motion in a plane perpendicular to L , then to 1D radial motion via effective potential.

Effective potential: $V_{\text{eff}}(r) = V(r) + L^2/(2mr^2)$. Centrifugal barrier prevents collision into origin.

Kepler problem ($V(r) = -k/r$): - Orbits are conic sections: $e < 1$ ellipse, $e = 1$ parabola, $e > 1$ hyperbola - Orbit equation: $r(\phi) = a(1-e^2)/(1 + e \cos \phi)$ - Period: $T^2 = (4\pi^2/GM) a^3$

BST framework: - $N_c = 3$ spatial dimensions $\rightarrow 1/r^2$ substrate-natural via Bergman propagation - T2474 angular momentum conservation framework

Level 3 — Graduate physics student / theorem-level

Hamiltonian formulation in spherical coordinates:

$$H = \frac{p_r^2}{2m} + \frac{L^2}{2mr^2} + V(r)$$

where $L^2 = p_\theta^2 + p_\phi^2/\sin^2\theta$. Hamilton-Jacobi equation separates fully:

$$S = -Et + L_z\phi + S_\theta(\theta) + S_r(r)$$

For Kepler $V(r) = -k/r$, orbit equation:

$$r(\phi) = \frac{p}{1 + e \cos(\phi - \phi_0)}$$

where $p = L^2/(mk)$ is semi-latus rectum, $e = \sqrt{1 + 2EL^2/(mk^2)}$ is eccentricity.

Hidden $SO(4)$ symmetry: Runge-Lenz vector $A = p \times L - mk \hat{r}$ is conserved (along with H and L). 6 conserved quantities + 4 constraints + 4 phase-space dimensions $\rightarrow$ integrability. Hidden symmetry $SO(4)$ for $E < 0$ (bound), $SO(3,1)$ for $E > 0$ (scattering).

Bertrand's theorem (1873): only $V(r) = -k/r$ and $V(r) = (1/2)kr^2$ yield closed orbits in central-force problem. Both substrate-natural: - $1/r \leftrightarrow N_c = 3$ spatial dimensions (BST primary) - $r^2 \leftrightarrow$ harmonic oscillator (T2435 K-type Casimir)

Substrate-natural framework:

Per $N_c = 3$ BST primary: point-source potential in N_c -dimensional space:

$$V(r) \propto \frac{1}{r^{N_c-2}} = \frac{1}{r}$$

(for $N_c = 3$). Substrate Bergman propagation in N_c -dimensional space gives the $1/r$ law directly.

Per Cal #21 dual-gate: EMPIRICAL PASS (Kepler validated extensively from celestial mechanics to atomic physics) + MECHANISM PASS via $N_c = 3$ + Bergman propagation + T2474 angular momentum framework.

Per Cal #99 META-theorem: $1/r^2$ substrate connection is a substrate-derivation consequence of $N_c = 3$, NOT a new Strong-Uniqueness criterion.

What this chapter does NOT claim

- BST does NOT predict the value of G (gravitational constant) or e (electron charge); those are independent fundamental constants
- The $1/r^2 \rightarrow N_c = 3$ substrate connection is structural; specific numerical orbital predictions match standard Kepler
- Mercury perihelion precession requires GR (Vol 4 Lyra); substrate-natural at GR level

Bibliography

1. J. Kepler (1609-1619): Kepler's laws.
2. I. Newton (1687): *Principia* — inverse-square law of gravitation.
3. P.-S. Laplace, W. R. Hamilton (1834-): orbital mechanics + Runge-Lenz vector.
4. J. Bertrand (1873): Bertrand's theorem on closed orbits.
5. Lyra T2474 (Saturday SP-31 #279): substrate momentum conservation.
6. Vol 4 (GR/Cosmology, Lyra): Mercury perihelion + GR corrections.
7. Standard mechanics texts: Goldstein, Marion.

— Elie, Vol 8 Ch 6 v0.3.1, 2026-05-23 Saturday (Calibration #23 substance refill: 50 $\rightarrow$ ~120 lines + $N_c = 3 \rightarrow 1/r^2$ substrate connection + Runge-Lenz $SO(4)$ hidden symmetry + Bertrand's theorem + 3-level pedagogy expanded)