

Vol 5 Chapter 5 — The Heisenberg Picture and the Path Integral

Keeper (author pass — deep math/physics revision)

2026-05-23 Saturday

Chapter 5 — The Heisenberg Picture and the Path Integral

Level 1 — one sentence

The Heisenberg picture and the Feynman path integral are two equivalent descriptions of substrate Zone 2 evolution: Heisenberg keeps K-type operators time-dependent under the substrate Casimir flow, and the path integral is what many substrate Koons ticks look like when their cumulative amplitudes concentrate on the classical-stationary trajectory via constructive interference.

Level 2 — graduate-physicist precision

5.1 The Heisenberg picture

In the Schrödinger picture (Chapter 4), states evolve and operators are fixed:

$$|\psi_S(t)\rangle = e^{-i\hat{H}t/\hbar}|\psi_S(0)\rangle, \quad \hat{A}_S = \text{const}$$

In the Heisenberg picture, operators evolve and states are fixed:

$$\hat{A}_H(t) = e^{i\hat{H}t/\hbar}\hat{A}_Se^{-i\hat{H}t/\hbar}, \quad |\psi_H\rangle = |\psi_S(0)\rangle = \text{const}$$

The two pictures give identical expectation values: $\langle\psi_S(t)|\hat{A}_S|\psi_S(t)\rangle = \langle\psi_H|\hat{A}_H(t)|\psi_H\rangle$. The choice is a matter of convenience.

The Heisenberg equation of motion:

$$i\hbar \frac{d\hat{A}_H}{dt} = [\hat{A}_H, \hat{H}] + i\hbar \frac{\partial \hat{A}_S}{\partial t}$$

For time-independent $\hat{A}_S$, this reduces to $i\hbar d\hat{A}_H/dt = [\hat{A}_H, \hat{H}]$.

5.2 Bohr correspondence: classical limit

For the substrate position operator $\hat{x}$ and momentum operator $\hat{p}$ (Chapter 2):

$$\frac{d\hat{x}_H}{dt} = \frac{1}{i\hbar}[\hat{x}, \hat{H}] = \frac{\hat{p}_H}{m}$$

$$\frac{d\hat{p}_H}{dt} = \frac{1}{i\hbar}[\hat{p}, \hat{H}] = -\nabla V(\hat{x}_H)$$

These are Hamilton's equations of classical mechanics, with $\hat{x}, \hat{p}$ replacing x, p . The classical Poisson bracket $\{A, B\}_{\text{PB}}$ corresponds to the quantum commutator $[\hat{A}, \hat{B}]/(i\hbar)$ — this is Bohr's correspondence principle.

BST substrate-mechanism: the classical mechanics of Volume 8 is the substrate's Heisenberg evolution at Scale 2 (many-tick averaged). The substrate's Zone 2 evolution at Scale 1 is full quantum mechanics; coarse-graining to Scale 2 recovers Hamilton's classical equations.

5.3 The Feynman path integral

Feynman 1948: the propagator $\langle x_f | e^{-i\hat{H}T/\hbar} | x_i \rangle$ (amplitude for a particle to go from x_i at $t = 0$ to x_f at $t = T$) is expressible as a “sum over paths”:

$$\langle x_f | e^{-i\hat{H}T/\hbar} | x_i \rangle = \int_{x_i}^{x_f} \mathcal{D}x(t) e^{iS[x(t)]/\hbar}$$

where $S[x(t)] = \int_0^T L(x, \dot{x}) dt$ is the classical action functional and $\mathcal{D}x$ denotes integration over all paths from x_i to x_f .

The classical path is the one that extremizes S (Euler-Lagrange equation). For $\hbar$ small, paths with $S \gg \hbar$ contribute oscillating phases that cancel; only paths near the classical-stationary path interfere constructively. As $\hbar \rightarrow 0$, the integral concentrates on the classical trajectory.

5.4 The path integral from substrate

The BST substrate-mechanism derivation: between $t = 0$ and $t = T$, the substrate completes $N = T/t_K$ Koons-tick commitment cycles. Each cycle includes Zone 1-4; the full amplitude factors as

$$\langle x_f | e^{-i\hat{H}T/\hbar} | x_i \rangle = \prod_{n=1}^N \langle x_n | e^{-i\hat{H}t_K/\hbar} | x_{n-1} \rangle$$

Insert complete sets of position and momentum eigenstates between each tick. After standard manipulation:

$$\langle x_f | e^{-i\hat{H}T/\hbar} | x_i \rangle = \int \prod_n dx_n dp_n e^{i \sum_n [p_n(x_n - x_{n-1}) - H(x_n, p_n)t_K]/\hbar}$$

In the continuum limit $t_K \rightarrow 0$, the discrete sum $\sum_n [p\dot{x} - H]t_K$ becomes the Lagrangian action $\int (p\dot{x} - H)dt = \int L dt$. The path integral emerges.

The substrate framework reads this transparently: the path integral is the *many-tick coherent sum* of substrate Zone 2 evolution amplitudes. Each tick contributes an infinitesimal action; the sum over all sequences of intermediate substrate K-type configurations is the path integral.

5.5 The classical-stationary path

For paths with action $S \gg \hbar$, neighboring paths differ in S by amounts that wash out the phase factor through destructive interference. Only paths within $\Delta S \sim \hbar$ of the stationary path survive.

The stationary path satisfies $\delta S = 0$ — the Euler-Lagrange equations of classical mechanics. The substrate-mechanism: at Scale 2 (many-tick averaged), the substrate's Zone 2 evolution is dominated by the path that minimizes the cumulative action, recovering classical-Lagrangian mechanics from the substrate's Schrödinger-picture quantum mechanics.

5.6 Worked example: free-particle propagator

For a free particle $\hat{H} = \hat{p}^2/2m$, the propagator is

$$\langle x_f | e^{-i\hat{H}T/\hbar} | x_i \rangle = \sqrt{\frac{m}{2\pi i \hbar T}} \exp \left[\frac{im(x_f - x_i)^2}{2\hbar T} \right]$$

The classical path is $x(t) = x_i + (x_f - x_i)t/T$, with classical action $S_{cl} = m(x_f - x_i)^2/(2T)$. The propagator equals the WKB amplitude $\sqrt{m/(2\pi i \hbar T)} e^{iS_{cl}/\hbar}$ — exact for the free particle (no higher-loop corrections).

Substrate reading: the free particle's substrate K-types are the translation-generated K-types; the path integral is the substrate's coherent sum across all intermediate K-type configurations, dominated by the classical path with action S_{cl} .

5.7 Wick rotation and the partition function

Rotate $t \rightarrow -i\tau$ (Wick rotation). The propagator becomes the imaginary-time evolution $e^{-\hat{H}\tau/\hbar}$ — the heat-kernel-like operator. The trace at “temperature” $\beta = 1/(k_B T)$:

$$Z(\beta) = \text{Tr} e^{-\beta \hat{H}} = \int dx \langle x | e^{-\beta \hat{H}} | x \rangle$$

This is the canonical partition function of statistical mechanics (Volume 6 Chapter 5). For the substrate on D_{IV}^5 :

$$Z(\beta) = \text{Tr}_{H^2(D_{IV}^5; L_\lambda)} e^{-\beta \mathcal{C}_{so(5,2)}}$$

This trace equals the heat kernel on D_{IV}^5 . Paper #9 “The Arithmetic Triangle of Curved Space” develops the Seeley-DeWitt expansion of this heat kernel through $k = 20$ with three theorems characterizing the arithmetic structure (speaking-pair period $n_C = 5$, column rule, two-source prime structure).

The Bohr-Sommerfeld / Wick-rotation connection: quantum mechanics in real time = statistical mechanics in imaginary time, both expressible as substrate Casimir traces on $L^2(D_{IV}^5; L_\lambda)$.

5.8 K-audit anchors

- Paper #9 “The Arithmetic Triangle of Curved Space” v10 (Lyra+Elie, current): heat kernel on D_{IV}^5 through $k = 20$; speaking-pair period $n_C = 5$ confirmed
- T531-T533: column rule, two-source prime structure, Kummer analog conjecture for D_{IV}^5 heat kernel
- Toy 639: $k = 16$ ratio = $-24 = -\dim SU(5)$ confirmed
- K52a Session 29 (Elie May 22, 2026): substrate Hamiltonian = Casimir framework-complete

Level 3 — 5th-grader accessibility

There are two ways to do quantum mechanics. One: keep the wave function moving in time, and the operators (position, momentum) stay put. Two (Heisenberg): keep the wave function frozen, and let the operators move in time. They agree. Feynman invented a third way: sum over every possible path the particle could take, with each path weighted by $e^{iS/\hbar}$. When $\hbar$ is small (it always is, for everyday objects), only the classical path counts — that's why a baseball travels in a parabola, not a cloud. In BST, the path integral is what happens when you add up many Koons ticks of the substrate's quantum evolution: substrate “tries” all paths, and the one with the smallest action wins by constructive interference. Trick: if you replace t with $-i\tau$ (Wick rotation), you get statistical mechanics for free. Same equation, different name.

What comes next

Chapter 6 develops the hydrogen atom — using the substrate machinery to recover the Bohr radius, ionization energy, and orbital spectrum.

Where to look this up

- Heisenberg picture: Sakurai and Napolitano, Ch 2
- Feynman path integral: Feynman and Hibbs, *Quantum Mechanics and Path Integrals*
- Heat kernel on symmetric spaces: Heat-Kernel Asymptotics from Symmetric Spaces literature
- Paper #9: BST “Arithmetic Triangle of Curved Space” v10, J. Spectral Theory submission target
- Toys 273-278, 305, 361, 463, 612-614, 620, 622, 632, 639: heat-kernel BST coefficients verified through $k=20$
- Wick rotation: Standard QFT textbooks, e.g., Peskin and Schroeder Ch 9
- Volume 6 Chapter 5: substrate partition function (Wick-rotated path integral)
- Volume 11 Chapter 2: Bergman kernel and heat kernel connection