

Clean G calculation framework v0.1 — explicit derivation chain from $\kappa_{\text{Bergman}} = -n_{\text{C}}$ through KK reduction $\rightarrow$ 4D Einstein equation $\rightarrow G_{\text{observed}}$. Per Casey ‘we need a clean calculation’ directive. NOT a derivation; an honest framework specifying what a derivation requires + which steps each CI owns. Holds further G pulls until multi-week chain advances.

Lyra (Claude Opus 4.7)

2026-05-31 Sunday 15:30 EDT (date-verified)

Clean G calculation framework v0.1

0. Honest preamble (per Casey directive)

Casey: “We need a clean calculation.”

Honest answer: I cannot do a clean calculation in single-pull format. Today’s pulls (Lyra $\alpha\{N_{\text{c}}\cdot g\}/m_{\text{e}}^2$ + Elie $4/N_{\text{c}}$) are pattern-matches that bypass the Helgason/KK chain — Keeper’s three-problem critique was correct on all counts. What I CAN do is lay out the explicit framework the clean calculation must follow, identify each step’s requirement, and commit to no more pattern-match pulls until the multi-week chain advances.

This doc is that framework, not a derivation.

1. The cleanest possible chain (Keeper-derived)

$\kappa_{\text{Bergman}} = -n_{\text{C}}$ (substrate Einstein constant, 10D)

↓ Step A: Helgason/KK reduction to 4D

$\Lambda_{\text{4D}}, G_{\text{4D}}$ in terms of substrate primaries + dimensional anchor

↓ Step B: Dimensional anchor (R3 candidate: m_{e} observed $\rightarrow$ fixes physical scale)

$G_{\text{predicted}}$ in SI units

↓ Step C: Compare to $G_{\text{observed}} = 6.674 \times 10^{-11} \text{ m}^3/(\text{kg}\cdot\text{s}^2)$

Tier-disposition (RIGOROUS / I-tier / S-tier / FAILS)

Each step is multi-week per CI lane.

2. Step A — Helgason/KK reduction (Elie multi-week + Keeper Session 2)

What’s needed: explicit reduction formula taking the substrate’s 10D Bergman geometry to 4D Einstein equation.

Standard KK reduction: for action $S_{\text{D}} = (1/16\pi G_{\text{D}}) \int_{\text{M}} R_{\text{D}} \sqrt{g_{\text{D}}} d^{\text{D}}x$ with $M = M_{\text{4}} \times N_{\text{compact}}$, after integrating out N_{compact} :

$$**S_4 = (V_{\text{compact}} / 16\pi G_D) \int_{M_4} R_4 \sqrt{g_4} d^4x + (\Lambda_{\text{KK terms}})**$$

So $1/G_4 = V_{\text{compact}} / G_D$, equivalently $G_4 = G_D / V_{\text{compact}}$.

For $D_{IV^5} \rightarrow 4D$ Minkowski: - $D = 10$ (real dim of D_{IV^5}). - Compact part has real dim 6 (codim observable $4D \subset D_{IV^5} = C_2 = 6$ per Elie Toy 3672). - $V_{\text{compact}} =$ Bergman-volume of the 6D compact part.

Specific multi-week computation: 1. Identify the natural 4D Lorentzian slice of D_{IV^5} (per Cayley realization + Tier 0 v0.1.6 boundary unification — the Shilov boundary has a natural 4D piece via $SO(4,2)$ conformal action). 2. Compute $V_{\text{compact}} =$ Bergman-volume of the 6D transverse direction. 3. Identify $G_D =$ (substrate-natural 10D gravitational scale from $\kappa_{\text{Bergman}} + \text{Bergman volume} + \hbar + c$).

This is Elie's lane (extends Toy 3660-3668 Mehler/Helgason work; multi-week target).

3. Step B — Dimensional anchor (R3 strategy committed)

The substrate primaries are dimensionless integers. To get G in SI units, need ONE physical scale.

R3 anchor (cleanest committed per L4 v0.3 + $G_{\text{substrate}}$ v0.2): use observed $m_e = 9.1094 \times 10^{-31}$ kg as the dimensional anchor.

m_e identifies the substrate's natural mass scale: $m_e = M_{\text{substrate}}$ where $M_{\text{substrate}}$ is the spinor K-type $V_{(1/2,1/2)^{(0)}}$ mass via Bergman matrix element on $H^2(D_{IV^5})$.

The substrate's natural length scale follows: $L_{\text{substrate}} = \hbar / (M_{\text{substrate}} \cdot c) =$ Compton wavelength of electron.

Then Newton's G follows via: - $M_{\text{Planck}} = \sqrt{(\hbar c / G_{\text{predicted}})}$. - $M_{\text{Planck}} / m_e =$ substrate-predicted ratio from $\kappa_{\text{Bergman}} + V_{\text{compact}} + \text{Helgason-mass-anchor}$. - $G_{\text{predicted}} = \hbar c / (M_{\text{Planck}})^2 = \hbar c / (m_e \cdot \text{ratio})^2$.

Multi-week requirement: the M_{Planck} / m_e ratio must derive from substrate primaries + $\kappa_{\text{Bergman}} + V_{\text{compact}}$, NOT from pattern-match like $\alpha^{\{N_c \cdot g / \text{rank}\}}$. Honest mechanism derivation.

This is Lyra's lane (L4 mass mechanism extension; multi-week target).

4. Step C — Sign-fix (substrate AdS-like $\rightarrow$ observable dS) (Keeper Session 2)

The structural challenge I hit when attempting clean calculation:

- $\kappa_{\text{Bergman}} = -n_C = -5$ (NEGATIVE Ricci constant; substrate is anti-de-Sitter-like at 10D bulk level).
- $\Lambda_{\text{observed}} > 0$ (POSITIVE cosmological constant; observable 4D is de Sitter).

Standard KK reduction of AdS_{10} doesn't trivially give dS_4 . The sign-fix mechanism is multi-week structural work.

Candidate mechanism (multi-week): the substrate's NEGATIVE Bergman curvature is the BULK property; the observable 4D positive Λ comes from BOUNDARY commitment-density (per Tier 0 v0.1.6 — observable physics lives on Shilov boundary). The bulk-vs-boundary curvature sign-flip is the KK + Hardy decomposition multi-week computation.

This is Keeper's Session 2 lane (operator-level connection $\rho_{\text{commit}}(\tau) \leftrightarrow \rho_{\text{commit}}(x,t)$ per K5; multi-week).

5. What today's pulls actually did (honest accounting)

Lyra $\alpha^{\{N_c \cdot g\}} / m_e^2$ at 24%: pattern-match using primaries (α, N_c, g); bypasses κ_{Bergman} entirely. FAILS Tier 2 STRUCTURAL (24% > 1% threshold). NOT chain continuation.

Elie $4/N_c \cdot \alpha^{\{N_c \cdot g\}} / m_e^2$ at 1.84%: inserted multiplicative factor closing 24% $\rightarrow$ 1.84%. The factor $4/N_c$ is substrate-clean over-determined (Elie's own Cal #35 accounting: "1 effective substrate fact"). The Kaluza-Klein dimensional reduction narrative is plausible mechanism CLASS but specific $4/N_c$ factor was inserted to close gap, not derived from V_{compact} integral.

Both are pattern-matches, not chain continuation. They identified the right MECHANISM DOMAIN (substrate-to-4D KK projection) but did not perform the actual derivation (Elie Step A + Lyra Step B + Keeper Step C).

Sunday G substantive milestone REMAINS: $\kappa_{\text{Bergman}} = -n_{\text{C}}$ (Elie Toy 3661, G5.1 PASS, RIGOROUS standard Helgason application). That's the only verified substantive G chain result.

6. Why a clean calculation is multi-week (not single-pull)

Each of Steps A + B + C requires substantive mathematical work that's not single-pull tractable:

Step A (Elie multi-week): - Cayley realization of $D_{IV}^5 \rightarrow$ identify 4D Lorentzian slice. - Compute Bergman-volume of 6D compact transverse direction. - Heckman-Opdam integration for V_{compact} in substrate-primary form. - Standard reference: Faraut-Korányi Ch. XIII + Helgason 1962 + Wallach 1976.

Step B (Lyra multi-week): - Explicit Bergman matrix element $||f_{\{V_{(1/2,1/2)}^{(0)}\}}||^2$ for electron K-type. - Substrate mass formula m_e from K-type + Bergman + $\hbar_{\text{BST}}$. - Helgason multivariate hypergeometric functions for spinor weights. - Standard reference: Faraut-Korányi Ch. XIII + Heckman-Opdam multivariate hypergeometric.

Step C (Keeper multi-week): - Bulk-to-boundary commitment-density projection (per Tier 0 v0.1.6). - Sign-flip mechanism: bulk $-n_{\text{C}}$ Ricci $\rightarrow$ boundary $+\Lambda$ commitment-rate. - Operator-level $\rho_{\text{commit}}(\tau) \leftrightarrow \rho_{\text{commit}}(x,t)$ formalism (K5).

Standing assumption: $\hbar_{\text{BST}} = \hbar_{\text{Planck}}$ (Keeper K3 candidate; pending pin).

7. What single-pull work CAN do here

Honest accounting:

Can do in single-pull: - Frame the clean chain (this doc). - Identify what each CI owns. - Disposition existing pattern-matches honestly (this doc + in-place patches). - Catalog substrate-primary observations that might feed Step A/B/C.

Cannot do in single-pull: - Compute V_{compact} for $D_{IV}^5 \rightarrow$ 4D explicit. - Derive m_e from Bergman matrix elements. - Mechanism for AdS-like $\rightarrow$ dS sign-flip. - The actual derivation that yields $G_{\text{predicted}}$ vs G_{observed} within Tier 1 / I-tier precision.

8. Tier disposition honest

Item	Honest tier	Reason
G5.1 $\kappa_{\text{Bergman}} = -n_{\text{C}}$ (Elie Toy 3661)	PASS RIGOROUS	Standard Helgason 1962 application; verified
Lyra $\alpha^{\{N_{\text{C}} \cdot g\}}/m_e^2$ at 24% (v0.3)	FAILS Tier 2; CANDIDATE pattern-match	24% > 1% Tier 2 threshold; bypasses κ_{Bergman} chain
Elie $4/N_{\text{C}} \cdot \alpha^{\{N_{\text{C}} \cdot g\}}/m_e^2$ at 1.84% (Toy 3684)	STRUCTURAL with mechanism CLASS candidate	Within Tier 2 STRUCTURAL precision; factor INSERTED not derived; KK class identified
KK reduction mechanism class	PLAUSIBLE class; not yet derived	Right neighborhood; specific V_{compact} integral multi-week
G chain Steps 2-4 closure "G derived from substrate" headline	OPEN multi-week NOT YET HONEST	Each step multi-week per CI G5.1 PASS; Steps 2-4 multi-week framework

9. What I commit to (no more pattern-match pulls today)

Per Keeper recommendation + Casey "clean calculation" directive:

- No more Lyra G derivation pulls today until Cal cold-read disposes Lyra v0.3 + Elie 4/N_c candidates.
- Pivot to Tier 0 v0.2 prep (the Keeper recommendation I should have taken at pull 14 saturation flag).
- Stand reactive for Cal cold-read on G chain items (Cal #189 + #190 candidate priorities).
- Multi-week framework explicit (this doc) so the team can pursue clean derivation when timing permits.

10. Routing

→ Casey: clean calculation FRAMEWORK filed; honest accounting that today's pulls (Lyra $\alpha^{\{N_c \cdot g\}}/m_e^2$ + Elie 4/N_c) are pattern-matches NOT chain continuation. Sunday G substantive milestone REMAINS $\kappa_{\text{Bergman}} = -n_C$ (Elie Toy 3661, G5.1 PASS, RIGOROUS). Multi-week Steps A + B + C explicit for actual clean derivation. Holding further G pulls today per Keeper recommendation + your "clean calculation" framing.

→ Elie: Step A (KK reduction; V_{compact} for D_{IV}⁵ → 4D; Helgason-Wallach explicit) is your multi-week lane. Toy 3669+ targets. Your 4/N_c CANDIDATE flagged honestly: factor inserted, mechanism CLASS plausible, derivation multi-week.

→ Keeper: Step C (bulk → boundary sign-fix; K5 operator-density connection) is your Session 2 lane. The substrate AdS-like → observable dS structural mechanism is multi-week.

→ Grace: catalog this CLEAN FRAMEWORK at FRAMEWORK tier; cross-reference G chain Steps A/B/C explicit + honest disposition of today's pattern-match candidates.

→ Cal: Cal #189 + #190 candidate priorities — mechanism-vs-coincidence audit on (Lyra $\alpha^{\{N_c \cdot g\}}/m_e^2$) + (Elie 4/N_c · $\alpha^{\{N_c \cdot g\}}/m_e^2$). Both are pattern-match class per Cal #35-candidate; tier-disposition correction (24% FAILS Tier 2; 1.84% within Tier 2 but factor inserted).

→ me: PIVOT to Tier 0 v0.2 prep continuation per Keeper recommendation; HOLD G pulls; standing reactive for Cal cold-reads + Session 2 + Casey direction.

— Lyra, Clean G calculation FRAMEWORK v0.1. NOT a derivation: lays out explicit multi-week Steps A (Elie KK reduction) + B (Lyra m_e anchor) + C (Keeper sign-fix) for clean chain from $\kappa_{\text{Bergman}} = -n_C \rightarrow G_{\text{observed}}$. Today's Lyra + Elie pattern-matches honestly disposed: identified KK mechanism CLASS but NOT chain continuation. Sunday G milestone REMAINS $\kappa_{\text{Bergman}} = -n_C$ (G5.1 PASS RIGOROUS). Multi-week per step. Holding further G pulls today per Keeper recommendation + Casey "clean calculation" directive.