

Vol 6 Chapter 6 — Classical Statistical Mechanics

Keeper (author pass — deep math/physics revision)

2026-05-24 Sunday

Chapter 6 — Classical Statistical Mechanics

Extended-reach chapter (May 2026), identification tier. Stamped 2026-09-13 (Keeper, K1895): the same five integers appear here as identifications — formulas that match, mechanisms not claimed — presented on a separate shelf from the derived core so that a reader can trust the core without accepting the reach. Not re-verified line by line since May; the presentation scan found no retired reading in this file except where a dated FIRED note says so. Where this chapter and the Spine (./Spine_DIV5_QM_GR_SM/) or the theorem registry disagree, they win.

Level 1 — one sentence

Classical statistical mechanics — Maxwell-Boltzmann velocity distribution, equipartition $\langle E_{\text{mode}} \rangle = (1/2)k_B T$, ergodic-hypothesis time-averages-equal-ensemble-averages — is the substrate's high-temperature large-K-type-count limit at Scale 2, where the quantum partition function (Chapter 5) collapses to a classical phase-space integral.

Level 2 — graduate-physicist precision

6.1 The classical limit of the partition function

For a single particle of mass m in 3D with Hamiltonian $\hat{H} = \hat{p}^2/2m + V(\hat{x})$, the partition function:

$$Z(\beta) = \int \frac{d^3x d^3p}{h^3} e^{-\beta H(x,p)}$$

(classical limit of $\text{Tr} e^{-\beta \hat{H}}$). The factor h^3 comes from the quantum-classical correspondence — Planck's constant doesn't fully disappear; it appears as the phase-space volume per state.

For N identical particles:

$$Z_N(\beta) = \frac{1}{N!} \int \prod_{i=1}^N \frac{d^3 x_i d^3 p_i}{h^3} e^{-\beta \sum_i H_i}$$

with $N!$ from indistinguishability (Gibbs paradox resolution, Chapter 3 Section 3.8).

6.2 Maxwell-Boltzmann velocity distribution

For an ideal gas (no interactions), the momentum integral factors. The probability distribution for finding a particle with momentum $\vec{p}$:

$$P(\vec{p}) = \left(\frac{1}{2\pi m k_B T} \right)^{3/2} e^{-p^2/(2m k_B T)}$$

In terms of velocity $\vec{v} = \vec{p}/m$:

$$P(\vec{v}) = \left(\frac{m}{2\pi k_B T} \right)^{3/2} e^{-mv^2/(2k_B T)}$$

Maxwell-Boltzmann distribution (Maxwell 1860, Boltzmann 1871). Speed distribution: $P(v)dv = 4\pi v^2 P(\vec{v})dv$, giving the characteristic skewed distribution.

Standard moments: - $\langle v \rangle = \sqrt{8k_B T/\pi m}$ (mean speed) - $\langle v^2 \rangle = 3k_B T/m$ (mean square speed) - $v_{\text{rms}} = \sqrt{3k_B T/m}$ (root-mean-square) - $v_{\text{mp}} = \sqrt{2k_B T/m}$ (most probable)

For nitrogen ($m \approx 28$ amu) at room temperature: $v_{\text{rms}} \approx 517$ m/s. Substrate reading: the substrate's K-type momentum spectrum at thermal coupling T has this characteristic distribution.

6.3 Equipartition theorem

For each quadratic degree of freedom in H (like $p^2/2m$ for kinetic, $kx^2/2$ for harmonic potential), the equilibrium contribution to average energy is exactly $(1/2)k_B T$:

$$\left\langle \frac{1}{2} m v_x^2 \right\rangle = \frac{1}{2} k_B T, \quad \left\langle \frac{1}{2} k x^2 \right\rangle = \frac{1}{2} k_B T$$

A 3D ideal gas has 3 translational quadratic dofs $\rightarrow \langle E \rangle = (3/2)k_B T$ per particle. A diatomic gas adds 2 rotational quadratic dofs $\rightarrow \langle E \rangle = (5/2)k_B T$. Plus 2 vibrational (kinetic + potential) at high $T \rightarrow (7/2)k_B T$.

This explains the heat capacity $C_V = (3/2)R$ for monatomic, $(5/2)R$ for diatomic at room temperature, $(7/2)R$ at high T where vibrational modes activate.

Substrate-mechanism reading: each “quadratic degree of freedom” corresponds to a substrate K-type direction with quadratic Casimir contribution; thermal averaging gives $(1/2)k_B T$ per direction.

6.4 Equipartition breakdown at low T

Why don’t vibrational modes contribute at room temperature for diatomic gases? Because their characteristic energy $\hbar\omega_{\text{vib}}$ is large compared to $k_B T$ — the modes are *frozen out*. Equipartition assumes the classical continuum; quantum-mechanically, modes with $\hbar\omega \gg k_B T$ are stuck in their ground state and don’t contribute to C_V .

Substrate reading: the substrate’s K-type spectrum is discrete (Volume 5 Chapter 4); equipartition holds only when the K-type spacing is much smaller than $k_B T$ (continuous limit). For high-energy modes, the substrate doesn’t have thermal access to excited K-types; the modes contribute zero.

This is the Einstein-Debye correction to classical equipartition — Chapter 7 of this volume develops the full quantum statistical mechanics treatment.

6.5 Ergodic hypothesis

The ergodic hypothesis: for a system in thermal equilibrium, time averages equal ensemble averages.

$$\bar{f}_{\text{time}} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(x(t), p(t)) dt = \int dx dp f(x, p) \rho_{\text{eq}}(x, p) = \langle f \rangle_{\text{ensemble}}$$

This is what allows replacing time-evolution simulations with phase-space-integral computations. Birkhoff (1931) and von Neumann (1932) gave conditions for ergodicity; for typical interacting systems, ergodicity holds though rigorous proofs are scarce.

Substrate-mechanism reading: ergodicity is the substrate’s accessibility of all energy-shell K-types over long times. The substrate’s many-tick evolution (Volume 0 Chapter 3 Scale 2) explores its accessible K-type configurations; sampling sufficient ticks recovers the equilibrium distribution.

6.6 Kinetic theory and transport

Maxwell-Boltzmann distribution enables classical kinetic theory:

- Mean free path: $\ell = 1/(\sqrt{2}n\sigma)$ where n is density, σ collision cross section
- Viscosity: $\eta = (1/3)n\langle v \rangle \ell$ (Maxwell 1860; surprisingly independent of n for dilute gases)
- Thermal conductivity: $\kappa = (1/3)n\langle v \rangle \ell c_v$ where c_v is heat capacity per particle

- Diffusion coefficient: $D = (1/3)\langle v \rangle \ell$

The Boltzmann transport equation governs nonequilibrium transport; classical statistical mechanics's apparatus for irreversible processes.

6.7 Worked example: pressure of ideal gas

Pressure on a wall from particles bombarding it. For particles of mass m moving with velocity $v_x > 0$ hitting wall: momentum transfer per collision = $2mv_x$ (elastic). Number of particles per unit area per second with velocity $v_x = (N/V)v_x P(v_x)$.

Total pressure:

$$P = \int_0^{\infty} 2mv_x \cdot \frac{N}{V} v_x P(v_x) dv_x = \frac{N}{V} m \langle v_x^2 \rangle = \frac{N}{V} k_B T$$

Recovers ideal gas law $PV = Nk_B T$. Substrate reading: the substrate's K-type momentum exchange with the wall's K-types gives the macroscopic pressure as the integrated K-type-momentum flux.

6.8 The H-theorem

Boltzmann's H-theorem: for a dilute gas, the function $H = \int f \ln f d^3v$ (where f is the velocity distribution) is monotonically non-increasing in time. As the gas approaches equilibrium, H decreases toward its minimum at the Maxwell-Boltzmann distribution; $H_{\text{equilibrium}} = -S/k_B$ recovers Boltzmann entropy.

Substrate-mechanism reading: the H-theorem is the substrate's many-tick Scale-2 entropy increase — exactly Zone 3 commitment information-discard summed over many cycles.

The Loschmidt paradox (time-reversal in microscopic dynamics vs irreversibility of H-theorem) was resolved at the substrate level: Zone 3 commitment is the substrate's intrinsic time-asymmetric step (Chapter 3 Section 3.5).

6.9 K-audit anchors and connections

- Casey's Principle: equipartition is substrate counting K-types per direction
- Maxwell 1860, Boltzmann 1871: foundational classical-stat-mech
- Volume 0 Chapter 3: Scale-2 averaging
- Chapter 5 of this volume: partition function (classical limit derivation)

Level 3 — 5th-grader accessibility

Air molecules in your room are moving fast — about 500 m/s on average. They don't all move the same speed; their speeds follow the Maxwell-Boltzmann distribution, a bell-curve-like shape that has more slow particles than fast ones. The

average kinetic energy per direction is exactly $(1/2)k_B T$ — that’s the equipartition theorem. So a 3D particle has $(3/2)k_B T$ of kinetic energy on average. From this you can derive the ideal gas law ($PV = Nk_B T$), viscosity, diffusion, heat conduction — all of classical kinetic theory. In BST, this is what happens at the substrate’s Scale 2 (many-Koons-tick averaging): at room temperature, billions of substrate K-type configurations are thermally accessible; the substrate’s count over those configurations gives the Maxwell-Boltzmann distribution. At low temperature, fewer K-types are accessible, and “equipartition breaks down” — vibrational modes freeze out because the substrate can’t thermally reach their excited K-types.

What comes next

Chapter 7 develops quantum statistical mechanics — Bose-Einstein and Fermi-Dirac distributions, Bose-Einstein condensation, Fermi gases, blackbody radiation, and the substrate-mechanism extensions.

Where to look this up

- Maxwell-Boltzmann: Maxwell 1860; Boltzmann 1871
- Equipartition: standard derivations; Pathria Ch 3
- H-theorem: Boltzmann 1872; Lebowitz 1993 review
- Kinetic theory: Chapman and Cowling, *The Mathematical Theory of Non-Uniform Gases*
- BST anchors: Casey’s Principle; Volume 0 Chapter 3 Scale 2
- Chapter 5 of this volume: partition function (master engine)