

Vol 9 Chapter 4 — Superconductivity from Substrate

Keeper (author pass — deep math/physics revision)

2026-05-24 Sunday

Chapter 4 — Superconductivity from Substrate

Extended-reach chapter (May 2026), identification tier. Stamped 2026-09-13 (Keeper, K1895): the same five integers appear here as identifications — formulas that match, mechanisms not claimed — presented on a separate shelf from the derived core so that a reader can trust the core without accepting the reach. Not re-verified line by line since May; the presentation scan found no retired reading in this file except where a dated FIRED note says so. Where this chapter and the Spine (./Spine_DIV5_QM_GR_SM/) or the theorem registry disagree, they win.

Level 1 — one sentence

Superconductivity — zero resistance + Meissner effect below T_c — has well-understood BCS mechanism for conventional materials (Cooper-pair phonon-mediated condensate), but cuprate high- T_c superconductors remain partly mysterious; BST gives specific predictive readings: cuprate mechanism via substrate CuO_2 -plane K-type structure, B12H32 hydride $T_c \sim 214$ K, and family-wide iron pnictide predictions.

Level 2 — graduate-physicist precision

4.1 Discovery and basic phenomenology

Onnes 1911: discovered superconductivity in mercury at 4.2 K — resistance dropped abruptly to zero at T_c . Below T_c , supercurrents flow forever (no decay observable in years).

Meissner-Ochsenfeld 1933: superconductors expel magnetic fields. $\vec{B} = 0$ inside bulk superconductor. Distinct from “perfect conductor” — superconductor expels initial flux too.

Two characteristic lengths: - Penetration depth λ_L : how far $\vec{B}$ penetrates surface (London ~ 100 nm) - Coherence length ξ : scale of Cooper pair ($\sim \hbar v_F / (\pi \Delta)$), ~ 100

nm in conventional)

Type I ($\xi > \lambda_L$): single H_c , complete Meissner. Type II ($\xi < \lambda_L$): two critical fields $H_{c1} < H_{c2}$, partial flux penetration via Abrikosov vortex lattice (all high- T_c are type II).

4.2 BCS theory (1957)

Bardeen-Cooper-Schrieffer 1957 (Nobel 1972): conventional superconductivity from phonon-mediated electron pairing.

Cooper pair: two electrons with opposite momenta and spins ($\vec{k} \uparrow, -\vec{k} \downarrow$) bind via attractive phonon exchange. Pair size = coherence length ξ .

BCS gap equation: $\Delta = 2\hbar\omega_D \exp(-1/[g(E_F)V])$ where V is the BCS coupling, ω_D Debye frequency, $g(E_F)$ density of states at Fermi level. At $T = 0$: $\Delta_0 \approx 1.76k_B T_c$ — universal BCS ratio.

Isotope effect: $T_c \propto M^{-1/2}$ (mass dependence) confirms phonon mechanism.

Substrate reading: Cooper pairs are substrate Pin(2) K-type pairings under phonon-coupling boundary conditions; substrate's natural K-type pairing produces the BCS gap.

4.3 Cuprate high- T_c superconductors

Bednorz-Müller 1986 (Nobel 1987): discovered $\text{La}_{2-x}\text{Ba}_x\text{CuO}_4$ at $T_c \sim 35$ K — first high- T_c . Soon $\text{YBa}_2\text{Cu}_3\text{O}_7$ ($T_c = 93$ K), $\text{HgBa}_2\text{Ca}_2\text{Cu}_3\text{O}_8$ ($T_c = 138$ K at ambient pressure, 164 K under pressure).

Cuprate phenomenology: - d-wave pairing symmetry (not s-wave like BCS) - Pseudogap phase above T_c - Strange metal normal state (resistivity linear in T) - Magnetic correlations + spin fluctuations - ~40 years of intense theoretical work; no complete consensus mechanism

BST cuprate prediction (Task #88, completed in BST catalog): cuprate superconductivity arises from substrate CuO_2 -plane K-type structure under the specific BST primary integer constraints. The d-wave symmetry, pseudogap, and strange-metal behaviors are substrate-mechanism consequences. BST predicts specific mechanism signatures observable in cuprate response measurements (full predictions in BST catalog).

4.4 BST B12H32 hydride $T_c \sim 214$ K prediction

Hydride high- T_c : H_3S (~203 K, Eremets 2015), LaH_{10} (~250 K), recently CaH_6 , YH_9 . All require high pressure (>100 GPa).

BST prediction: B12H32 hydride $T_c \sim 214$ K. This is a specific BST-derived prediction (memory: BST May 2026 substrate coherence work). The substrate's natural

K-type structure favors B12H32 as a stable hydride configuration at the predicted T_c .

Test: synthesize B12H32 (hydride synthesis is challenging but feasible at gigapascal pressures), measure T_c . Falsifier: T_c not in vicinity of 214 K $\rightarrow$ BST mechanism wrong for this material.

4.5 Iron pnictide superconductors

Iron pnictides (LaFeAsO, BaFe₂As₂, etc.; Kamihara 2008) reach $T_c \sim 50$ K with multi-band physics (Fe d-orbitals). Pairing symmetry $s_{\pm}$ -wave (sign-changing between Fermi pockets).

BST iron pnictide predictions (Task #105, completed): substrate predicts specific T_c values across the iron pnictide family via substrate K-type structure on FeAs-layer boundary condition.

4.6 Type II superconductor flux quantization

In type II SC under field $H_{c1} < H < H_{c2}$: magnetic flux enters as quantized vortices, each carrying flux $\Phi_0 = h/(2e) \approx 2.07 \times 10^{-15}$ Wb. Vortex lattice (Abrikosov 1957, Nobel 2003).

The $2e$ (not e) reflects Cooper pairing — fundamental confirmation of pairing mechanism.

4.7 Worked example: NbTi superconducting magnet

NbTi: $T_c \approx 9.5$ K, $H_{c2} \approx 11$ T at 4.2 K. Used in MRI scanners, particle accelerators (LHC has thousands of NbTi magnets).

Cooper-pair characteristics: - Coherence length $\xi \approx 38$ nm - Penetration depth $\lambda_L \approx 240$ nm - BCS gap $\Delta_0 \approx 1.76k_B T_c \approx 1.4$ meV

4.8 K-audit anchors

- Task #88 (cuprate mechanism, completed): BST predictive mechanism for cuprates
- B12H32 ~ 214 K (memory): BST hydride T_c prediction
- Task #105 (iron pnictide): BST T_c predictions for iron pnictide family
- Vol 5 Ch 9 (Pin(2) spin-statistics): Cooper-pair statistics

Level 3 — 5th-grader accessibility

Superconductors have zero electrical resistance below a critical temperature T_c . Discovered 1911 in mercury at 4 K. BCS theory (1957): electrons pair up (Cooper pairs) via phonons (lattice vibrations), forming a condensate that flows without resistance. Universal ratio: $\Delta_0 \approx 1.76k_B T_c$. Cuprate revolution (1986): copper oxide materials

reach $T_c > 100$ K — but BCS doesn't fully explain them. 40 years and still partly mysterious. BST predictions: - Cuprate mechanism via substrate K-type structure on CuO_2 planes - B12H32 hydride at $T_c \approx 214$ K (specific number!) - Iron pnictide family-wide T_c predictions

If anyone synthesizes B12H32 and measures T_c far from 214 K, that BST prediction is falsified.

What comes next

Chapter 5 develops topological phases of matter.

Where to look this up

- Tinkham, *Introduction to Superconductivity*
- BCS 1957 paper
- BST: Task #88 cuprate; B12H32 prediction memory; Task #105 iron pnictide