

Vol 10 Chapter 11 — Asymptotic Analysis and WKB

Keeper (author pass — deep math/physics revision)

2026-05-24 Sunday

Chapter 11 — Asymptotic Analysis and WKB

Extended-reach chapter (May 2026), identification tier. Stamped 2026-09-13 (Keeper, K1895): the same five integers appear here as identifications — formulas that match, mechanisms not claimed — presented on a separate shelf from the derived core so that a reader can trust the core without accepting the reach. Not re-verified line by line since May; the presentation scan found no retired reading in this file except where a dated FIRED note says so. Where this chapter and the Spine (./Spine_DIV5_QM_GR_SM/) or the theorem registry disagree, they win.

Level 1 — one sentence

Asymptotic analysis treats expansions valid in some limit (small parameter, large argument), with WKB as the canonical example (semiclassical expansion in powers of $\hbar$), saddle-point/Laplace methods for $\int e^{f(z)/\hbar} dz$, and stationary-phase for oscillatory integrals — all foundational for path integrals (Ch 10), turning points in QM, and asymptotic limits of special functions.

Level 2 — graduate-physicist precision

11.1 Asymptotic expansions

A formal series $\sum a_n/x^n$ is asymptotic to $f(x)$ as $x \rightarrow \infty$ if for each N :

$$\left| f(x) - \sum_{n=0}^N a_n/x^n \right| = o(1/x^N)$$

May not converge — but partial sums approximate well.

Stirling: $\ln \Gamma(s+1) \sim s \ln s - s + (1/2) \ln(2\pi s) + 1/(12s) - 1/(360s^3) + \dots$

11.2 Laplace's method (saddle-point)

For $I(\lambda) = \int_a^b e^{\lambda f(t)} g(t) dt$ with λ large and f having a unique maximum at $t_0 \in (a, b)$:

$$I(\lambda) \sim g(t_0) e^{\lambda f(t_0)} \sqrt{\frac{2\pi}{\lambda |f''(t_0)|}}$$

Generalizes to complex contour integrals (steepest descent).

11.3 Stationary phase

For oscillatory integrals $I(\lambda) = \int g(t) e^{i\lambda f(t)} dt$ with λ large:

$$I(\lambda) \sim g(t_0) e^{i\lambda f(t_0)} \sqrt{\frac{2\pi}{\lambda |f''(t_0)|}} e^{i\pi/4 \cdot \text{sgn}(f''(t_0))}$$

at stationary points $f'(t_0) = 0$. Foundation for path-integral saddle-point.

11.4 WKB approximation

For Schrödinger equation $\hbar^2 \psi'' = -[E - V(x)]\psi$: try $\psi = \exp(iS/\hbar)$, expand $S = S_0 + \hbar S_1 + \dots$. At leading order:

$$S_0(x) = \pm \int^x p(x') dx', \quad p(x) = \sqrt{2m[E - V(x)]}$$

The WKB wave function: $\psi_{\text{WKB}}(x) \sim p(x)^{-1/2} e^{\pm i \int p dx / \hbar}$.

Gives semiclassical quantization (Bohr-Sommerfeld): $\oint p dx = 2\pi\hbar(n + 1/2)$.

Connection formulas across turning points $V(x) = E$: Airy function asymptotics.

11.5 Substrate connection

WKB is the small- $\hbar$ limit of the path integral (Ch 10): saddle-point + Gaussian fluctuations. In BST: substrate's many-Koons-tick coherent sum concentrates on classical-stationary path as $\hbar$ -effective becomes small — Scale-2 emergence of classical mechanics.

11.6 K-audit anchors

- Vol 5 Ch 5: path integral semiclassical limit
- Vol 8 Ch 3: classical Lagrangian as WKB leading order

Level 3 — 5th-grader accessibility

Asymptotic analysis: how do functions behave in some limit (small parameter, large argument)? Laplace's method: integrals like $\int e^{\lambda f(t)} dt$ are dominated by the peak of f as $\lambda \rightarrow \infty$. Stationary phase: oscillatory integrals are dominated by where the phase is stationary. WKB: Schrödinger equation in semiclassical limit gives $\psi \sim p^{-1/2} e^{\pm i \int p dx / \hbar}$. Quantization: $\oint p dx = 2\pi\hbar(n + 1/2)$. These methods bridge quantum to classical and reveal universal asymptotic forms of special functions.

What comes next

Chapter 12 develops numerical methods.

Where to look this up

- Bender-Orszag, *Advanced Mathematical Methods*
- BST: Vol 5 Ch 5