

Higher gerbes, loop spaces, and transgression

Chris Kottke, with R. Melrose

New College of Florida

Geometric and Spectral Methods in PDE
CMO-BIRS Oaxaca, December 2016

Gerbes

Higher gerbes

Relation to loop spaces

Line bundles : $H^2(X; \mathbb{Z})$

- ▶ A complex line bundle $L \rightarrow X$ has a Chern class $c_1(L) \in H^2(X; \mathbb{Z})$.
- ▶ Naturality:

$$\begin{aligned}c_1(\mathbb{C} \times X) &= 0, & c_1(L \otimes L') &= c_1(L) + c_1(L'), \\c_1(L^{-1}) &= -c_1(L), & c_1(f^*L) &= f^*c_1(L)\end{aligned}$$

- ▶ $c_1(L) = c_1(L')$ if and only if $L \cong L'$.

Line bundles : $H^2(X; \mathbb{Z})$

- ▶ A complex line bundle $L \rightarrow X$ has a Chern class $c_1(L) \in H^2(X; \mathbb{Z})$.
- ▶ Naturality:

$$\begin{aligned} c_1(\mathbb{C} \times X) &= 0, & c_1(L \otimes L') &= c_1(L) + c_1(L'), \\ c_1(L^{-1}) &= -c_1(L), & c_1(f^*L) &= f^*c_1(L) \end{aligned}$$

- ▶ $c_1(L) = c_2(L')$ if and only if $L \cong L'$.
- ▶ Seen by Čech cohomology: $[L] \in \check{C}^1(X; \mathbb{C}^*)$ satisfies $d[L] = 1$, unique up to dh , $h \in \check{C}^0(X; \mathbb{C}^*)$, so

$$[L] \in \check{H}^1(X; \mathbb{C}^*) \cong H^2(X; \mathbb{Z}).$$

Gerbes : $H^3(X; \mathbb{Z})$

- Various versions: Giraud, Brylinski, Hitchin and Chattergee, Murray.

Gerbes : $H^3(X; \mathbb{Z})$

- ▶ Various versions: Giraud, Brylinski, Hitchin and Chatterjee, Murray.
- ▶ Murray: a *bundle gerbe* (L, Y, X) is
 - ▶ a fiber bundle (more generally: *locally split*, i.e., surjective map admitting local sections)

$$p : Y \longrightarrow X,$$

- ▶ a line bundle

$$L \longrightarrow Y^{[2]} = Y \times_X Y = \{(y_1, y_2) : p(y_1) = p(y_2) \in X\}$$

- ▶ with a product

$$\phi : L_{(y_1, y_2)} \otimes L_{(y_2, y_3)} \xrightarrow{\cong} L_{(y_1, y_3)}, \quad (y_1, y_2, y_3) \in Y^{[3]}$$

- ▶ satisfying associativity:

$$\phi \circ (1 \otimes \phi) = \phi \circ (\phi \otimes 1) : L_{(y_1, y_2)} \otimes L_{(y_2, y_3)} \otimes L_{(y_3, y_4)} \cong L_{(y_1, y_4)},$$

$$(y_1, y_2, y_3, y_4) \in Y^{[4]}$$

- ▶ (L, Y, X) has a *Dixmier Douady class* $DD(L, Y, X) \in H^3(X; \mathbb{Z})$.

Properties of gerbes

$$\begin{array}{ccccc} & & & & L \\ & & & & \downarrow \\ X & \xleftarrow{p} & Y & \xleftarrow[p_0]{p_1} & Y^{[2]} \end{array}$$

Properties of gerbes

$$\begin{array}{ccccc}
 & & Q & & L \\
 & & \downarrow & & \downarrow \\
 X & \xleftarrow{p} & Y & \xrightleftharpoons[p_0]{p_1} & Y^{[2]}
 \end{array}$$

- Trivialization: an isomorphism $L \cong \delta Q := p_0^* Q \otimes p_1^* Q^{-1}$ for some $Q \longrightarrow Y$.

Properties of gerbes

$$\begin{array}{ccccc}
 & & & & L \\
 & & & & \downarrow \\
 X & \xleftarrow{p} & Y & \xleftarrow[p_0]{p_1} & Y^{[2]}
 \end{array}$$

- ▶ Trivialization: an isomorphism $L \cong \delta Q := p_0^* Q \otimes p_1^* Q^{-1}$ for some $Q \longrightarrow Y$.
- ▶ Inverse: $(L, Y, X)^{-1} = (L^{-1}, Y, X)$.
- ▶ Product: $(L, Y, X) \otimes (L', Y', X) = (\pi_1^* L \otimes \pi_2^* L', Y \times_X Y', X)$
- ▶ Pullback: $f^*(L, Y, X) = (f^* L, f^* Y, X')$

Properties of gerbes

$$\begin{array}{ccccc}
 & & & & L \\
 & & & & \downarrow \\
 X & \xleftarrow{p} & Y & \xleftarrow[p_0]{p_1} & Y^{[2]}
 \end{array}$$

- ▶ Trivialization: an isomorphism $L \cong \delta Q := p_0^* Q \otimes p_1^* Q^{-1}$ for some $Q \longrightarrow Y$.
- ▶ Inverse: $(L, Y, X)^{-1} = (L^{-1}, Y, X)$.
- ▶ Product: $(L, Y, X) \otimes (L', Y', X) = (\pi_1^* L \otimes \pi_2^* L', Y \times_X Y', X)$
- ▶ Pullback: $f^*(L, Y, X) = (f^* L, f^* Y, X')$
- ▶ Relation with DD class:
 - ▶ $DD(L) = 0$ if and only if L is trivial.
 - ▶ $DD(L^{-1}) = -DD(L)$
 - ▶ $DD(L \otimes L') = DD(L) + DD(L')$
 - ▶ $DD(f^* L) = f^* DD(L)$.
 - ▶ $DD(L) = DD(L')$ if and only if L and L' are *stably isomorphic*, i.e., $L \otimes Q \cong L' \otimes Q'$ for trivial gerbes Q and Q' .

Example: lifting bundle gerbes

- ▶ $E \longrightarrow X$ principal G bundle, where G has a central extension

$$1 \longrightarrow \mathbb{C}^* \longrightarrow \widehat{G} \longrightarrow G \longrightarrow 1$$

- ▶ $\widehat{G} \longrightarrow G$ defines an associated line bundle $L = \widehat{G} \times_{\mathbb{C}^*} \mathbb{C} \longrightarrow G$
- ▶ Difference map $u : E^{[2]} \longrightarrow G$, where “ $u(y_0, y_1) = y_0^{-1} y_1$ ” i.e., $u(y_0, y_1) = g$ such that $y_1 = y_0 g$.
- ▶ $(u^* L, E, X)$ is the *lifting bundle gerbe* for E .

$$\begin{array}{ccc}
 u^* L & & L \\
 \downarrow & & \downarrow \\
 E^{[2]} & \xrightarrow{u} & G \\
 \downarrow & & \\
 X & &
 \end{array}$$

- ▶ $DD(u^* L, E, X) \in H^3(X; \mathbb{Z})$ is the obstruction to lifting E to a \widehat{G} bundle $\widehat{E} \longrightarrow X$.

Gerbes as simplicial line bundles

$$X \longleftarrow Y \rightrightarrows Y^{[2]} \Rrightarrow Y^{[3]} \Rrightarrow Y^{[4]} \dots$$

Gerbes as simplicial line bundles

$$X \longleftarrow Y \rightrightarrows Y^{[2]} \Rrightarrow Y^{[3]} \Rrightarrow Y^{[4]} \dots$$

- ▶ A *simplicial space* (over X) is a sequence $\{Y_n : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \longrightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \longrightarrow Y_n$, $j = 0, \dots, n-1$ (commuting with maps to X), satisfying relations derived from those of standard simplices.

Gerbes as simplicial line bundles

$$X \longleftarrow Y_0 \rightrightarrows Y_1 \rightrightarrows Y_2 \rightrightarrows Y_3 \quad \dots$$

- ▶ A *simplicial space* (over X) is a sequence $\{Y_n : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \longrightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \longrightarrow Y_n$, $j = 0, \dots, n-1$ (commuting with maps to X), satisfying relations derived from those of standard simplices.

Gerbes as simplicial line bundles

$$\begin{array}{ccccccc}
 & & L & & \delta L & & \delta^2 L \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y_0 & \rightrightarrows & Y_1 & \rightrightarrows & Y_2 & \rightrightarrows & Y_3 & \cdots
 \end{array}$$

- ▶ A *simplicial space* (over X) is a sequence $\{Y_n : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \longrightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \longrightarrow Y_n$, $j = 0, \dots, n-1$ (commuting with maps to X), satisfying relations derived from those of standard simplices.
- ▶ [Brylinski-McLaughlin]: A *simplicial line bundle* is a line bundle $L \longrightarrow Y_1$ with a trivialization of $\delta L = p_0^* L \otimes p_1^* L^{-1} \otimes p_2^* L$ pulling back to the canonical trivialization of $\delta \delta L$.

Gerbes as simplicial line bundles

$$\begin{array}{ccccccc}
 & & L & & \delta L & & \delta^2 L \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y_0 & \rightrightarrows & Y_1 & \rightrightarrows & Y_2 & \rightrightarrows & Y_3 & \cdots
 \end{array}$$

- ▶ A *simplicial space* (over X) is a sequence $\{Y_n : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \rightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \rightarrow Y_n$, $j = 0, \dots, n-1$ (commuting with maps to X), satisfying relations derived from those of standard simplices.
- ▶ [Brylinski-McLaughlin]: A *simplicial line bundle* is a line bundle $L \rightarrow Y_1$ with a trivialization of $\delta L = p_0^* L \otimes p_1^* L^{-1} \otimes p_2^* L$ pulling back to the canonical trivialization of $\delta \delta L$.
- ▶ In case Y_\bullet consists of fiber products $Y^{[\bullet-1]}$ of a locally split map $Y \rightarrow X$, this recovers the definition of a bundle gerbe.

Gerbes as simplicial line bundles

$$\begin{array}{ccccccc}
 & & L & & \delta L & & \delta^2 L \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\quad} & Y^{[2]} & \xleftarrow{\quad} & Y^{[3]} & \xleftarrow{\quad} & Y^{[4]} & \cdots
 \end{array}$$

- ▶ A *simplicial space* (over X) is a sequence $\{Y_n : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \longrightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \longrightarrow Y_n$, $j = 0, \dots, n-1$ (commuting with maps to X), satisfying relations derived from those of standard simplices.
- ▶ [Brylinski-McLaughlin]: A *simplicial line bundle* is a line bundle $L \longrightarrow Y_1$ with a trivialization of $\delta L = p_0^* L \otimes p_1^* L^{-1} \otimes p_2^* L$ pulling back to the canonical trivialization of $\delta \delta L$.
- ▶ In case Y_\bullet consists of fiber products $Y^{[\bullet-1]}$ of a locally split map $Y \longrightarrow X$, this recovers the definition of a bundle gerbe.

Dixmier-Douady class via Čech theory

Dixmier-Douady class via Čech theory

$Y \longrightarrow X$ locally split induces a double complex

$$\begin{array}{ccccccc}
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) & \xrightarrow{d} & \dots
 \end{array}$$

which is vertically *exact*.

Dixmier-Douady class via Čech theory

$Y \longrightarrow X$ locally split induces a double complex

$$\begin{array}{ccccccc}
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) & \xrightarrow{d} & \dots
 \end{array}$$

which is vertically *exact*. The Dixmier-Douady class is just a diagram chase:

$$\begin{array}{ccccc}
 & & 1 & & \\
 & & \uparrow & & \\
 & & [L] \rightarrow 1 & & \\
 & & \uparrow & & \uparrow \\
 & & \beta \rightarrow d\beta \rightarrow 1 & & \\
 & & & \uparrow & \uparrow \\
 & & & \alpha \rightarrow 1 &
 \end{array}$$

Dixmier-Douady class via Čech theory

$Y \longrightarrow X$ locally split induces a double complex

$$\begin{array}{ccccccc}
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) & \xrightarrow{d} & \dots
 \end{array}$$

which is vertically *exact*. The Dixmier-Douady class is just a diagram chase:

$$\begin{array}{ccccc}
 & & 1 & & \\
 & & \uparrow & & \\
 & & [L] \rightarrow 1 & & \\
 & & \uparrow & & \uparrow \\
 & & \beta \rightarrow d\beta \rightarrow 1 & & \\
 & & & \uparrow & \uparrow \\
 & & & \alpha \rightarrow 1 &
 \end{array}$$

Then $DD(L) = [\alpha] \in \check{H}^2(X; \mathbb{C}^*) \cong H^3(X; \mathbb{Z})$.

Dixmier-Douady class via Čech theory

$Y \longrightarrow X$ locally split induces a double complex

$$\begin{array}{ccccccc}
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) & \xrightarrow{d} & \dots
 \end{array}$$

which is vertically *exact*. The Dixmier-Douady class is just a diagram chase:

$$\begin{array}{ccccc}
 & & 1 & & \\
 & & \uparrow & & \\
 & & [L] \rightarrow 1 & & \\
 & & \uparrow & & \uparrow \\
 & & \beta \rightarrow d\beta \rightarrow 1 & & \\
 & & & \uparrow & \uparrow \\
 & & & \alpha \rightarrow 1 &
 \end{array}$$

Then $DD(L) = [\alpha] \in \check{H}^2(X; \mathbb{C}^*) \cong H^3(X; \mathbb{Z})$. Also Y supports a bundle gerbe with class $[\alpha] \in H^3(X; \mathbb{Z})$ iff $\delta[\alpha] = 0 \in H^3(Y; \mathbb{Z})$.

2-gerbes : $H^4(X; \mathbb{Z})$

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.

2-gerbes : $H^4(X; \mathbb{Z})$

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.
- ▶ A *bundle 2-gerbe* (L, Z, Y, X) is a “simplicial bundle gerbe”

$$\begin{array}{ccccccc}
 & & \mathbb{L} & & \delta\mathbb{L} & & \delta^2\mathbb{L} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\times 2} & Y^{[2]} & \xleftarrow{\times 3} & Y^{[3]} & \xleftarrow{\times 4} & Y^{[4]} & \xleftarrow{\times 5} & Y^{[5]}
 \end{array}$$

- ▶ A fiber bundle $Y \longrightarrow X$,
- ▶ A gerbe $\mathbb{L} = (L, Z, Y^{[2]})$,
- ▶ A trivialization of $\delta\mathbb{L} = p_0^*\mathbb{L} \otimes p_1^*\mathbb{L}^{-1} \otimes p_2^*\mathbb{L}$ over $Y^{[3]}$,

2-gerbes : $H^4(X; \mathbb{Z})$

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.
- ▶ A *bundle 2-gerbe* (L, Z, Y, X) is a “simplicial bundle gerbe”

$$\begin{array}{ccccccc}
 & & \mathbb{L} & & \delta\mathbb{L} & & \delta^2\mathbb{L} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\times 2} & Y^{[2]} & \xleftarrow{\times 3} & Y^{[3]} & \xleftarrow{\times 4} & Y^{[4]} & \xleftarrow{\times 5} & Y^{[5]}
 \end{array}$$

- ▶ A fiber bundle $Y \longrightarrow X$,
- ▶ A gerbe $\mathbb{L} = (L, Z, Y^{[2]})$,
- ▶ A trivialization of $\delta\mathbb{L} = p_0^*\mathbb{L} \otimes p_1^*\mathbb{L}^{-1} \otimes p_2^*\mathbb{L}$ over $Y^{[3]}$,
- ▶ A 2-morphism (did I mention gerbes have 2-morphisms?) relating the induced trivialization of $\delta^2\mathbb{L}$ to the canonical one,
- ▶ A coherency condition on pulled back 2-morphisms over $Y^{[5]}$.

2-gerbes : $H^4(X; \mathbb{Z})$

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.
- ▶ A *bundle 2-gerbe* (L, Z, Y, X) is a “simplicial bundle gerbe”

$$\begin{array}{ccccccc}
 & & \mathbb{L} & & \delta\mathbb{L} & & \delta^2\mathbb{L} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\times 2} & Y^{[2]} & \xleftarrow{\times 3} & Y^{[3]} & \xleftarrow{\times 4} & Y^{[4]} & \xleftarrow{\times 5} & Y^{[5]}
 \end{array}$$

- ▶ A fiber bundle $Y \longrightarrow X$,
- ▶ A gerbe $\mathbb{L} = (L, Z, Y^{[2]})$,
- ▶ A trivialization of $\delta\mathbb{L} = p_0^*\mathbb{L} \otimes p_1^*\mathbb{L}^{-1} \otimes p_2^*\mathbb{L}$ over $Y^{[3]}$,
- ▶ A 2-morphism (did I mention gerbes have 2-morphisms?) relating the induced trivialization of $\delta^2\mathbb{L}$ to the canonical one,
- ▶ A coherency condition on pulled back 2-morphisms over $Y^{[5]}$.
- ▶ (L, Z, Y, X) has a well-defined class $DD(L, Z, Y, X) \in H^4(X; \mathbb{Z})$.

2-gerbes : $H^4(X; \mathbb{Z})$

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.
- ▶ A *bundle 2-gerbe* (L, Z, Y, X) is a “simplicial bundle gerbe”

$$\begin{array}{ccccccc}
 & & \mathbb{L} & & \delta\mathbb{L} & & \delta^2\mathbb{L} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\times 2} & Y^{[2]} & \xleftarrow{\times 3} & Y^{[3]} & \xleftarrow{\times 4} & Y^{[4]} & \xleftarrow{\times 5} & Y^{[5]}
 \end{array}$$

- ▶ A fiber bundle $Y \longrightarrow X$,
- ▶ A gerbe $\mathbb{L} = (L, Z, Y^{[2]})$,
- ▶ A trivialization of $\delta\mathbb{L} = p_0^*\mathbb{L} \otimes p_1^*\mathbb{L}^{-1} \otimes p_2^*\mathbb{L}$ over $Y^{[3]}$,
- ▶ A 2-morphism (did I mention gerbes have 2-morphisms?) relating the induced trivialization of $\delta^2\mathbb{L}$ to the canonical one,
- ▶ A coherency condition on pulled back 2-morphisms over $Y^{[5]}$.
- ▶ (L, Z, Y, X) has a well-defined class $DD(L, Z, Y, X) \in H^4(X; \mathbb{Z})$.
- ▶ For higher gerbes $(H^{\geq 5}(X; \mathbb{Z}))$, higher and more complicated coherency conditions will appear.
- ▶ Y and Z are not on equal footing.

Bigerbes

A nicer version of 2-gerbes:

$$\begin{array}{ccc} Z & \longleftarrow & W_{00} \\ \downarrow & & \downarrow \\ X & \longleftarrow & Y \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ fiber bundles (or locally split).
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ fiber bundles forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger. Set $W_{00} = W$.

Bigerbes

A nicer version of 2-gerbes:

$$\begin{array}{ccc} Z & \longleftarrow & W_{00} \\ \downarrow & & \downarrow \\ X & \longleftarrow & Y \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ fiber bundles (or locally split).
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ fiber bundles forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger. Set $W_{00} = W$.
- ▶ Fill out the diagram by fiber products.

Bigerbes

A nicer version of 2-gerbes:

$$\begin{array}{ccccccc}
 & & Z^{[3]} & & & & \\
 & & \Downarrow & & & & \\
 & & Z^{[2]} & & & & \\
 & & \Downarrow & & & & \\
 & Z & \xleftarrow{\quad} & W_{00} & & & \\
 & \downarrow & & \downarrow & & & \\
 X & \xleftarrow{\quad} & Y & \xleftarrow{\quad} & Y^{[2]} & \xleftarrow{\quad} & Y^{[3]}
 \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ fiber bundles (or locally split).
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ fiber bundles forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger. Set $W_{00} = W$.
- ▶ Fill out the diagram by fiber products.

Bigerbes

A nicer version of 2-gerbes:

$$\begin{array}{ccccccc}
 Z^{[3]} & \longleftarrow & W_{02} & \rightrightarrows & W_{12} & \rightrightarrows & W_{22} \\
 \Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
 Z^{[2]} & \longleftarrow & W_{01} & \rightrightarrows & W_{11} & \rightrightarrows & W_{12} \\
 \Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
 Z & \longleftarrow & W_{00} & \rightrightarrows & W_{10} & \rightrightarrows & W_{20} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \rightrightarrows & Y^{[2]} & \rightrightarrows & Y^{[3]}
 \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ fiber bundles (or locally split).
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ fiber bundles forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger. Set $W_{00} = W$.
- ▶ Fill out the diagram by fiber products.


Bigerbes

A nicer version of 2-gerbes:

$$\begin{array}{ccccccc}
 Z^{[3]} & \longleftarrow & W_{02} & \rightrightarrows & W_{12} & \rightrightarrows & W_{22} \\
 \Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
 Z^{[2]} & \longleftarrow & W_{01} & \rightrightarrows & W_{11} & \rightrightarrows & W_{12} \\
 \Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
 Z & \longleftarrow & W_{00} & \rightrightarrows & W_{10} & \rightrightarrows & W_{20} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \rightrightarrows & Y^{[2]} & \rightrightarrows & Y^{[3]}
 \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ fiber bundles (or locally split).
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ fiber bundles forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger. Set $W_{00} = W$.
- ▶ Fill out the diagram by fiber products.
- ▶ $W_{\bullet\bullet}$ forms a *bisimplicial space* over X .


Bigerbes



Definition

A bundle bigerbe is a “bisimplicial line bundle” over $W_{\bullet\bullet}$, i.e., a line bundle L over W_{11} , with trivializations of $\delta_0 L$ and $\delta_1 L$, such that the induced trivializations of $\delta_0 \delta_1 L$ agree and which induce the canonical trivializations of $\delta_1^2 L$ and $\delta_0^2 L$.

Bigerbes



Definition

A bundle bigerbe is a “bisimplicial line bundle” over $W_{\bullet\bullet}$, i.e., a line bundle L over W_{11} , with trivializations of $\delta_0 L$ and $\delta_1 L$, such that the induced trivializations of $\delta_0 \delta_1 L$ agree and which induce the canonical trivializations of $\delta_1^2 L$ and $\delta_0^2 L$.

- ▶ Products, inverses, pull backs straightforward to define.
- ▶ A *trivialization* is an isomorphism $L \cong \delta_0 \delta_1 Q$ for a line bundle Q over W_{00} .

Theorem

A bundle bigerbe (L, W, X) has a well-defined Dixmier-Douady class $DD(L) \in H^4(X; \mathbb{Z})$, with

$$DD(L^{-1}) = -DD(L),$$

$$DD(L \otimes L') = DD(L) + DD(L'),$$

$$DD(f^*L) = f^*DD(L).$$

$DD(L) = 0$ if and only if L is trivial. $DD(L) = DD(L')$ if and only if L and L' are stably isomorphic.

Theorem

A bundle bigerbe (L, W, X) has a well-defined Dixmier-Douady class $DD(L) \in H^4(X; \mathbb{Z})$, with

$$DD(L^{-1}) = -DD(L),$$

$$DD(L \otimes L') = DD(L) + DD(L'),$$

$$DD(f^*L) = f^*DD(L).$$

$DD(L) = 0$ if and only if L is trivial. $DD(L) = DD(L')$ if and only if L and L' are stably isomorphic.

- Definition generalizes in a straightforward manner to higher degree (Exercise).

Theorem

A bundle multigerbe L of degree n has a well-defined Dixmier-Douady class $DD(L) \in H^{2+n}(X; \mathbb{Z})$, with

$$DD(L^{-1}) = -DD(L),$$

$$DD(L \otimes L') = DD(L) + DD(L'),$$

$$DD(f^*L) = f^*DD(L).$$

$DD(L) = 0$ if and only if L is trivial. $DD(L) = DD(L')$ if and only if L and L' are stably isomorphic.

- Definition generalizes in a straightforward manner to higher degree (Exercise).

Existence: based loop spaces

- ▶ Suppose X is connected, and take $Y = \mathcal{P}_*X$, the based path space.
- ▶ Then $Y^{[2]} = \mathcal{P}_*^{[2]}X \cong \Omega X$, the based loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe (L, \mathcal{P}_*X, X) , i.e., a simplicial line bundle L on ΩX .

Existence: based loop spaces

- ▶ Suppose X is connected, and take $Y = \mathcal{P}_*X$, the based path space.
- ▶ Then $Y^{[2]} = \mathcal{P}_*^{[2]}X \cong \Omega X$, the based loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe (L, \mathcal{P}_*X, X) , i.e., a simplicial line bundle L on ΩX .
- ▶ In this case, the gerbe product is known as the “fusion product” (Stolz-Teichner, Waldorf), and L is a “fusion line bundle”.

Existence: based loop spaces

- ▶ Suppose X is connected, and take $Y = \mathcal{P}_*X$, the based path space.
- ▶ Then $Y^{[2]} = \mathcal{P}_*^{[2]}X \cong \Omega X$, the based loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe (L, \mathcal{P}_*X, X) , i.e., a simplicial line bundle L on ΩX .
- ▶ In this case, the gerbe product is known as the “fusion product” (Stolz-Teichner, Waldorf), and L is a “fusion line bundle”.
- ▶ Likewise, if X is simply connected, with $Y = Z = \mathcal{P}_*X$ and $W_{00} = \mathcal{P}_*\mathcal{P}_*X$, Then $W_{11} = \Omega^2 X$, the double based loop space of X .
- ▶ Every class in $H^4(X; \mathbb{Z})$ is represented by a bisimplicial (or doubly fusion) line bundle $L \longrightarrow \Omega^2 X$.

Existence: based loop spaces

- ▶ Suppose X is connected, and take $Y = \mathcal{P}_*X$, the based path space.
- ▶ Then $Y^{[2]} = \mathcal{P}_*^{[2]}X \cong \Omega X$, the based loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe (L, \mathcal{P}_*X, X) , i.e., a simplicial line bundle L on ΩX .
- ▶ In this case, the gerbe product is known as the “fusion product” (Stolz-Teichner, Waldorf), and L is a “fusion line bundle”.
- ▶ Likewise, if X is simply connected, with $Y = Z = \mathcal{P}_*X$ and $W_{00} = \mathcal{P}_*\mathcal{P}_*X$, Then $W_{11} = \Omega^2 X$, the double based loop space of X .
- ▶ Every class in $H^4(X; \mathbb{Z})$ is represented by a bisimplicial (or doubly fusion) line bundle $L \rightarrow \Omega^2 X$.

Proposition

If X is k -connected, then every class in $H^{3+k}(X; \mathbb{Z})$ is represented by a multisimplicial line bundle on $\Omega^{2+k}X$.

Existence: free loop spaces

- ▶ Alternatively, take $Y = \mathcal{P}X$, the free path space, fibering over X^2 .
- ▶ Then $Y^{[2]} = \mathcal{P}^{[2]}X \cong \mathcal{L}X$, the free loop space.

Existence: free loop spaces

- ▶ Alternatively, take $Y = \mathcal{P}X$, the free path space, fibering over X^2 .
- ▶ Then $Y^{[2]} = \mathcal{P}^{[2]}X \cong \mathcal{L}X$, the free loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a simplicial line bundle $L = (L, \mathcal{P}X, X^2)$ on $\mathcal{L}X$, with the additional condition of a trivialization of the alternating product of pullbacks to the “figure-of-eight” loop space [K-Melrose, 2013].
- ▶ Figure-of-eight is yet another simplicial condition “over” the simplicial space

$$X \rightrightarrows X^2 \rightrightarrows X^3 \quad \dots$$

guaranteeing that the class in $H^3(X^2; \mathbb{Z})$ comes from $H^3(X; \mathbb{Z})$.

Existence: free loop spaces

- ▶ Alternatively, take $Y = \mathcal{P}X$, the free path space, fibering over X^2 .
- ▶ Then $Y^{[2]} = \mathcal{P}^{[2]}X \cong \mathcal{L}X$, the free loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a simplicial line bundle $L = (L, \mathcal{P}X, X^2)$ on $\mathcal{L}X$, with the additional condition of a trivialization of the alternating product of pullbacks to the “figure-of-eight” loop space [K-Melrose, 2013].
- ▶ Figure-of-eight is yet another simplicial condition “over” the simplicial space

$$X \rightrightarrows X^2 \rightrightarrows X^3 \quad \dots$$

guaranteeing that the class in $H^3(X^2; \mathbb{Z})$ comes from $H^3(X; \mathbb{Z})$.

Proposition

Every class in $H^{3+k}(X; \mathbb{Z})$ is represented by a multisimplicial (and multi figure-of-eight) line bundle on $\mathcal{L}^{2+k}X$.

Transgression and loop-fusion cohomology

Transgression and loop-fusion cohomology

- ▶ Take $\alpha \in H^3(X; \mathbb{Z})$ and $L \rightarrow \mathcal{L}X$ with $DD(L, \mathcal{P}X, X^2) = \alpha$.
- ▶ $c_1(L) \in H^2(\mathcal{L}X; \mathbb{Z})$ is the *transgression* of α :

$$\begin{array}{ccc}
 H^k(X; \mathbb{Z}) & \xrightarrow{\text{ev}^*} & H^k(\mathbb{S}^1 \times \mathcal{L}X; \mathbb{Z}) \\
 & \searrow T & \downarrow \int_{\mathbb{S}^1} \\
 & & H^{k-1}(\mathcal{L}X; \mathbb{Z})
 \end{array}$$

- ▶ Loses information since it forgets the simplicial properties of L .

Transgression and loop-fusion cohomology

- ▶ Take $\alpha \in H^3(X; \mathbb{Z})$ and $L \rightarrow \mathcal{L}X$ with $DD(L, \mathcal{P}X, X^2) = \alpha$.
- ▶ $c_1(L) \in H^2(\mathcal{L}X; \mathbb{Z})$ is the *transgression* of α :

$$\begin{array}{ccc}
 H^k(X; \mathbb{Z}) & \xrightarrow{\text{ev}^*} & H^k(\mathbb{S}^1 \times \mathcal{L}X; \mathbb{Z}) \\
 & \searrow T & \downarrow \int_{\mathbb{S}^1} \\
 & & H^{k-1}(\mathcal{L}X; \mathbb{Z})
 \end{array}$$

- ▶ Loses information since it forgets the simplicial properties of L .
- ▶ [K.-Melrose, 2015]: “Loop-fusion” Čech cohomology $H_{\text{lf}}^\bullet(\mathcal{L}X; \mathbb{Z})$ such that transgression factors through an isomorphism

$$H^k(X; \mathbb{Z}) \xrightarrow{\cong} H_{\text{lf}}^{k-1}(\mathcal{L}X; \mathbb{Z}) \longrightarrow H^{k-1}(\mathcal{L}X; \mathbb{Z}).$$

Transgression and loop-fusion cohomology

- ▶ Take $\alpha \in H^3(X; \mathbb{Z})$ and $L \rightarrow \mathcal{L}X$ with $DD(L, \mathcal{P}X, X^2) = \alpha$.
- ▶ $c_1(L) \in H^2(\mathcal{L}X; \mathbb{Z})$ is the *transgression* of α :

$$\begin{array}{ccc}
 H^k(X; \mathbb{Z}) & \xrightarrow{\text{ev}^*} & H^k(\mathbb{S}^1 \times \mathcal{L}X; \mathbb{Z}) \\
 & \searrow T & \downarrow \int_{\mathbb{S}^1} \\
 & & H^{k-1}(\mathcal{L}X; \mathbb{Z})
 \end{array}$$

- ▶ Loses information since it forgets the simplicial properties of L .
- ▶ [K.-Melrose, 2015]: “Loop-fusion” Čech cohomology $H_{\text{lf}}^\bullet(\mathcal{L}X; \mathbb{Z})$ such that transgression factors through an isomorphism

$$H^k(X; \mathbb{Z}) \xrightarrow{\cong} H_{\text{lf}}^{k-1}(\mathcal{L}X; \mathbb{Z}) \longrightarrow H^{k-1}(\mathcal{L}X; \mathbb{Z}).$$

Theorem

On $\mathcal{L}^\ell X$ there is a well-defined loop-fusion cohomology $\check{H}_{\text{lf}}^\bullet(\mathcal{L}^\ell X; \mathbb{Z})$ through which iterated transgression factors as an isomorphism:

$$H_{\text{lf}}^k(\mathcal{L}^\ell X; \mathbb{Z}) \xrightarrow{\cong} H_{\text{lf}}^{k-n}(\mathcal{L}^{\ell+n} X; \mathbb{Z}).$$

Spin structures on loop space

- ▶ Let X be a spin manifold and $E \rightarrow X$ the principal $G = \text{Spin}_n$ bundle.
- ▶ Then $\mathcal{L}E \rightarrow \mathcal{L}X$ is a $\mathcal{L}G$ bundle, and $\mathcal{L}G$ has a central extension

$$1 \rightarrow \text{U}(1) \rightarrow \widehat{\mathcal{L}G} \rightarrow \mathcal{L}G \rightarrow 1$$

Spin structures on loop space

- ▶ Let X be a spin manifold and $E \rightarrow X$ the principal $G = \mathrm{Spin}_n$ bundle.
- ▶ Then $\mathcal{L}E \rightarrow \mathcal{L}X$ is a $\mathcal{L}G$ bundle, and $\mathcal{L}G$ has a central extension

$$1 \rightarrow \mathrm{U}(1) \rightarrow \widehat{\mathcal{L}G} \rightarrow \mathcal{L}G \rightarrow 1$$

- ▶ A lift of $\mathcal{L}E$ to a $\widehat{\mathcal{L}G}$ -bundle is a “spin structure” on loop space [Atiyah, Witten].

Spin structures on loop space

- ▶ Let X be a spin manifold and $E \rightarrow X$ the principal $G = \text{Spin}_n$ bundle.
- ▶ Then $\mathcal{L}E \rightarrow \mathcal{L}X$ is a $\mathcal{L}G$ bundle, and $\mathcal{L}G$ has a central extension

$$1 \rightarrow \text{U}(1) \rightarrow \widehat{\mathcal{L}G} \rightarrow \mathcal{L}G \rightarrow 1$$

- ▶ A lift of $\mathcal{L}E$ to a $\widehat{\mathcal{L}G}$ -bundle is a “spin structure” on loop space [Atiyah, Witten].

Proposition

The lifting bundle gerbe $(u^\widehat{\mathcal{L}G}, \mathcal{L}E, \mathcal{L}X)$ is a bundle bigerbe associated to the bisimplicial space generated by*

$$\begin{array}{ccc} E^2 & \longleftarrow & \mathcal{P}E \\ \downarrow & & \downarrow \\ X^2 & \longleftarrow & \mathcal{P}X \end{array}$$

with Dixmier-Douady class $\frac{1}{2}p_1(X) \in H^4(X; \mathbb{Z})$.

- ▶ c.f. McLaughlin, Redden, Waldorf, K.-Melrose.

Questions and future directions

- ▶ Connection structures, representations of differential cohomology.
- ▶ On $\mathcal{L}X$ (and generally $\mathcal{L}^k X$), equivariance of L with respect to action of $\text{Diffeo}^+(\mathbb{S}^1)$ (and its central extension) [c.f. Brylinski]. This may have an important role to play in elliptic cohomology theories.
- ▶ Loop-fusion K-theory of $\mathcal{L}X$ and $\mathcal{L}^\ell X$.