

Multigerbes: a new theory of higher gerbes

Chris Kottke

Joint work in progress with R. Melrose

New College of Florida

Workshop on Geometric Quantization
BIRS, April 2018

- ▶ A complex line bundle $L \rightarrow X$ has a Chern class $c_1(L) \in H^2(X; \mathbb{Z})$.
- ▶ Naturality:

$$\begin{aligned}c_1(\underline{\mathbb{C}}) &= 0, & c_1(L \otimes L') &= c_1(L) + c_1(L'), \\c_1(L^{-1}) &= -c_1(L), & c_1(f^*L) &= f^*c_1(L)\end{aligned}$$

- ▶ $c_1(L) = c_2(L')$ if and only if $L \cong L'$.

- ▶ A complex line bundle $L \rightarrow X$ has a Chern class $c_1(L) \in H^2(X; \mathbb{Z})$.
- ▶ Naturality:

$$\begin{aligned} c_1(\underline{\mathbb{C}}) &= 0, & c_1(L \otimes L') &= c_1(L) + c_1(L'), \\ c_1(L^{-1}) &= -c_1(L), & c_1(f^*L) &= f^*c_1(L) \end{aligned}$$

- ▶ $c_1(L) = c_2(L')$ if and only if $L \cong L'$.
- ▶ Explicit in Čech cohomology: $[L] \in \check{C}^1(X; \mathbb{C}^*)$ satisfies $d[L] = 0$, unique up to dh , $h \in \check{C}^0(X; \mathbb{C}^*)$, so

$$[L] \in \check{H}^1(X; \mathbb{C}^*) \cong H^2(X; \mathbb{Z}).$$

- Various versions: Giraud, Brylinski, Hitchin and Chatterjee, Murray.

- ▶ Various versions: Giraud, Brylinski, Hitchin and Chatterjee, Murray.
- ▶ Murray: a *bundle gerbe* (L, Y, X) is
 - ▶ a *locally split* map (meaning surjective with local sections)

$$p : Y \longrightarrow X,$$

- ▶ a line bundle

$$L \longrightarrow Y^{[2]} = Y \times_X Y = \{(y_1, y_2) : p(y_1) = p(y_2) \in X\}$$

- ▶ with a product

$$\phi : L_{(y_1, y_2)} \otimes L_{(y_2, y_3)} \xrightarrow{\cong} L_{(y_1, y_3)}, \quad (y_1, y_2, y_3) \in Y^{[3]}$$

- ▶ satisfying associativity:

$$\begin{aligned} \phi \circ (1 \otimes \phi) &= \phi \circ (\phi \otimes 1) : L_{(y_1, y_2)} \otimes L_{(y_2, y_3)} \otimes L_{(y_3, y_4)} \cong L_{(y_1, y_4)}, \\ (y_1, y_2, y_3, y_4) &\in Y^{[4]} \end{aligned}$$

- ▶ (L, Y, X) has a *Dixmier Douady class* $DD(L, Y, X) \in H^3(X; \mathbb{Z})$.

Properties of gerbes

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccc} & & & & L \\ & & & & \downarrow \\ X & \xleftarrow{p} & Y & \xrightleftharpoons[p_0]{p_1} & Y^{[2]} \end{array}$$

Properties of gerbes

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccc} & & Q & & L \\ & & \downarrow & & \downarrow \\ X & \xleftarrow{p} & Y & \xrightleftharpoons[p_0]{p_1} & Y^{[2]} \end{array}$$

- Trivialization: an isomorphism $L \cong \delta Q := p_0^* Q \otimes p_1^* Q^{-1}$ for some line bundle $Q \rightarrow Y$.

$$\begin{array}{ccccc}
 & & & & L \\
 & & & & \downarrow \\
 X & \xleftarrow{p} & Y & \xleftarrow[p_0]{p_1} & Y^{[2]}
 \end{array}$$

- ▶ Trivialization: an isomorphism $L \cong \delta Q := p_0^* Q \otimes p_1^* Q^{-1}$ for some line bundle $Q \rightarrow Y$.
- ▶ Inverse: $(L, Y, X)^{-1} = (L^{-1}, Y, X)$.
- ▶ Product: $(L, Y, X) \otimes (L', Y', X) = (\pi_1^* L \otimes \pi_2^* L', Y \times_X Y', X)$
- ▶ Pullback: $f^*(L, Y, X) = (f^* L, f^* Y, X')$

$$\begin{array}{ccccc}
 & & & & L \\
 & & & & \downarrow \\
 X & \xleftarrow{p} & Y & \xleftarrow[p_0]{p_1} & Y^{[2]}
 \end{array}$$

- ▶ Trivialization: an isomorphism $L \cong \delta Q := p_0^* Q \otimes p_1^* Q^{-1}$ for some line bundle $Q \rightarrow Y$.
- ▶ Inverse: $(L, Y, X)^{-1} = (L^{-1}, Y, X)$.
- ▶ Product: $(L, Y, X) \otimes (L', Y', X) = (\pi_1^* L \otimes \pi_2^* L', Y \times_X Y', X)$
- ▶ Pullback: $f^*(L, Y, X) = (f^* L, f^* Y, X')$
- ▶ Relation with DD class:
 - ▶ $DD(L) = 0$ if and only if L is trivial.
 - ▶ $DD(L^{-1}) = -DD(L)$
 - ▶ $DD(L \otimes L') = DD(L) + DD(L')$
 - ▶ $DD(f^* L) = f^* DD(L)$.
 - ▶ $DD(L) = DD(L')$ if and only if L and L' are *stably isomorphic*, i.e., $L \otimes Q \cong L' \otimes Q'$ for trivial gerbes Q and Q' .
 - ▶ Better: $DD(L) = DD(L')$ if and only if there is a 1-isomorphism (a la Waldorf) $(L, Y, X) \rightarrow (L', Y', X)$.

- $E \longrightarrow X$ principal G bundle, where G has a central extension

$$1 \longrightarrow \mathrm{U}(1) \longrightarrow \widehat{G} \longrightarrow G \longrightarrow 1$$

- $\widehat{G} \longrightarrow G$ defines an associated line bundle $L = \widehat{G} \times_{\mathrm{U}(1)} \mathbb{C} \longrightarrow G$
- Difference map $u : E^{[2]} \longrightarrow G$, where $u(y_0, y_1) = g$ such that $y_1 = y_0 g$.
- (u^*L, E, X) is the *lifting bundle gerbe* for E .

$$\begin{array}{ccc} u^*L & & L \\ \downarrow & & \downarrow \\ E^{[2]} & \xrightarrow{u} & G \\ \downarrow & & \\ X & & \end{array}$$

- $DD(u^*L, E, X) \in H^3(X; \mathbb{Z})$ is the obstruction to lifting E to a \widehat{G} bundle $\widehat{E} \longrightarrow X$.

$$X \longleftarrow Y \rightrightarrows Y^{[2]} \Rrightarrow Y^{[3]} \Rrightarrow Y^{[4]} \cdots$$

- These higher fiber products define a *simplicial space* over X , i.e., a sequence $\{Y_n = Y^{[n+1]} : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \longrightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \longrightarrow Y_n$, $j = 0, \dots, n-1$ (all commuting with maps to X), satisfying the relations of standard simplices.

$$X \longleftarrow Y \rightrightarrows Y^{[2]} \Rrightarrow Y^{[3]} \Rrightarrow Y^{[4]} \cdots$$

- These higher fiber products define a *simplicial space* over X , i.e., a sequence $\{Y_n = Y^{[n+1]} : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \longrightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \longrightarrow Y_n$, $j = 0, \dots, n-1$ (all commuting with maps to X), satisfying the relations of standard simplices.

$$X \longleftarrow Y_0 \rightrightarrows Y_1 \rightrightarrows Y_2 \rightrightarrows Y_3 \cdots$$

- These higher fiber products define a *simplicial space* over X , i.e., a sequence $\{Y_n = Y^{[n+1]} : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \longrightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \longrightarrow Y_n$, $j = 0, \dots, n-1$ (all commuting with maps to X), satisfying the relations of standard simplices.

$$\begin{array}{ccccccc}
 & & L & & \delta L & & \delta^2 L \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y_0 & \rightrightarrows & Y_1 & \rightrightarrows & Y_2 & \rightrightarrows & Y_3 & \cdots
 \end{array}$$

- These higher fiber products define a *simplicial space* over X , i.e., a sequence $\{Y_n = Y^{[n+1]} : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \rightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \rightarrow Y_n$, $j = 0, \dots, n-1$ (all commuting with maps to X), satisfying the relations of standard simplices.
- [Brylinski-McLaughlin]: A *simplicial line bundle* is a line bundle $L \rightarrow Y_1$ with a trivialization of $\delta L = p_0^* L \otimes p_1^* L^{-1} \otimes p_2^* L$ pulling back to the canonical trivialization of $\delta^2 L$.

$$\begin{array}{ccccccc}
 & & L & & \delta L & & \delta^2 L \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y_0 & \rightrightarrows & Y_1 & \rightrightarrows & Y_2 & \rightrightarrows & Y_3 & \cdots
 \end{array}$$

- These higher fiber products define a *simplicial space* over X , i.e., a sequence $\{Y_n = Y^{[n+1]} : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \longrightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \longrightarrow Y_n$, $j = 0, \dots, n-1$ (all commuting with maps to X), satisfying the relations of standard simplices.
- [Brylinski-McLaughlin]: A *simplicial line bundle* is a line bundle $L \longrightarrow Y_1$ with a trivialization of $\delta L = p_0^* L \otimes p_1^* L^{-1} \otimes p_2^* L$ pulling back to the canonical trivialization of $\delta^2 L$.
- In case Y_\bullet consists of fiber products $Y^{[\bullet-1]}$ of a locally split map $Y \longrightarrow X$, this precisely recovers the definition of a bundle gerbe.

Gerbes as simplicial line bundles

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccccc}
 & & L & & \delta L & & \delta^2 L \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \rightrightarrows & Y^{[2]} & \rightrightarrows & Y^{[3]} & \rightrightarrows & Y^{[4]} & \cdots
 \end{array}$$

- ▶ These higher fiber products define a *simplicial space* over X , i.e., a sequence $\{Y_n = Y^{[n+1]} : n \in \mathbb{N}_0\}$ of spaces with *face maps* $p_j : Y_n \rightarrow Y_{n-1}$, $j = 0, \dots, n$ and *degeneracy maps* $s_j : Y_{n-1} \rightarrow Y_n$, $j = 0, \dots, n-1$ (all commuting with maps to X), satisfying the relations of standard simplices.
- ▶ [Brylinski-McLaughlin]: A *simplicial line bundle* is a line bundle $L \rightarrow Y_1$ with a trivialization of $\delta L = p_0^* L \otimes p_1^* L^{-1} \otimes p_2^* L$ pulling back to the canonical trivialization of $\delta^2 L$.
- ▶ In case Y_\bullet consists of fiber products $Y^{[\bullet-1]}$ of a locally split map $Y \rightarrow X$, this precisely recovers the definition of a bundle gerbe.

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccccc} & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\ \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \dots \\ & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\ \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) & \xrightarrow{d} & \dots \\ & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\ \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) & \xrightarrow{d} & \dots \end{array}$$

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccc} \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\ \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) \xrightarrow{d} \dots \\ \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\ \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) \xrightarrow{d} \dots \\ \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\ \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) \xrightarrow{d} \dots \end{array}$$

This double complex is vertically *exact* for $Y \rightarrow X$ locally split. In particular the total cohomology is isomorphic to $\check{H}^\bullet(X; \mathbb{C}^*)$.

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccc}
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) \xrightarrow{d} \dots \\
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) \xrightarrow{d} \dots \\
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) \xrightarrow{d} \dots
 \end{array}$$

This double complex is vertically *exact* for $Y \rightarrow X$ locally split. In particular the total cohomology is isomorphic to $\check{H}^\bullet(X; \mathbb{C}^*)$.

- The Čech chains are with respect to pairs of “admissible covers” of (X, Y) to which $p : Y \rightarrow X$ and its local sections are adapted, including higher intersections.

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccc}
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) \xrightarrow{d} \dots \\
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) \xrightarrow{d} \dots \\
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) \xrightarrow{d} \dots
 \end{array}$$

This double complex is vertically *exact* for $Y \rightarrow X$ locally split. In particular the total cohomology is isomorphic to $\check{H}^\bullet(X; \mathbb{C}^*)$.

- ▶ The Čech chains are with respect to pairs of “admissible covers” of (X, Y) to which $p : Y \rightarrow X$ and its local sections are adapted, including higher intersections.
- ▶ Fiber products of the pair of covers give covers of $Y^{[k]}$ for $k > 1$.

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccc}
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) \xrightarrow{d} \dots \\
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) \xrightarrow{d} \dots \\
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) \xrightarrow{d} \dots
 \end{array}$$

This double complex is vertically *exact* for $Y \rightarrow X$ locally split. In particular the total cohomology is isomorphic to $\check{H}^\bullet(X; \mathbb{C}^*)$.

- ▶ The Čech chains are with respect to pairs of “admissible covers” of (X, Y) to which $p : Y \rightarrow X$ and its local sections are adapted, including higher intersections.
- ▶ Fiber products of the pair of covers give covers of $Y^{[k]}$ for $k > 1$.
- ▶ The local sections of p induce chain homotopy contractions of each vertical complex.

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccc}
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) \xrightarrow{d} \dots \\
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) \xrightarrow{d} \dots \\
 \delta \uparrow & & \delta \uparrow & & \delta \uparrow \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) \xrightarrow{d} \dots
 \end{array}$$

This double complex is vertically *exact* for $Y \rightarrow X$ locally split. In particular the total cohomology is isomorphic to $\check{H}^\bullet(X; \mathbb{C}^*)$.

- ▶ The Čech chains are with respect to pairs of “admissible covers” of (X, Y) to which $p : Y \rightarrow X$ and its local sections are adapted, including higher intersections.
- ▶ Fiber products of the pair of covers give covers of $Y^{[k]}$ for $k > 1$.
- ▶ The local sections of p induce chain homotopy contractions of each vertical complex.
- ▶ Take the direct limit over all admissible pairs of covers. Over X and Y this is equivalent to the direct limit over all covers.

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccccc}
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) & \xrightarrow{d} & \dots
 \end{array}$$

The Dixmier-Douady class is the image in $\check{H}^2(X; \mathbb{C}^*)$ of the pure cocycle $-[L] \in \check{C}^1(Y^{[2]}; \mathbb{C}^*) \subset \check{C}^\bullet(Y^{[\bullet]}; \mathbb{C}^*)$ in $\check{H}^2(X; \mathbb{C}^*)$.

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccccc}
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) & \xrightarrow{d} & \dots
 \end{array}$$

The Dixmier-Douady class is the image in $\check{H}^2(X; \mathbb{C}^*)$ of the pure cocycle $-[L] \in \check{C}^1(Y^{[2]}; \mathbb{C}^*) \subset \check{C}^\bullet(Y^{[\bullet]}; \mathbb{C}^*)$ in $\check{H}^2(X; \mathbb{C}^*)$.

$$\begin{array}{ccccc}
 0 & & & & \\
 \uparrow & & & & \\
 -[L] & \rightarrow & 0 & & \\
 \uparrow & & \uparrow & & \\
 \beta & \rightarrow & d\beta & \rightarrow & 0 \\
 & & \uparrow & & \uparrow \\
 & & \alpha & \rightarrow & 0
 \end{array}$$

so $DD(L) = [\alpha] \in \check{H}^2(X; \mathbb{C}^*) \cong H^3(X; \mathbb{Z})$.

Dixmier-Douady class via Čech-simplicial double complex

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

$$\begin{array}{ccccccc}
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y^{[2]}; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(Y; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(Y; \mathbb{C}^*) & \xrightarrow{d} & \dots \\
 & \delta \uparrow & & \delta \uparrow & & \delta \uparrow & \\
 \check{C}^0(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^1(X; \mathbb{C}^*) & \xrightarrow{d} & \check{C}^2(X; \mathbb{C}^*) & \xrightarrow{d} & \dots
 \end{array}$$

The Dixmier-Douady class is the image in $\check{H}^2(X; \mathbb{C}^*)$ of the pure cocycle $-[L] \in \check{C}^1(Y^{[2]}; \mathbb{C}^*) \subset \check{C}^\bullet(Y^{[\bullet]}; \mathbb{C}^*)$ in $\check{H}^2(X; \mathbb{C}^*)$.

$$\begin{array}{ccccc}
 0 & & & & \\
 \uparrow & & & & \\
 -[L] & \rightarrow & 0 & & \\
 \uparrow & & \uparrow & & \\
 \beta & \rightarrow & d\beta & \rightarrow & 0 \\
 & & \uparrow & & \uparrow \\
 & & \alpha & \rightarrow & 0
 \end{array}$$

so $DD(L) = [\alpha] \in \check{H}^2(X; \mathbb{C}^*) \cong H^3(X; \mathbb{Z})$. It also follows that Y supports a bundle gerbe with class $[\alpha] \in H^3(X; \mathbb{Z})$ iff $p^*[\alpha] = 0 \in H^3(Y; \mathbb{Z})$.

- Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.
- ▶ A *bundle 2-gerbe* (L, Z, Y, X) is a “simplicial bundle gerbe”

$$\begin{array}{ccccccc}
 & & \mathbb{L} & & \delta\mathbb{L} & & \delta^2\mathbb{L} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\times 2} & Y^{[2]} & \xleftarrow{\times 3} & Y^{[3]} & \xleftarrow{\times 4} & Y^{[4]} & \xleftarrow{\times 5} & Y^{[5]}
 \end{array}$$

- ▶ A locally split map $Y \longrightarrow X$,
- ▶ A gerbe $\mathbb{L} = (L, Z, Y^{[2]})$,
- ▶ A trivialization of $\delta\mathbb{L} = p_0^*\mathbb{L} \otimes p_1^*\mathbb{L}^{-1} \otimes p_2^*\mathbb{L}$ over $Y^{[3]}$,

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.
- ▶ A *bundle 2-gerbe* (L, Z, Y, X) is a “simplicial bundle gerbe”

$$\begin{array}{ccccccc}
 & & \mathbb{L} & & \delta\mathbb{L} & & \delta^2\mathbb{L} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\times 2} & Y^{[2]} & \xleftarrow{\times 3} & Y^{[3]} & \xleftarrow{\times 4} & Y^{[4]} & \xleftarrow{\times 5} & Y^{[5]}
 \end{array}$$

- ▶ A locally split map $Y \longrightarrow X$,
- ▶ A gerbe $\mathbb{L} = (L, Z, Y^{[2]})$,
- ▶ A trivialization of $\delta\mathbb{L} = p_0^*\mathbb{L} \otimes p_1^*\mathbb{L}^{-1} \otimes p_2^*\mathbb{L}$ over $Y^{[3]}$,
- ▶ A 2-morphism (did I mention gerbes have 2-morphisms?) relating the induced trivialization of $\delta^2\mathbb{L}$ to the canonical one,
- ▶ A coherency condition on pulled back 2-morphisms over $Y^{[5]}$.

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.
- ▶ A *bundle 2-gerbe* (L, Z, Y, X) is a “simplicial bundle gerbe”

$$\begin{array}{ccccccc}
 & & \mathbb{L} & & \delta\mathbb{L} & & \delta^2\mathbb{L} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\times 2} & Y^{[2]} & \xleftarrow{\times 3} & Y^{[3]} & \xleftarrow{\times 4} & Y^{[4]} & \xleftarrow{\times 5} & Y^{[5]}
 \end{array}$$

- ▶ A locally split map $Y \longrightarrow X$,
- ▶ A gerbe $\mathbb{L} = (L, Z, Y^{[2]})$,
- ▶ A trivialization of $\delta\mathbb{L} = p_0^*\mathbb{L} \otimes p_1^*\mathbb{L}^{-1} \otimes p_2^*\mathbb{L}$ over $Y^{[3]}$,
- ▶ A 2-morphism (did I mention gerbes have 2-morphisms?) relating the induced trivialization of $\delta^2\mathbb{L}$ to the canonical one,
- ▶ A coherency condition on pulled back 2-morphisms over $Y^{[5]}$.
- ▶ (L, Z, Y, X) has a well-defined characteristic class $C(L, Z, Y, X) \in H^4(X; \mathbb{Z})$.

- ▶ Stevenson: gerbes have pullbacks, trivializations, morphisms, so we can play the same game again.
- ▶ A *bundle 2-gerbe* (L, Z, Y, X) is a “simplicial bundle gerbe”

$$\begin{array}{ccccccc}
 & & \mathbb{L} & & \delta\mathbb{L} & & \delta^2\mathbb{L} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\times 2} & Y^{[2]} & \xleftarrow{\times 3} & Y^{[3]} & \xleftarrow{\times 4} & Y^{[4]} & \xleftarrow{\times 5} & Y^{[5]}
 \end{array}$$

- ▶ A locally split map $Y \longrightarrow X$,
- ▶ A gerbe $\mathbb{L} = (L, Z, Y^{[2]})$,
- ▶ A trivialization of $\delta\mathbb{L} = p_0^*\mathbb{L} \otimes p_1^*\mathbb{L}^{-1} \otimes p_2^*\mathbb{L}$ over $Y^{[3]}$,
- ▶ A 2-morphism (did I mention gerbes have 2-morphisms?) relating the induced trivialization of $\delta^2\mathbb{L}$ to the canonical one,
- ▶ A coherency condition on pulled back 2-morphisms over $Y^{[5]}$.
- ▶ (L, Z, Y, X) has a well-defined characteristic class $C(L, Z, Y, X) \in H^4(X; \mathbb{Z})$.
- ▶ For higher gerbes ($H^{\geq 5}(X; \mathbb{Z})$), higher and more complicated coherency conditions will appear.
- ▶ The roles of Y and Z are very asymmetric.

A new version of 2-gerbes:

$$\begin{array}{ccc} & Z & \\ & \downarrow & \\ X & \longleftarrow & Y \end{array}$$

- Start with $Y \longrightarrow X$ and $Z \longrightarrow X$ locally split.

A new version of 2-gerbes:

$$\begin{array}{ccc} Z & \longleftarrow & W \\ \downarrow & & \downarrow \\ X & \longleftarrow & Y \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ locally split.
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ locally split forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger.

A new version of 2-gerbes:

$$\begin{array}{ccccccc}
 & & Z^{[3]} & & & & \\
 & & \Downarrow & & & & \\
 & & \Downarrow & & & & \\
 & & \Downarrow & & & & \\
 & & Z^{[2]} & & & & \\
 & & \Downarrow & & & & \\
 & & \Downarrow & & & & \\
 & & Z & \longleftarrow & W & & \\
 & & \downarrow & & \downarrow & & \\
 & & X & \longleftarrow & Y & \Longleftarrow & Y^{[2]} \Longleftarrow Y^{[3]}
 \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ locally split.
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ locally split forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger.
- ▶ Fill out the diagram by fiber products.

A new version of 2-gerbes:


$$\begin{array}{ccccccc}
 Z^{[3]} & \longleftarrow & W^{[1,3]} & \xleftarrow{\quad} & W^{[2,3]} & \xleftarrow{\quad} & W^{[3,3]} \\
 \Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
 Z^{[2]} & \longleftarrow & W^{[1,2]} & \xleftarrow{\quad} & W^{[2,2]} & \xleftarrow{\quad} & W^{[2,3]} \\
 \Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
 Z & \longleftarrow & W & \xleftarrow{\quad} & W^{[2,1]} & \xleftarrow{\quad} & W^{[3,1]} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\quad} & Y^{[2]} & \xleftarrow{\quad} & Y^{[3]}
 \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ locally split.
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ locally split forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger.
- ▶ Fill out the diagram by fiber products.

A new version of 2-gerbes:


$$\begin{array}{ccccccc}
 Z^{[3]} & \longleftarrow & W^{[1,3]} & \xleftarrow{\quad} & W^{[2,3]} & \xleftarrow{\quad} & W^{[3,3]} \\
 \Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
 Z^{[2]} & \longleftarrow & W^{[1,2]} & \xleftarrow{\quad} & W^{[2,2]} & \xleftarrow{\quad} & W^{[2,3]} \\
 \Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
 Z & \longleftarrow & W & \xleftarrow{\quad} & W^{[2,1]} & \xleftarrow{\quad} & W^{[3,1]} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 X & \longleftarrow & Y & \xleftarrow{\quad} & Y^{[2]} & \xleftarrow{\quad} & Y^{[3]}
 \end{array}$$

- ▶ Start with $Y \rightarrow X$ and $Z \rightarrow X$ locally split.
- ▶ Take $W \rightarrow Y$, $W \rightarrow Z$ locally split forming a commutative square. Minimal choice: $W = Y \times_X Z$, but typically W will be larger.
- ▶ Fill out the diagram by fiber products.
- ▶ $W^{[\bullet, \bullet]}$ forms a *bisimplicial space* over X .



Definition

A bundle bigerbe is a “bisimplicial line bundle” over $W^{[\bullet, \bullet]}$, i.e., a line bundle L over $W^{[2,2]}$, with trivializations of $\delta_0 L$ and $\delta_1 L$, such that the induced trivializations of $\delta_0 \delta_1 L$ agree and which induce the canonical trivializations of $\delta_1^2 L$ and $\delta_0^2 L$.



Definition

A bundle bigerbe is a “bisimplicial line bundle” over $W^{[\bullet, \bullet]}$, i.e., a line bundle L over $W^{[2,2]}$, with trivializations of $\delta_0 L$ and $\delta_1 L$, such that the induced trivializations of $\delta_0 \delta_1 L$ agree and which induce the canonical trivializations of $\delta_1^2 L$ and $\delta_0^2 L$.

- ▶ Products, inverses, pull backs straightforward to define.
- ▶ A *trivialization* is an isomorphism $L \cong \delta_1 Q$ (equivalently $L \cong \delta_0 Q'$) for a line bundle Q over $W^{[1,2]}$ (Q' over $W^{[2,1]}$).

Theorem

A bundle bigerbe (L, W, X) has a well-defined characteristic class $C(L) \in H^4(X; \mathbb{Z})$, with

$$\begin{aligned}C(L^{-1}) &= -C(L), \\C(L \otimes L') &= C(L) + C(L'), \\C(f^* L) &= f^* C(L).\end{aligned}$$

$C(L) = 0$ if and only if L is trivial. $C(L) = C(L')$ if and only if L and L' are stably isomorphic.

Theorem

A bundle bigerbe (L, W, X) has a well-defined characteristic class $C(L) \in H^4(X; \mathbb{Z})$, with

$$\begin{aligned}C(L^{-1}) &= -C(L), \\C(L \otimes L') &= C(L) + C(L'), \\C(f^* L) &= f^* C(L).\end{aligned}$$

$C(L) = 0$ if and only if L is trivial. $C(L) = C(L')$ if and only if L and L' are stably isomorphic.

- This generalizes in a straightforward manner to higher degree (Exercise), leading to *bundle multigerbes*.

Theorem

A bundle multigerbe L of degree n has a well-defined characteristic class $C(L) \in H^{2+n}(X; \mathbb{Z})$, with

$$\begin{aligned}C(L^{-1}) &= -C(L), \\C(L \otimes L') &= C(L) + C(L'), \\C(f^* L) &= f^* C(L).\end{aligned}$$

$C(L) = 0$ if and only if L is trivial. $C(L) = C(L')$ if and only if L and L' are stably isomorphic.

- This generalizes in a straightforward manner to higher degree (Exercise), leading to *bundle multigerbes*.

Taking Čech cochains with respect to a certain class of covers, we obtain a triple complex $(\check{C}^\bullet(W^{[\bullet,\bullet]}; \mathbb{C}^*), d, \delta_0, \delta_1)$.

$$\begin{array}{ccccccc}
 \check{C}^p(Z[3]) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,3]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,3]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,3]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(Z[2]) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,2]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,2]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,2]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(Z) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,1]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,1]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,1]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(X) & \xrightarrow{\delta_0} & \check{C}^p(Y) & \xrightarrow{\delta_0} & \check{C}^p(Y^{[2]}) & \xrightarrow{\delta_0} & \check{C}^p(Y^{[3]})
 \end{array}$$

- ▶ The simplicial complexes $(\check{C}^p(W^{[\bullet,q]}; \mathbb{C}^*), \delta_0)$ and $(\check{C}^p(W^{[q,\bullet]}; \mathbb{C}^*), \delta_1)$ are exact; in fact they admit chain homotopy contractions (commuting with each other, but not with the Čech differential).

Taking Čech cochains with respect to a certain class of covers, we obtain a triple complex $(\check{C}^\bullet(W^{[\bullet,\bullet]}; \mathbb{C}^*), d, \delta_0, \delta_1)$.

$$\begin{array}{ccccccc}
 \check{C}^p(Z[3]) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,3]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,3]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,3]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(Z[2]) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,2]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,2]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,2]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(Z) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,1]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,1]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,1]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(X) & \xrightarrow{\delta_0} & \check{C}^p(Y) & \xrightarrow{\delta_0} & \check{C}^p(Y^{[2]}) & \xrightarrow{\delta_0} & \check{C}^p(Y^{[3]})
 \end{array}$$

- ▶ The simplicial complexes $(\check{C}^p(W^{[\bullet,q]}; \mathbb{C}^*), \delta_0)$ and $(\check{C}^p(W^{[q,\bullet]}; \mathbb{C}^*), \delta_1)$ are exact; in fact they admit chain homotopy contractions (commuting with each other, but not with the Čech differential).
- ▶ The tototal cohomology of $(\check{C}^\bullet(W^{[\bullet,\bullet]}; \mathbb{C}^*), d, \delta_0, \delta_1)$ is isomorphic to $\check{H}^\bullet(X; \mathbb{C}^*)$.

Taking Čech cochains with respect to a certain class of covers, we obtain a triple complex $(\check{C}^\bullet(W^{[\bullet,\bullet]}; \mathbb{C}^*), d, \delta_0, \delta_1)$.

$$\begin{array}{ccccccc}
 \check{C}^p(Z[3]) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,3]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,3]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,3]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(Z[2]) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,2]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,2]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,2]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(Z) & \xrightarrow{\delta_0} & \check{C}^p(W^{[1,1]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[2,1]}) & \xrightarrow{\delta_0} & \check{C}^p(W^{[3,1]}) \\
 \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow & & \delta_1 \uparrow \\
 \check{C}^p(X) & \xrightarrow{\delta_0} & \check{C}^p(Y) & \xrightarrow{\delta_0} & \check{C}^p(Y^{[2]}) & \xrightarrow{\delta_0} & \check{C}^p(Y^{[3]})
 \end{array}$$

- ▶ The simplicial complexes $(\check{C}^p(W^{[\bullet,q]}; \mathbb{C}^*), \delta_0)$ and $(\check{C}^p(W^{[q,\bullet]}; \mathbb{C}^*), \delta_1)$ are exact; in fact they admit chain homotopy contractions (commuting with each other, but not with the Čech differential).
- ▶ The total cohomology of $(\check{C}^\bullet(W^{[\bullet,\bullet]}; \mathbb{C}^*), d, \delta_0, \delta_1)$ is isomorphic to $\check{H}^\bullet(X; \mathbb{C}^*)$.
- ▶ $C(L) \in \check{H}^3(X; \mathbb{C}^*) \cong H^4(X; \mathbb{Z})$ is the image of the pure cocycle $[L] \in \check{C}^1(W^{[2,2]}; \mathbb{C}^*)$.

- ▶ Suppose X is connected, and take $Y = \mathcal{P}_*X$, the based path space.
- ▶ Then $Y^{[2]} = \mathcal{P}_*^{[2]}X \cong \Omega X$, the based loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe (L, \mathcal{P}_*X, X) , i.e., a simplicial line bundle L on ΩX .

- ▶ Suppose X is connected, and take $Y = \mathcal{P}_*X$, the based path space.
- ▶ Then $Y^{[2]} = \mathcal{P}_*^{[2]}X \cong \Omega X$, the based loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe (L, \mathcal{P}_*X, X) , i.e., a simplicial line bundle L on ΩX .
- ▶ In this case, the gerbe product is also known as the “fusion product” (Stolz-Teichner, Waldorf), and L is a “fusion line bundle”.

- ▶ Suppose X is connected, and take $Y = \mathcal{P}_*X$, the based path space.
- ▶ Then $Y^{[2]} = \mathcal{P}_*^{[2]}X \cong \Omega X$, the based loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe (L, \mathcal{P}_*X, X) , i.e., a simplicial line bundle L on ΩX .
- ▶ In this case, the gerbe product is also known as the “fusion product” (Stolz-Teichner, Waldorf), and L is a “fusion line bundle”.
- ▶ Likewise, if X is simply connected, with $Y = Z = \mathcal{P}_*X$ and $W = \mathcal{P}_*\mathcal{P}_*X$, Then $W^{[2,2]} = \Omega^2 X$, the double based loop space of X .
- ▶ Every class in $H^4(X; \mathbb{Z})$ is represented by a bundle bigerbe $(L, \mathcal{P}_*X, \mathcal{P}_*\mathcal{P}_*X, X)$, equivalently a “doubly fusion” line bundle $L \longrightarrow \Omega^2 X$. (c.f. Carey Johnson Murray Stevenson Wang)

- ▶ Suppose X is connected, and take $Y = \mathcal{P}_*X$, the based path space.
- ▶ Then $Y^{[2]} = \mathcal{P}_*^{[2]}X \cong \Omega X$, the based loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe (L, \mathcal{P}_*X, X) , i.e., a simplicial line bundle L on ΩX .
- ▶ In this case, the gerbe product is also known as the “fusion product” (Stolz-Teichner, Waldorf), and L is a “fusion line bundle”.
- ▶ Likewise, if X is simply connected, with $Y = Z = \mathcal{P}_*X$ and $W = \mathcal{P}_*\mathcal{P}_*X$, Then $W^{[2,2]} = \Omega^2 X$, the double based loop space of X .
- ▶ Every class in $H^4(X; \mathbb{Z})$ is represented by a bundle bigerbe $(L, \mathcal{P}_*X, \mathcal{P}_*\mathcal{P}_*X, X)$, equivalently a “doubly fusion” line bundle $L \longrightarrow \Omega^2 X$. (c.f. Carey Johnson Murray Stevenson Wang)

Proposition

If X is k -connected, then every class in $H^{3+k}(X; \mathbb{Z})$ is represented by a multigerbe on $\Omega^{2+k}X$, (aka a $2 + k$ -fold fusion line bundle).

- ▶ Alternatively, take $Y = \mathcal{P}X$, the free path space, fibering over X^2 .
- ▶ Then $Y^{[2]} = \mathcal{P}^{[2]}X \cong \mathcal{L}X$, the free loop space.

- ▶ Alternatively, take $Y = \mathcal{P}X$, the free path space, fibering over X^2 .
- ▶ Then $Y^{[2]} = \mathcal{P}^{[2]}X \cong \mathcal{L}X$, the free loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe $L = (L, \mathcal{P}X, X^2)$ on $\mathcal{L}X$, with the additional condition of a trivialization of the alternating product of pullbacks to the “figure-of-eight” loop space [K-Melrose, 2013].
- ▶ Figure-of-eight is yet another simplicial condition “over” the simplicial space

$$X \rightrightarrows X^2 \rightrightarrows X^3 \cdots$$

guaranteeing that the class in $H^3(X^2; \mathbb{Z})$ comes from $H^3(X; \mathbb{Z})$.

- ▶ Alternatively, take $Y = \mathcal{P}X$, the free path space, fibering over X^2 .
- ▶ Then $Y^{[2]} = \mathcal{P}^{[2]}X \cong \mathcal{L}X$, the free loop space.
- ▶ Every class in $H^3(X; \mathbb{Z})$ is represented by a bundle gerbe $L = (L, \mathcal{P}X, X^2)$ on $\mathcal{L}X$, with the additional condition of a trivialization of the alternating product of pullbacks to the “figure-of-eight” loop space [K-Melrose, 2013].
- ▶ Figure-of-eight is yet another simplicial condition “over” the simplicial space

$$X \rightrightarrows X^2 \rightrightarrows X^3 \cdots$$

guaranteeing that the class in $H^3(X^2; \mathbb{Z})$ comes from $H^3(X; \mathbb{Z})$.

Proposition

Every class in $H^{3+k}(X; \mathbb{Z})$ is represented by a multisimplicial (and multi figure-of-eight) line bundle on $\mathcal{L}^{2+k}X$.

Transgression and loop-fusion cohomology

Multigerbes

Chris
Kottke

Gerbes

Higher
gerbes

Relation to
loop
spaces

- ▶ Take $\alpha \in H^3(X; \mathbb{Z})$ and $L \rightarrow \mathcal{L}X$ with $DD(L, \mathcal{P}X, X^2) = \alpha$.
- ▶ $c_1(L) \in H^2(\mathcal{L}X; \mathbb{Z})$ is the *transgression* of α :

$$\begin{array}{ccc} H^k(X; \mathbb{Z}) & \xrightarrow{\text{ev}^*} & H^k(\mathbb{S}^1 \times \mathcal{L}X; \mathbb{Z}) \\ & \searrow \tau & \downarrow \int_{\mathbb{S}^1} \\ & & H^{k-1}(\mathcal{L}X; \mathbb{Z}) \end{array}$$

- ▶ Loses information since it forgets the simplicial properties of L .

- ▶ Take $\alpha \in H^3(X; \mathbb{Z})$ and $L \rightarrow \mathcal{L}X$ with $DD(L, \mathcal{P}X, X^2) = \alpha$.
- ▶ $c_1(L) \in H^2(\mathcal{L}X; \mathbb{Z})$ is the *transgression* of α :

$$\begin{array}{ccc} H^k(X; \mathbb{Z}) & \xrightarrow{\text{ev}^*} & H^k(\mathbb{S}^1 \times \mathcal{L}X; \mathbb{Z}) \\ & \searrow \tau & \downarrow \int_{\mathbb{S}^1} \\ & & H^{k-1}(\mathcal{L}X; \mathbb{Z}) \end{array}$$

- ▶ Loses information since it forgets the simplicial properties of L .
- ▶ [K.-Melrose, 2015]: “Loop-fusion” Čech cohomology $H_{\text{lf}}^\bullet(\mathcal{L}X; \mathbb{Z})$ such that transgression factors through an isomorphism

$$H^k(X; \mathbb{Z}) \xrightarrow{\cong} H_{\text{lf}}^{k-1}(\mathcal{L}X; \mathbb{Z}) \longrightarrow H^{k-1}(\mathcal{L}X; \mathbb{Z}).$$

- ▶ Take $\alpha \in H^3(X; \mathbb{Z})$ and $L \rightarrow \mathcal{L}X$ with $DD(L, \mathcal{P}X, X^2) = \alpha$.
- ▶ $c_1(L) \in H^2(\mathcal{L}X; \mathbb{Z})$ is the *transgression* of α :

$$\begin{array}{ccc} H^k(X; \mathbb{Z}) & \xrightarrow{\text{ev}^*} & H^k(\mathbb{S}^1 \times \mathcal{L}X; \mathbb{Z}) \\ & \searrow \tau & \downarrow \int_{\mathbb{S}^1} \\ & & H^{k-1}(\mathcal{L}X; \mathbb{Z}) \end{array}$$

- ▶ Loses information since it forgets the simplicial properties of L .
- ▶ [K.-Melrose, 2015]: “Loop-fusion” Čech cohomology $H_{\text{lf}}^\bullet(\mathcal{L}X; \mathbb{Z})$ such that transgression factors through an isomorphism

$$H^k(X; \mathbb{Z}) \xrightarrow{\cong} H_{\text{lf}}^{k-1}(\mathcal{L}X; \mathbb{Z}) \longrightarrow H^{k-1}(\mathcal{L}X; \mathbb{Z}).$$

Theorem

On $\mathcal{L}^\ell X$ there is a well-defined loop-fusion cohomology $\check{H}_{\text{lf}}^\bullet(\mathcal{L}^\ell X; \mathbb{Z})$ through which iterated transgression factors as an isomorphism:

$$H_{\text{lf}}^k(\mathcal{L}^\ell X; \mathbb{Z}) \xrightarrow{\cong} H_{\text{lf}}^{k-n}(\mathcal{L}^{\ell+n} X; \mathbb{Z}).$$

- ▶ Let X be a manifold and $E \rightarrow X$ a principal G bundle for a simple, simply connected Lie group G (e.g. $G = \text{Spin}$).
- ▶ Then $\mathcal{L}E \rightarrow \mathcal{L}X$ is a $\mathcal{L}G$ bundle, and $\mathcal{L}G$ has a central extension

$$1 \rightarrow \text{U}(1) \rightarrow \widehat{\mathcal{L}G} \rightarrow \mathcal{L}G \rightarrow 1$$

- ▶ Let X be a manifold and $E \rightarrow X$ a principal G bundle for a simple, simply connected Lie group G (e.g. $G = \text{Spin}$).
- ▶ Then $\mathcal{L}E \rightarrow \mathcal{L}X$ is a $\mathcal{L}G$ bundle, and $\mathcal{L}G$ has a central extension

$$1 \rightarrow \text{U}(1) \rightarrow \widehat{\mathcal{L}G} \rightarrow \mathcal{L}G \rightarrow 1$$

- ▶ A lift of $\mathcal{L}E$ to a $\widehat{\mathcal{L}G}$ -bundle may be thought of as a “spin structure” on loop space [Atiyah, Witten].

- ▶ Let X be a manifold and $E \rightarrow X$ a principal G bundle for a simple, simply connected Lie group G (e.g. $G = \text{Spin}$).
- ▶ Then $\mathcal{L}E \rightarrow \mathcal{L}X$ is a $\mathcal{L}G$ bundle, and $\mathcal{L}G$ has a central extension

$$1 \rightarrow \text{U}(1) \rightarrow \widehat{\mathcal{L}G} \rightarrow \mathcal{L}G \rightarrow 1$$

- ▶ A lift of $\mathcal{L}E$ to a $\widehat{\mathcal{L}G}$ -bundle may be thought of as a “spin structure” on loop space [Atiyah, Witten].

Proposition

The lifting bundle gerbe $(u^\widehat{\mathcal{L}G}, \mathcal{L}E, \mathcal{L}X)$ is a bundle bigerbe associated to the bisimplicial space generated by*

$$\begin{array}{ccc} E^2 & \longleftarrow & \mathcal{P}E \\ \downarrow & & \downarrow \\ X^2 & \longleftarrow & \mathcal{P}X \end{array}$$

with Dixmier-Douady class $\frac{1}{2}p_1(E) \in H^4(X; \mathbb{Z})$.

- ▶ c.f. McLaughlin, Redden, CJMSW, Waldorf, K.-Melrose.

- ▶ Connection structures, representations of differential cohomology when X, Y, Z, W are manifolds.
- ▶ Satisfactory notion of morphisms for multigerbes.
- ▶ On $\mathcal{L}X$ (and generally $\mathcal{L}^k X$), equivariance of L with respect to action of $\text{Diffeo}^+(\mathbb{S}^1)$ (and its central extension) [c.f. Brylinski].
- ▶ Loop-fusion K-theory of $\mathcal{L}X$ and $\mathcal{L}^\ell X$.