

# Computational Fluid Dynamics - Part I

## Discretization Techniques - Elliptic Problems

Marc Gerritsma

`M.I.Gerritsma@TUDelft.nl`

Department of Aerospace Engineering  
Delft University of Technology

September 25, 2007/ MSc Aerospace Engineering

# Outline

- 1 Finite Difference Formulation
- 2 Finite Volume Method
- 3 Finite Element Method
- 4 Summary

# Outline of this lecture

## What we do and do not do

- General techniques for elliptic equations
- Finite Difference Method, Finite Volume Method and Finite Element Method
- Mixed problems
- Transformation to general domains
- We will **not** discuss Section 3.1.1 from the lecture notes.

# Outline of this lecture

## What we do and do not do

- General techniques for elliptic equations
- Finite Difference Method, Finite Volume Method and Finite Element Method
- Mixed problems
- Transformation to general domains
- We will **not** discuss Section 3.1.1 from the lecture notes.

# Outline of this lecture

## What we do and do not do

- General techniques for elliptic equations
- Finite Difference Method, Finite Volume Method and Finite Element Method
- Mixed problems
- Transformation to general domains
- We will **not** discuss Section 3.1.1 from the lecture notes.

# Outline of this lecture

## What we do and do not do

- General techniques for elliptic equations
- Finite Difference Method, Finite Volume Method and Finite Element Method
- Mixed problems
- Transformation to general domains
- We will **not** discuss Section 3.1.1 from the lecture notes.

# Outline of this lecture

## What we do and do not do

- General techniques for elliptic equations
- Finite Difference Method, Finite Volume Method and Finite Element Method
- Mixed problems
- Transformation to general domains
- We will **not** discuss Section 3.1.1 from the lecture notes.

# Outline of this lecture

## What we do and do not do

- General techniques for elliptic equations
- Finite Difference Method, Finite Volume Method and Finite Element Method
- Mixed problems
- Transformation to general domains
- We will **not** discuss Section 3.1.1 from the lecture notes.

## Approximation

- **Quantitative** approximation (small errors)
- **Qualitative** approximation (preserve symmetries of differential equations)



# Outline of this lecture

## What we do and do not do

- General techniques for elliptic equations
- Finite Difference Method, Finite Volume Method and Finite Element Method
- Mixed problems
- Transformation to general domains
- We will **not** discuss Section 3.1.1 from the lecture notes.

## Approximation

- **Quantitative** approximation (small errors)
- **Qualitative** approximation (preserve symmetries of differential equations)

# Elliptic problems

## The Laplace equation

### The Laplace equation

The Laplace equation for  $u$  is given by

$$\Delta u = \nabla \cdot \nabla u = 0 \quad \Rightarrow \quad \begin{cases} \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 & 2D \\ \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0 & 3D \end{cases}$$

# Elliptic problems

## The Laplace equation

### The Laplace equation

The Laplace equation for  $u$  is given by

$$\Delta u = \nabla \cdot \nabla u = 0 \quad \Rightarrow \quad \begin{cases} \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 & 2D \\ \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0 & 3D \end{cases}$$

### First order system

Introducing the vector  $\vec{w}$  we can write this as a first order system (mixed problem):

$$\vec{\nabla} \cdot \vec{w} = 0 ,$$

$$\vec{w} = \vec{\nabla} u .$$

# Finite Difference Method

## Laplace Equation I

### 2D FD method

- Assume we have a 2D **uniform** grid
- Discretize solution at  $u_{i,j} = u(i * \Delta x, j * \Delta y)$
- Relate the value in neighboring point by **Taylor series expansion**

$$u_{i-1} = u_i - \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i - \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

$$u_{i+1} = u_i + \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i + \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

### Graphical Representation

# Finite Difference Method

## Laplace Equation I


### 2D FD method

- Assume we have a 2D **uniform** grid
- Discretize solution at  $u_{i,j} = u(i * \Delta x, j * \Delta y)$
- Relate the value in neighboring point by **Taylor series expansion**

$$u_{i-1} = u_i - \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i - \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

$$u_{i+1} = u_i + \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i + \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

### Graphical Representation



# Finite Difference Method

## Laplace Equation I


### 2D FD method

- Assume we have a 2D **uniform** grid
- Discretize solution at  $u_{i,j} = u(i * \Delta x, j * \Delta y)$
- Relate the value in neighboring point by **Taylor series expansion**

$$u_{i-1} = u_i - \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i - \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

$$u_{i+1} = u_i + \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i + \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

### Graphical Representation



# Finite Difference Method

## Laplace Equation I


### 2D FD method

- Assume we have a 2D **uniform** grid
- Discretize solution at  $u_{i,j} = u(i * \Delta x, j * \Delta y)$
- Relate the value in neighboring point by **Taylor series expansion**

$$u_{i-1} = u_i - \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i - \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

$$u_{i+1} = u_i + \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i + \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

### Graphical Representation



# Finite Difference Method

## Laplace Equation I


### 2D FD method

- Assume we have a 2D **uniform** grid
- Discretize solution at  $u_{i,j} = u(i * \Delta x, j * \Delta y)$
- Relate the value in neighboring point by **Taylor series expansion**

$$u_{i-1} = u_i - \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i - \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

$$u_{i+1} = u_i + \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i + \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

### Graphical Representation





# Finite Difference Method

## Laplace Equation I


### 2D FD method

- Assume we have a 2D **uniform** grid
- Discretize solution at  $u_{i,j} = u(i * \Delta x, j * \Delta y)$
- Relate the value in neighboring point by **Taylor series expansion**

$$u_{i-1} = u_i - \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i - \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

$$u_{i+1} = u_i + \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i + \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

### Graphical Representation



# Finite Difference Method

## Laplace Equation I


### 2D FD method

- Assume we have a 2D **uniform** grid
- Discretize solution at  $u_{i,j} = u(i * \Delta x, j * \Delta y)$
- Relate the value in neighboring point by **Taylor series expansion**

$$u_{i-1} = u_i - \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i - \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

$$u_{i+1} = u_i + \Delta x \left. \frac{\partial u}{\partial x} \right|_i + \frac{\Delta x^2}{2} \left. \frac{\partial^2 u}{\partial x^2} \right|_i + \frac{\Delta x^3}{6} \left. \frac{\partial^3 u}{\partial x^3} \right|_i + \frac{\Delta x^4}{24} \left. \frac{\partial^4 u}{\partial x^4} \right|_i + O(\Delta x^5)$$

### Graphical Representation



### Exact expansion $\partial^2 u / \partial x^2$

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} = \frac{\partial^2 u}{\partial x^2} + \frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} + O(\Delta x^4)$$

# Finite Difference Method

## Laplace Equation II

Exact expansion  $\partial^2 u / \partial x^2$

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} = \frac{\partial^2 u}{\partial x^2} + \frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} + O(\Delta x^4) .$$

# Finite Difference Method

## Laplace Equation II

Exact expansion  $\partial^2 u / \partial x^2$

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} = \frac{\partial^2 u}{\partial x^2} + \frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} + O(\Delta x^4) .$$

Approximation  $\partial^2 u / \partial x^2$

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} \approx \frac{\partial^2 u}{\partial x^2} .$$

# Finite Difference Method

## Laplace Equation II

Exact expansion  $\partial^2 u / \partial x^2$

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} = \frac{\partial^2 u}{\partial x^2} + \frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} + O(\Delta x^4) .$$

Approximation  $\partial^2 u / \partial x^2$

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} \approx \frac{\partial^2 u}{\partial x^2} .$$

Truncation error  $\partial^2 u / \partial x^2$

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} - \frac{\partial^2 u}{\partial x^2} = \underbrace{\frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4}}_{\text{Truncation error}} + O(\Delta x^4) .$$

# Finite Difference Method

## Laplace Equation III

### Laplace equation

$$0 = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = \frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{\Delta y^2} - \underbrace{\frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} - \frac{\Delta y^2}{12} \frac{\partial^4 u}{\partial y^4}}_{\text{Truncation Error}} + O(\Delta x^4, \Delta y^4) .$$

# Finite Difference Method

## Laplace Equation III

### Laplace equation

$$0 = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = \frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{\Delta y^2} - \underbrace{\frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} - \frac{\Delta y^2}{12} \frac{\partial^4 u}{\partial y^4}}_{\text{Truncation Error}} + O(\Delta x^4, \Delta y^4) .$$

### Consistency

A numerical scheme is called **consistent** if the truncation error goes to zero for the mesh to zero

$$\lim_{\Delta x, \Delta y \rightarrow 0} \left[ \frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} - \frac{\Delta y^2}{12} \frac{\partial^4 u}{\partial y^4} + O(\Delta x^4, \Delta y^4) \right] = 0 .$$

# Finite Difference Method

## Laplace Equation IV

### Modified/equivalent differential equation

By solving

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{\Delta y^2} = 0 ,$$

we actually approximate

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} + \frac{\Delta y^2}{12} \frac{\partial^4 u}{\partial y^4} = 0 ,$$

up to  $O(\Delta x^4, \Delta y^4)$ .



# Finite Difference Method

## Laplace Equation IV

### Modified/equivalent differential equation

By solving

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{\Delta y^2} = 0 ,$$

we actually approximate

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} + \frac{\Delta y^2}{12} \frac{\partial^4 u}{\partial y^4} = 0 ,$$

up to  $O(\Delta x^4, \Delta y^4)$ .

### Dirichlet Boundary Conditions

If on the lefthand side of the domain  $u_{0,j} = u_j^*$  is prescribed then

$$\frac{u_{2,j} - 2u_{1,j}}{\Delta x^2} + \frac{u_{1,j+1} - 2u_{1,j} + u_{1,j-1}}{\Delta y^2} = -\frac{u_j^*}{\Delta x^2} .$$

Analogously for other boundaries

# Finite Difference Method

## Laplace Equation IV

### Modified/equivalent differential equation

By solving

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{\Delta y^2} = 0 ,$$

we actually approximate

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\Delta x^2}{12} \frac{\partial^4 u}{\partial x^4} + \frac{\Delta y^2}{12} \frac{\partial^4 u}{\partial y^4} = 0 ,$$

up to  $O(\Delta x^4, \Delta y^4)$ .

### Neumann Boundary Conditions

If on the lefthand side of the domain  $\partial u / \partial x(0, j) = A_j$  is prescribed then

$$\frac{2u_{1,j} - 2u_{0,j}}{\Delta x^2} + \frac{u_{0,j+1} - 2u_{0,j} + u_{0,j-1}}{\Delta y^2} = \frac{2A_j}{\Delta x} .$$

Analogously for other boundaries

# Finite Difference Method

## Laplace Equation V

### Example on a $3 \times 3$ -grid, Dirichlet

Let  $\Delta x = \Delta y = h$

$$\frac{1}{h^2} \begin{pmatrix} -4 & 1 & 0 & 1 \\ 1 & -4 & 1 & 0 \\ 0 & 1 & -4 & 1 \\ 1 & 0 & 1 & -4 \end{pmatrix} \begin{pmatrix} u_{1,1} \\ u_{2,1} \\ u_{2,2} \\ u_{1,2} \end{pmatrix} = \frac{1}{h^2} \begin{pmatrix} -u_{0,1}^* - u_{1,0}^* \\ -u_{2,0}^* - u_{3,1}^* \\ -u_{3,2}^* - u_{2,3}^* \\ -u_{1,3}^* - u_{0,2}^* \end{pmatrix}$$

# Finite Difference Method

## Laplace Equation V

### Example on a $3 \times 3$ -grid, Dirichlet

Let  $\Delta x = \Delta y = h$

$$\frac{1}{h^2} \begin{pmatrix} -4 & 1 & 0 & 1 \\ 1 & -4 & 1 & 0 \\ 0 & 1 & -4 & 1 \\ 1 & 0 & 1 & -4 \end{pmatrix} \begin{pmatrix} u_{1,1} \\ u_{2,1} \\ u_{2,2} \\ u_{1,2} \end{pmatrix} = \frac{1}{h^2} \begin{pmatrix} -u_{0,1}^* - u_{1,0}^* \\ -u_{2,0}^* - u_{3,1}^* \\ -u_{3,2}^* - u_{2,3}^* \\ -u_{1,3}^* - u_{0,2}^* \end{pmatrix}$$

### Symmetry

Note that the matrix in this example is **symmetric**. This was to be expected, because the Laplace operator is a **symmetric operator**.

# Finite Difference Method

## General domain

### Transformation


- Suppose we can map square domain onto  $x(s, t)$  and  $y(s, t)$

$$\begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial s} & \frac{\partial y}{\partial s} \\ \frac{\partial x}{\partial t} & \frac{\partial y}{\partial t} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix}$$

$$\begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix} = \frac{1}{J} \begin{pmatrix} \frac{\partial y}{\partial t} & -\frac{\partial y}{\partial s} \\ -\frac{\partial x}{\partial t} & \frac{\partial x}{\partial s} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix}$$

$$J = x_s y_t - x_t y_s.$$

### Mapping



# Finite Difference Method

## General domain

### Transformation


- Suppose we can map square domain onto  $x(s, t)$  and  $y(s, t)$

$$\begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial s} & \frac{\partial y}{\partial s} \\ \frac{\partial x}{\partial t} & \frac{\partial y}{\partial t} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix}$$

$$\begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix} = \frac{1}{J} \begin{pmatrix} \frac{\partial y}{\partial t} & -\frac{\partial y}{\partial s} \\ -\frac{\partial x}{\partial t} & \frac{\partial x}{\partial s} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix}$$

$$J = x_s y_t - x_t y_s.$$

### Mapping



# Finite Difference Method

## General domain

### Transformation


- Suppose we can map square domain onto  $x(s, t)$  and  $y(s, t)$

$$\begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial s} & \frac{\partial y}{\partial s} \\ \frac{\partial x}{\partial t} & \frac{\partial y}{\partial t} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix}$$

$$\begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix} = \frac{1}{J} \begin{pmatrix} \frac{\partial y}{\partial t} & -\frac{\partial y}{\partial s} \\ -\frac{\partial x}{\partial t} & \frac{\partial x}{\partial s} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix}$$

$$J = x_s y_t - x_t y_s.$$

### Mapping



# Finite Difference Method

## General domain

### Transformation


- Suppose we can map square domain onto  $x(s, t)$  and  $y(s, t)$

$$\begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial s} & \frac{\partial y}{\partial s} \\ \frac{\partial x}{\partial t} & \frac{\partial y}{\partial t} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix}$$

$$\begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix} = \frac{1}{J} \begin{pmatrix} \frac{\partial y}{\partial t} & -\frac{\partial y}{\partial s} \\ -\frac{\partial x}{\partial t} & \frac{\partial x}{\partial s} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix}$$

$$J = x_s y_t - x_t y_s.$$

### Mapping





# Finite Difference Method

## General domain

### Transformation


- Suppose we can map square domain onto  $x(s, t)$  and  $y(s, t)$

$$\begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial s} & \frac{\partial y}{\partial s} \\ \frac{\partial x}{\partial t} & \frac{\partial y}{\partial t} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix}$$

$$\begin{pmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{pmatrix} = \frac{1}{J} \begin{pmatrix} \frac{\partial y}{\partial t} & -\frac{\partial y}{\partial s} \\ -\frac{\partial x}{\partial t} & \frac{\partial x}{\partial s} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial s} \\ \frac{\partial}{\partial t} \end{pmatrix}$$

$$J = x_s y_t - x_t y_s.$$

### Mapping



### Laplace equation

$$\begin{aligned} & \frac{1}{J} \frac{\partial y}{\partial t} \frac{\partial}{\partial s} \left[ \frac{1}{J} \frac{\partial y}{\partial t} \frac{\partial u}{\partial s} - \frac{1}{J} \frac{\partial y}{\partial s} \frac{\partial u}{\partial t} \right] - \frac{1}{J} \frac{\partial y}{\partial s} \frac{\partial}{\partial t} \left[ \frac{1}{J} \frac{\partial y}{\partial t} \frac{\partial u}{\partial s} - \frac{1}{J} \frac{\partial y}{\partial s} \frac{\partial u}{\partial t} \right] + \\ & \frac{1}{J} \frac{\partial x}{\partial s} \frac{\partial}{\partial t} \left[ \frac{1}{J} \frac{\partial x}{\partial s} \frac{\partial u}{\partial t} - \frac{1}{J} \frac{\partial x}{\partial t} \frac{\partial u}{\partial s} \right] - \frac{1}{J} \frac{\partial x}{\partial t} \frac{\partial}{\partial s} \left[ \frac{1}{J} \frac{\partial x}{\partial s} \frac{\partial u}{\partial t} - \frac{1}{J} \frac{\partial x}{\partial t} \frac{\partial u}{\partial s} \right] = 0 \end{aligned}$$

# Finite Volume Method

## General Philosophy

- The finite volume method is a method for [conservation laws](#), see Lecture 1.
- The idea is to divide the computational domain into volumes and impose the conservation laws on these volumes.
- Use the [divergence theorem](#) to convert volume integrals to boundary integrals
- Evaluate fluxes at boundaries.

# Finite Volume Method

## General Philosophy

- The finite volume method is a method for **conservation laws**, see Lecture 1.
- The idea is to divide the computational domain into volumes and impose the conservation laws on these volumes.
- Use the **divergence theorem** to convert volume integrals to boundary integrals
- Evaluate fluxes at boundaries.

# Finite Volume Method

## General Philosophy

- The finite volume method is a method for [conservation laws](#), see Lecture 1.
- The idea is to divide the computational domain into volumes and impose the conservation laws on these volumes.
- Use the [divergence theorem](#) to convert volume integrals to boundary integrals
- Evaluate fluxes at boundaries.

# Finite Volume Method

## General Philosophy

- The finite volume method is a method for [conservation laws](#), see Lecture 1.
- The idea is to divide the computational domain into volumes and impose the conservation laws on these volumes.
- Use the [divergence theorem](#) to convert volume integrals to boundary integrals
- Evaluate fluxes at boundaries.

# Finite Volume Method

## General Philosophy

- The finite volume method is a method for [conservation laws](#), see Lecture 1.
- The idea is to divide the computational domain into volumes and impose the conservation laws on these volumes.
- Use the [divergence theorem](#) to convert volume integrals to boundary integrals
- Evaluate fluxes at boundaries.

# Finite Volume Method

## Laplace Equation I

### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$

$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume
- This is still **exact**. The numerical approximation enters into the scheme if **approximate** the **integrals** and the **derivatives**  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation

# Finite Volume Method

## Laplace Equation I


### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$

$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume
- This is still [exact](#). The numerical approximation enters into the scheme if [approximate](#) the [integrals](#) and the [derivatives](#)  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation





# Finite Volume Method

## Laplace Equation I


### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$

$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume
- This is still **exact**. The numerical approximation enters into the scheme if **approximate** the **integrals** and the **derivatives**  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation



# Finite Volume Method

## Laplace Equation I


### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$

$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume
- This is still *exact*. The numerical approximation enters into the scheme if *approximate* the *integrals* and the *derivatives*  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation



# Finite Volume Method

## Laplace Equation I

### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- This is still **exact**. The numerical approximation enters into the scheme if **approximate the integrals** and the **derivatives**  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation



# Finite Volume Method

## Laplace Equation I

### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- This is still **exact**. The numerical approximation enters into the scheme if **approximate** the **integrals** and the **derivatives**  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation



# Finite Volume Method

## Laplace Equation I

### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- This is still **exact**. The numerical approximation enters into the scheme if **approximate** the **integrals** and the **derivatives**  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation



# Finite Volume Method

## Laplace Equation I

### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- This is still **exact**. The numerical approximation enters into the scheme if **approximate** the **integrals** and the **derivatives**  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation



# Finite Volume Method

## Laplace Equation I

### 2D FV method

- Use again a grid with the unknowns defined at the vertices
- Surround the point  $(i, j)$  with a volume  $\Omega_{i,j}$
- Integrate the Laplace equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \nabla \cdot \nabla u \, d\Omega = \int_{\partial\Omega_{i,j}} \nabla u \cdot \mathbf{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- This is still **exact**. The numerical approximation enters into the scheme if **approximate** the **integrals** and the **derivatives**  $\partial u / \partial x$ ,  $\partial u / \partial y$  numerically.

### Graphical Representation



# Finite Volume Method

## Laplace Equation II

### Discretization

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- Approximate integrals by midpoint rule: Let  $y = f(x)$ ,  $x \in [a, b]$  then

$$\int_a^b f(x) dx = f\left(\frac{a+b}{2}\right) h + \frac{h^2}{24} f''(c) \approx f\left(\frac{a+b}{2}\right) h, \quad h = b - a, \quad c \in (a, b)$$

- The derivatives at the midpoints are approximated by finite differences:
- So the finite volume discretization becomes:

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0.$$



# Finite Volume Method

## Laplace Equation II

### Discretization

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- Approximate integrals by midpoint rule: Let  $y = f(x)$ ,  $x \in [a, b]$  then

$$\int_a^b f(x) dx = f\left(\frac{a+b}{2}\right) h + \frac{h^2}{24} f''(c) \approx f\left(\frac{a+b}{2}\right) h, \quad h = b - a, \quad c \in (a, b)$$

- The derivatives at the midpoints are approximated by finite differences:
- So the finite volume discretization becomes:

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0.$$

# Finite Volume Method

## Laplace Equation II

### Discretization

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- Approximate integrals by midpoint rule: Let  $y = f(x)$ ,  $x \in [a, b]$  then

$$\int_a^b f(x) dx = f\left(\frac{a+b}{2}\right) h + \frac{h^2}{24} f''(c) \approx f\left(\frac{a+b}{2}\right) h, \quad h = b - a, \quad c \in (a, b)$$

- The derivatives at the midpoints are approximated by finite differences:

$$\left. \frac{\partial u}{\partial x} \right|_{\text{midpoint I}} \approx \frac{u_{i+1,j} - u_{i,j}}{\Delta x}$$

- So the finite volume discretization becomes:

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0.$$

# Finite Volume Method

## Laplace Equation II

### Discretization

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- Approximate integrals by midpoint rule: Let  $y = f(x)$ ,  $x \in [a, b]$  then

$$\int_a^b f(x) dx = f\left(\frac{a+b}{2}\right) h + \frac{h^2}{24} f''(c) \approx f\left(\frac{a+b}{2}\right) h, \quad h = b - a, \quad c \in (a, b)$$

- The derivatives at the midpoints are approximated by finite differences:

$$\left. \frac{\partial u}{\partial y} \right|_{\text{midpoint II}} \approx \frac{u_{i,j+1} - u_{i,j}}{\Delta y}$$

- So the finite volume discretization becomes:

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0.$$

# Finite Volume Method

## Laplace Equation II

### Discretization

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- Approximate integrals by midpoint rule: Let  $y = f(x)$ ,  $x \in [a, b]$  then

$$\int_a^b f(x) dx = f\left(\frac{a+b}{2}\right) h + \frac{h^2}{24} f''(c) \approx f\left(\frac{a+b}{2}\right) h, \quad h = b - a, \quad c \in (a, b)$$

- The derivatives at the midpoints are approximated by finite differences:

$$\left. \frac{\partial u}{\partial x} \right|_{\text{midpoint III}} \approx \frac{u_{i,j} - u_{i-1,j}}{\Delta x}$$

- So the finite volume discretization becomes:

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0.$$

# Finite Volume Method

## Laplace Equation II

### Discretization

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- Approximate integrals by midpoint rule: Let  $y = f(x)$ ,  $x \in [a, b]$  then

$$\int_a^b f(x) dx = f\left(\frac{a+b}{2}\right) h + \frac{h^2}{24} f''(c) \approx f\left(\frac{a+b}{2}\right) h, \quad h = b - a, \quad c \in (a, b)$$

- The derivatives at the midpoints are approximated by finite differences:

$$\left. \frac{\partial u}{\partial y} \right|_{\text{midpoint IV}} \approx \frac{u_{i,j} - u_{i,j-1}}{\Delta y}$$

- So the finite volume discretization becomes:

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0.$$

# Finite Volume Method

## Laplace Equation II

### Discretization

$$0 = \int_I \frac{\partial u}{\partial x} dy + \int_{II} \frac{\partial u}{\partial y} dx - \int_{III} \frac{\partial u}{\partial x} dy - \int_{IV} \frac{\partial u}{\partial y} dx$$

- Approximate integrals by midpoint rule: Let  $y = f(x)$ ,  $x \in [a, b]$  then

$$\int_a^b f(x) dx = f\left(\frac{a+b}{2}\right) h + \frac{h^2}{24} f''(c) \approx f\left(\frac{a+b}{2}\right) h, \quad h = b - a, \quad c \in (a, b)$$

- The derivatives at the midpoints are approximated by finite differences:

$$\left. \frac{\partial u}{\partial y} \right|_{\text{midpoint IV}} \approx \frac{u_{i,j} - u_{i,j-1}}{\Delta y}$$

- So the finite volume discretization becomes:

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0.$$

# Finite Volume Method

## Laplace Equation III

### Comparison with FDM

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0 .$$

- If we divide the Finite Volume scheme by the area of  $\Omega_{i,j} = \Delta x \Delta y$  we obtain

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - u_{i,j} + u_{i,j-1}}{\Delta y^2} = 0 .$$

This is, **in this particular case**, the same as the Finite Difference scheme obtained earlier.

- Note that we started with a second order PDE, but due to the conversion to boundary integrals we only need to evaluate first order derivatives.
- Note that the midpoint rule is **second order accurate** and the approximation of the derivatives is **second order accurate**, so the **overall scheme is second order**.

# Finite Volume Method

## Laplace Equation III

### Comparison with FDM

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0 .$$

- If we divide the Finite Volume scheme by the area of  $\Omega_{i,j} = \Delta x \Delta y$  we obtain

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - u_{i,j} + u_{i,j-1}}{\Delta y^2} = 0 .$$

This is, **in this particular case**, the same as the Finite Difference scheme obtained earlier.

- Note that we started with a second order PDE, but due to the conversion to boundary integrals we only need to evaluate first order derivatives.
- Note that the midpoint rule is **second order accurate** and the approximation of the derivatives is **second order accurate**, so the **overall scheme is second order**.



# Finite Volume Method

## Laplace Equation III

### Comparison with FDM

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0 .$$

- If we divide the Finite Volume scheme by the area of  $\Omega_{i,j} = \Delta x \Delta y$  we obtain

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - u_{i,j} + u_{i,j-1}}{\Delta y^2} = 0 .$$

This is, **in this particular case**, the same as the Finite Difference scheme obtained earlier.

- Note that we started with a second order PDE, but due to the conversion to boundary integrals we only need to evaluate first order derivatives.
- Note that the midpoint rule is **second order accurate** and the approximation of the derivatives is **second order accurate**, so the **overall scheme is second order**.

# Finite Volume Method

## Laplace Equation III

### Comparison with FDM

$$\frac{u_{i+1,j} - u_{i,j}}{\Delta x} \Delta y + \frac{u_{i,j+1} - u_{i,j}}{\Delta y} \Delta x - \frac{u_{i,j} - u_{i-1,j}}{\Delta x} \Delta y - \frac{u_{i,j} - u_{i,j-1}}{\Delta y} \Delta x = 0 .$$

- If we divide the Finite Volume scheme by the area of  $\Omega_{i,j} = \Delta x \Delta y$  we obtain

$$\frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} + \frac{u_{i,j+1} - u_{i,j} + u_{i,j-1}}{\Delta y^2} = 0 .$$

This is, **in this particular case**, the same as the Finite Difference scheme obtained earlier.

- Note that we started with a second order PDE, but due to the conversion to boundary integrals we only need to evaluate first order derivatives.
- Note that the midpoint rule is **second order accurate** and the approximation of the derivatives is **second order accurate**, so the **overall scheme is second order**.

# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$

$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$

$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume
- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation

# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$


$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$

$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume
- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation



# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$


$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$

$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume
- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation



# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$


$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$

$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume
- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation



# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$

$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I w_1 \, dy + \int_{II} w_2 \, dx - \int_{III} w_1 \, dy - \int_{IV} w_2 \, dx$$

- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation



# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$

$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I w_1 \, dy + \int_{II} w_2 \, dx - \int_{III} w_1 \, dy - \int_{IV} w_2 \, dx$$

- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation





# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$

$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I w_1 \, dy + \int_{II} w_2 \, dx - \int_{III} w_1 \, dy - \int_{IV} w_2 \, dx$$

- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation



# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$

$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I w_1 \, dy + \int_{II} w_2 \, dx - \int_{III} w_1 \, dy - \int_{IV} w_2 \, dx$$

- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation



# Finite Volume Method

## Laplace Equation IV

### Mixed Formulation I

- Recall the **mixed formulation** of the Laplace equation

$$\vec{\nabla} \cdot \vec{w} = 0, \quad \text{with } \vec{w} = (w_1, w_2)^T,$$

$$\vec{w} = \vec{\nabla} u.$$

- Use the same volume  $\Omega_{i,j}$
- Integrate the  $\vec{\nabla} \cdot \vec{w}$  equation over  $\Omega_{i,j}$


$$0 = \int_{\Omega_{i,j}} \vec{\nabla} \cdot \vec{w} \, d\Omega = \int_{\partial\Omega_{i,j}} \vec{w} \cdot \vec{n} \, dS$$

- Split this boundary integral over the sides of the volume

$$0 = \int_I w_1 \, dy + \int_{II} w_2 \, dx - \int_{III} w_1 \, dy - \int_{IV} w_2 \, dx$$

- Note that both components of the vector  $\vec{w}$  are located at different places in the grid.  $u$  at the vertices,  $w_1$  at the vertical edges,  $w_2$  at the horizontal edges. Such a grid is called a **staggered grid**.

### Graphical Representation




# Finite Volume Method

## Laplace Equation V

### Mixed Formulation II

- The staggered grid arises naturally from the finite volume approach. Not so easy to deduce in finite difference method.
- The equation  $\vec{\nabla} \cdot \vec{w} = 0$  is called the **conservation law** and the vector  $\vec{w}$  is called the **flux vector**.
- The equation  $\vec{w} = \vec{\nabla} u$  is called the **constitutive equation**, which relates the primary unknowns to the fluxes.
- If all unknowns  $u$ ,  $w_1$  and  $w_2$  were defined at the same point – no staggered grid – the linear system is singular and solution will be polluted with artificial oscillations.
- The placement of the various unknowns in the grid is important. For incompressible flows staggered grids play an important role.

### Graphical Representation




# Finite Volume Method

## Laplace Equation V

### Mixed Formulation II

- The staggered grid arises naturally from the finite volume approach. Not so easy to deduce in finite difference method.
- The equation  $\vec{\nabla} \cdot \vec{w} = 0$  is called the **conservation law** and the vector  $\vec{w}$  is called the **flux vector**.
- The equation  $\vec{w} = \vec{\nabla} u$  is called the **constitutive equation**, which relates the primary unknowns to the fluxes.
- If all unknowns  $u$ ,  $w_1$  and  $w_2$  were defined at the same point – no staggered grid – the linear system is singular and solution will be polluted with artificial oscillations.
- The placement of the various unknowns in the grid is important. For incompressible flows staggered grids play an important role.

### Graphical Representation




# Finite Volume Method

## Laplace Equation V

### Mixed Formulation II

- The staggered grid arises naturally from the finite volume approach. Not so easy to deduce in finite difference method.
- The equation  $\vec{\nabla} \cdot \vec{w} = 0$  is called the **conservation law** and the vector  $\vec{w}$  is called the **flux vector**.
- The equation  $\vec{w} = \vec{\nabla} u$  is called the **constitutive equation**, which relates the primary unknowns to the fluxes.
- If all unknowns  $u$ ,  $w_1$  and  $w_2$  were defined at the same point – no staggered grid – the linear system is singular and solution will be polluted with artificial oscillations.
- The placement of the various unknowns in the grid is important. For incompressible flows staggered grids play an important role.

### Graphical Representation




# Finite Volume Method

## Laplace Equation V

### Mixed Formulation II

- The staggered grid arises naturally from the finite volume approach. Not so easy to deduce in finite difference method.
- The equation  $\vec{\nabla} \cdot \vec{w} = 0$  is called the **conservation law** and the vector  $\vec{w}$  is called the **flux vector**.
- The equation  $\vec{w} = \vec{\nabla} u$  is called the **constitutive equation**, which relates the primary unknowns to the fluxes.
- If all unknowns  $u$ ,  $w_1$  and  $w_2$  were defined at the same point – no staggered grid – the linear system is singular and solution will be polluted with artificial oscillations.
- The placement of the various unknowns in the grid is important. For incompressible flows staggered grids play an important role.

### Graphical Representation




# Finite Volume Method

## Laplace Equation V

### Mixed Formulation II

- The staggered grid arises naturally from the finite volume approach. Not so easy to deduce in finite difference method.
- The equation  $\vec{\nabla} \cdot \vec{w} = 0$  is called the **conservation law** and the vector  $\vec{w}$  is called the **flux vector**.
- The equation  $\vec{w} = \vec{\nabla} u$  is called the **constitutive equation**, which relates the primary unknowns to the fluxes.
- If all unknowns  $u$ ,  $w_1$  and  $w_2$  were defined at the same point – no staggered grid – the linear system is singular and solution will be polluted with artificial oscillations.
- The placement of the various unknowns in the grid is important. For incompressible flows staggered grids play an important role.

### Graphical Representation






# Finite Volume Method

## Laplace Equation V

### Mixed Formulation II

- The staggered grid arises naturally from the finite volume approach. Not so easy to deduce in finite difference method.
- The equation  $\vec{\nabla} \cdot \vec{w} = 0$  is called the **conservation law** and the vector  $\vec{w}$  is called the **flux vector**.
- The equation  $\vec{w} = \vec{\nabla} u$  is called the **constitutive equation**, which relates the primary unknowns to the fluxes.
- If all unknowns  $u$ ,  $w_1$  and  $w_2$  were defined at the same point – no staggered grid – the linear system is singular and solution will be polluted with artificial oscillations.
- The placement of the various unknowns in the grid is important. For incompressible flows staggered grids play an important role.

### Graphical Representation



# Finite Element Method I

## What is different

- The Finite Difference Method is a **collocation method**. This is a purely **nodal** method in which all unknowns are defined in discrete points
- The Finite Volume Method is based on an **integral formulation**.
- The Finite Element Method is a **Residual based** method.
- The idea in Finite Element Methods is to expand the unknown solution in terms of known basis functions and to determine the coefficients of this expansion such that the residual is 'small'.

# Finite Element Method I

## What is different

- The Finite Difference Method is a **collocation method**. This is a purely **nodal** method in which all unknowns are defined in discrete points
- The Finite Volume Method is based on an **integral formulation**.
- The Finite Element Method is a **Residual based** method.
- The idea in Finite Element Methods is to expand the unknown solution in terms of known basis functions and to determine the coefficients of this expansion such that the residual is 'small'.

# Finite Element Method I

## What is different

- The Finite Difference Method is a **collocation method**. This is a purely **nodal** method in which all unknowns are defined in discrete points
- The Finite Volume Method is based on an **integral formulation**.
- The Finite Element Method is a **Residual based** method.
- The idea in Finite Element Methods is to expand the unknown solution in terms of known basis functions and to determine the coefficients of this expansion such that the residual is 'small'.

# Finite Element Method I

## What is different

- The Finite Difference Method is a **collocation method**. This is a purely **nodal** method in which all unknowns are defined in discrete points
- The Finite Volume Method is based on an **integral formulation**.
- The Finite Element Method is a **Residual based** method.
- The idea in Finite Element Methods is to expand the unknown solution in terms of known basis functions and to determine the coefficients of this expansion such that the residual is 'small'.

# Finite Element Method I

## What is different

- The Finite Difference Method is a **collocation method**. This is a purely **nodal** method in which all unknowns are defined in discrete points
- The Finite Volume Method is based on an **integral formulation**.
- The Finite Element Method is a **Residual based** method.
- The idea in Finite Element Methods is to expand the unknown solution in terms of known basis functions and to determine the coefficients of this expansion such that the residual is 'small'.

# Finite Element Method II

## One Element Case

### Basic idea

- Suppose we have  $N$  basis functions  $\phi_1(x, y), \dots, \phi_N(x, y)$  and we expand the solution as

$$u^N(x, y) = \sum_{i=1}^N \alpha_i \phi_i(x, y).$$

- The residual is then defined as

$$R(x, y) = \Delta u^N(x, y) = \frac{\partial^2 u^N}{\partial x^2} + \frac{\partial^2 u^N}{\partial y^2}$$

- Galerkin:** Set the projection of  $R$  onto the basis functions to zero:

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N, \vec{\nabla} \phi_i \, d\Omega + \int_{\partial\Omega} \frac{\partial u^N}{\partial n} \phi_i \, dS \quad i = 1, \dots, N.$$

- This gives us  $N$  equations for the  $N$  unknowns  $\alpha_i$ .

# Finite Element Method II

## One Element Case

### Basic idea

- Suppose we have  $N$  basis functions  $\phi_1(x, y), \dots, \phi_N(x, y)$  and we expand the solution as

$$u^N(x, y) = \sum_{i=1}^N \alpha_i \phi_i(x, y) .$$

- The residual is then defined as

$$R(x, y) = \Delta u^N(x, y) = \frac{\partial^2 u^N}{\partial x^2} + \frac{\partial^2 u^N}{\partial y^2}$$

- Galerkin:** Set the projection of  $R$  onto the basis functions to zero:

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N, \vec{\nabla} \phi_i \, d\Omega + \int_{\partial\Omega} \frac{\partial u^N}{\partial n} \phi_i \, dS \quad i = 1, \dots, N .$$

- This gives us  $N$  equations for the  $N$  unknowns  $\alpha_i$ .



# Finite Element Method II

## One Element Case

### Basic idea

- Suppose we have  $N$  basis functions  $\phi_1(x, y), \dots, \phi_N(x, y)$  and we expand the solution as

$$u^N(x, y) = \sum_{i=1}^N \alpha_i \phi_i(x, y) .$$

- The residual is then defined as

$$R(x, y) = \Delta u^N(x, y) = \frac{\partial^2 u^N}{\partial x^2} + \frac{\partial^2 u^N}{\partial y^2}$$

- Galerkin:** Set the projection of  $R$  onto the basis functions to zero:

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N, \vec{\nabla} \phi_i \, d\Omega + \int_{\partial\Omega} \frac{\partial u^N}{\partial n} \phi_i \, dS \quad i = 1, \dots, N .$$

- This gives us  $N$  equations for the  $N$  unknowns  $\alpha_i$ .

# Finite Element Method II

## One Element Case

### Basic idea

- Suppose we have  $N$  basis functions  $\phi_1(x, y), \dots, \phi_N(x, y)$  and we expand the solution as

$$u^N(x, y) = \sum_{i=1}^N \alpha_i \phi_i(x, y) .$$

- The residual is then defined as

$$R(x, y) = \Delta u^N(x, y) = \frac{\partial^2 u^N}{\partial x^2} + \frac{\partial^2 u^N}{\partial y^2}$$

- Galerkin:** Set the projection of  $R$  onto the basis functions to zero:

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N, \vec{\nabla} \phi_i \, d\Omega + \int_{\partial\Omega} \frac{\partial u^N}{\partial n} \phi_i \, dS \quad i = 1, \dots, N .$$

- This gives us  $N$  equations for the  $N$  unknowns  $\alpha_j$ .

# Finite Element Method II

## One Element Case

### Basic idea

- Suppose we have  $N$  basis functions  $\phi_1(x, y), \dots, \phi_N(x, y)$  and we expand the solution as

$$u^N(x, y) = \sum_{i=1}^N \alpha_i \phi_i(x, y) .$$

- The residual is then defined as

$$R(x, y) = \Delta u^N(x, y) = \frac{\partial^2 u^N}{\partial x^2} + \frac{\partial^2 u^N}{\partial y^2}$$

- Galerkin:** Set the projection of  $R$  onto the basis functions to zero:

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N, \vec{\nabla} \phi_i \, d\Omega + \int_{\partial\Omega} \frac{\partial u^N}{\partial n} \phi_i \, dS \quad i = 1, \dots, N .$$

- This gives us  $N$  equations for the  $N$  unknowns  $\alpha_i$ .

# Finite Element Method III

## One Element Case - continued

### Basic idea

- If Dirichlet boundary conditions are prescribed,  $\phi_i$  needs to be zero at the boundary. Therefore

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i d\Omega = - \int_{\Omega} \vec{\nabla} u^N : \vec{\nabla} \phi_i d\Omega \quad i = 1, \dots, N.$$

- If we insert the expansion for  $u^N$ , we get

$$\int_{\Omega} \vec{\nabla} \left( \sum_{j=1}^N \alpha_j \phi_j \right) : \vec{\nabla} \phi_i d\Omega = \sum_{j=1}^N \alpha_j \int_{\Omega} \vec{\nabla} \phi_j : \vec{\nabla} \phi_i d\Omega := K_{ij} \alpha_j = 0, \quad i = 1, \dots, N.$$

- The matrix  $K_{ij}$  is called **the element matrix**.
- The element matrix is again **symmetric**, i.e.  $K_{ij} = K_{ji}$ .

# Finite Element Method III

## One Element Case - continued

### Basic idea

- If Dirichlet boundary conditions are prescribed,  $\phi_i$  needs to be zero at the boundary. Therefore

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N : \vec{\nabla} \phi_i \, d\Omega \quad i = 1, \dots, N.$$

- If we insert the expansion for  $u^N$ , we get

$$\int_{\Omega} \vec{\nabla} \left( \sum_{j=1}^N \alpha_j \phi_j \right) : \vec{\nabla} \phi_i \, d\Omega = \sum_{j=1}^N \alpha_j \int_{\Omega} \vec{\nabla} \phi_j : \vec{\nabla} \phi_i \, d\Omega := K_{ij} \alpha_j = 0, \quad i = 1, \dots, N.$$

- The matrix  $K_{ij}$  is called **the element matrix**.
- The element matrix is again **symmetric**, i.e.  $K_{ij} = K_{ji}$ .

# Finite Element Method III

## One Element Case - continued

### Basic idea

- If Dirichlet boundary conditions are prescribed,  $\phi_i$  needs to be zero at the boundary. Therefore

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N : \vec{\nabla} \phi_i \, d\Omega \quad i = 1, \dots, N.$$

- If we insert the expansion for  $u^N$ , we get

$$\int_{\Omega} \vec{\nabla} \left( \sum_{j=1}^N \alpha_j \phi_j \right) : \vec{\nabla} \phi_i \, d\Omega = \sum_{j=1}^N \alpha_j \int_{\Omega} \vec{\nabla} \phi_j : \vec{\nabla} \phi_i \, d\Omega := K_{ij} \alpha_j = 0, \quad i = 1, \dots, N.$$

- The matrix  $K_{ij}$  is called **the element matrix**.
- The element matrix is again **symmetric**, i.e.  $K_{ij} = K_{ji}$ .

# Finite Element Method III

## One Element Case - continued

### Basic idea

- If Dirichlet boundary conditions are prescribed,  $\phi_i$  needs to be zero at the boundary. Therefore

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N : \vec{\nabla} \phi_i \, d\Omega \quad i = 1, \dots, N.$$

- If we insert the expansion for  $u^N$ , we get

$$\int_{\Omega} \vec{\nabla} \left( \sum_{j=1}^N \alpha_j \phi_j \right) : \vec{\nabla} \phi_i \, d\Omega = \sum_{j=1}^N \alpha_j \int_{\Omega} \vec{\nabla} \phi_j : \vec{\nabla} \phi_i \, d\Omega := K_{ij} \alpha_j = 0, \quad i = 1, \dots, N.$$

- The matrix  $K_{ij}$  is called **the element matrix**.
- The element matrix is again **symmetric**, i.e.  $K_{ij} = K_{ji}$ .

# Finite Element Method III

## One Element Case - continued

### Basic idea

- If Dirichlet boundary conditions are prescribed,  $\phi_i$  needs to be zero at the boundary. Therefore

$$0 = (R, \phi_i) = \int_{\Omega} \Delta u^N \phi_i \, d\Omega = - \int_{\Omega} \vec{\nabla} u^N : \vec{\nabla} \phi_i \, d\Omega \quad i = 1, \dots, N.$$

- If we insert the expansion for  $u^N$ , we get

$$\int_{\Omega} \vec{\nabla} \left( \sum_{j=1}^N \alpha_j \phi_j \right) : \vec{\nabla} \phi_i \, d\Omega = \sum_{j=1}^N \alpha_j \int_{\Omega} \vec{\nabla} \phi_j : \vec{\nabla} \phi_i \, d\Omega := K_{ij} \alpha_j = 0, \quad i = 1, \dots, N.$$

- The matrix  $K_{ij}$  is called **the element matrix**.
- The element matrix is again **symmetric**, i.e.  $K_{ij} = K_{ji}$ .



# Finite Element Method IV

## One Element Case - continued

### Linear basis functions

- Let us take the domain  $[0, 1] \times [0, 1]$  and the 4 basis functions

$$\phi_1(x, y) = (1 - x)(1 - y)$$

$$\phi_2(x, y) = x(1 - y)$$

$$\phi_3(x, y) = xy \text{ and}$$

$$\phi_4(x, y) = (1 - x)y.$$

- The element  $K_{ij}$  is then given by

$$K_{ij} = \int_0^1 \int_0^1 \left[ \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} + \frac{\partial \phi_i}{\partial y} \frac{\partial \phi_j}{\partial y} \right] dx dy$$

For instance,

$$K_{11} = \int_0^1 \int_0^1 \left[ (1 - y)^2 + (1 - x)^2 \right] dx dy = \frac{2}{3}.$$



$$K = \frac{1}{6} \begin{pmatrix} 4 & -1 & -2 & -1 \\ -1 & 4 & -1 & -2 \\ -2 & -1 & 4 & -1 \\ -1 & -2 & -1 & 4 \end{pmatrix}.$$

# Finite Element Method IV

## One Element Case - continued

### Linear basis functions

- Let us take the domain  $[0, 1] \times [0, 1]$  and the 4 basis functions

$$\phi_1(x, y) = (1 - x)(1 - y)$$

$$\phi_2(x, y) = x(1 - y)$$

$$\phi_3(x, y) = xy \text{ and}$$

$$\phi_4(x, y) = (1 - x)y.$$

- The element  $K_{ij}$  is then given by

$$K_{ij} = \int_0^1 \int_0^1 \left[ \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} + \frac{\partial \phi_i}{\partial y} \frac{\partial \phi_j}{\partial y} \right] dx dy$$

For instance,

$$K_{11} = \int_0^1 \int_0^1 \left[ (1 - y)^2 + (1 - x)^2 \right] dx dy = \frac{2}{3}.$$



$$K = \frac{1}{6} \begin{pmatrix} 4 & -1 & -2 & -1 \\ -1 & 4 & -1 & -2 \\ -2 & -1 & 4 & -1 \\ -1 & -2 & -1 & 4 \end{pmatrix}.$$

# Finite Element Method IV

## One Element Case - continued

### Linear basis functions

- Let us take the domain  $[0, 1] \times [0, 1]$  and the 4 basis functions

$$\phi_1(x, y) = (1 - x)(1 - y)$$

$$\phi_2(x, y) = x(1 - y)$$

$$\phi_3(x, y) = xy \text{ and}$$

$$\phi_4(x, y) = (1 - x)y.$$

- The element  $K_{ij}$  is then given by

$$K_{ij} = \int_0^1 \int_0^1 \left[ \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} + \frac{\partial \phi_i}{\partial y} \frac{\partial \phi_j}{\partial y} \right] dx dy$$

For instance,

$$K_{11} = \int_0^1 \int_0^1 \left[ (1 - y)^2 + (1 - x)^2 \right] dx dy = \frac{2}{3}.$$

$$K = \frac{1}{6} \begin{pmatrix} 4 & -1 & -2 & -1 \\ -1 & 4 & -1 & -2 \\ -2 & -1 & 4 & -1 \\ -1 & -2 & -1 & 4 \end{pmatrix}.$$

# Finite Element Method IV

## One Element Case - continued

### Linear basis functions

- Let us take the domain  $[0, 1] \times [0, 1]$  and the 4 basis functions

$$\phi_1(x, y) = (1 - x)(1 - y)$$

$$\phi_2(x, y) = x(1 - y)$$

$$\phi_3(x, y) = xy \text{ and}$$

$$\phi_4(x, y) = (1 - x)y.$$

- The element  $K_{ij}$  is then given by

$$K_{ij} = \int_0^1 \int_0^1 \left[ \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} + \frac{\partial \phi_i}{\partial y} \frac{\partial \phi_j}{\partial y} \right] dx dy$$

For instance,

$$K_{11} = \int_0^1 \int_0^1 \left[ (1 - y)^2 + (1 - x)^2 \right] dx dy = \frac{2}{3}.$$



$$K = \frac{1}{6} \begin{pmatrix} 4 & -1 & -2 & -1 \\ -1 & 4 & -1 & -2 \\ -2 & -1 & 4 & -1 \\ -1 & -2 & -1 & 4 \end{pmatrix}.$$

# Finite Element Method V

## Multiple elements

- Show that if we evaluate the basis functions on a domain  $[0, h] \times [0, h]$  we get the same element matrix.
- Consider the  $3 \times 3$  mesh on the right.
- Choose an element numbering, see Fig [blue numbers](#).
- Choose a global node numbering, see Fig [red numbers](#)
- Set up a table/matrix which relates local node numbers to global node numbers  $GM(e, LN)$

### Node numbering

# Finite Element Method V

## Multiple elements

- Show that if we evaluate the basis functions on a domain  $[0, h] \times [0, h]$  we get the same element matrix.
- Consider the  $3 \times 3$  mesh on the right.
- Choose an element numbering, see Fig [blue numbers](#).
- Choose a global node numbering, see Fig [red numbers](#)
- Set up a table/matrix which relates local node numbers to global node numbers  $GM(e, LN)$


### Node numbering

# Finite Element Method V

## Multiple elements

- Show that if we evaluate the basis functions on a domain  $[0, h] \times [0, h]$  we get the same element matrix.
- Consider the  $3 \times 3$  mesh on the right.
- Choose an element numbering, see Fig [blue numbers](#).
- Choose a global node numbering, see Fig [red numbers](#)
- Set up a table/matrix which relates local node numbers to global node numbers  $GM(e, LN)$

### Node numbering



# Finite Element Method V

## Multiple elements

- Show that if we evaluate the basis functions on a domain  $[0, h] \times [0, h]$  we get the same element matrix.
- Consider the  $3 \times 3$  mesh on the right.
- Choose an element numbering, see Fig **blue numbers**.
- Choose a global node numbering, see Fig **red numbers**
- Set up a table/matrix which relates local node numbers to global node numbers  $GM(e, LN)$

### Node numbering

1	2	3
1	2	3
1	2	3



# Finite Element Method V

## Multiple elements

- Show that if we evaluate the basis functions on a domain  $[0, h] \times [0, h]$  we get the same element matrix.
- Consider the  $3 \times 3$  mesh on the right.
- Choose an element numbering, see Fig **blue numbers**.
- Choose a global node numbering, see Fig **red numbers**
- Set up a table/matrix which relates local node numbers to global node numbers  $GM(e, LN)$

### Node numbering

13	14	15	16
7	8	9	
9	10	11	12
4	5	6	
5	6	7	8
1	2	3	
1	2	3	4

# Finite Element Method V


## Multiple elements

- Show that if we evaluate the basis functions on a domain  $[0, h] \times [0, h]$  we get the same element matrix.
- Consider the  $3 \times 3$  mesh on the right.
- Choose an element numbering, see Fig **blue numbers**.
- Choose a global node numbering, see Fig **red numbers**
- Set up a table/matrix which relates local node numbers to global node numbers  $GM(e, LN)$

### Gathering matrix

GM	1	2	3	4	5	6	7	8	9
1	1	2	3	5	6	7	9	10	11
2	2	3	4	6	7	8	10	11	12
3	6	7	8	10	11	12	14	15	16
4	5	6	7	9	10	11	13	14	15

### Node numbering



# Finite Element Method VI

## Multiple elements - continued

Gathering matrix

GM	1	2	3	4	5	6	7	8	9
1	1	2	3	5	6	7	9	10	11
2	2	3	4	6	7	8	10	11	12
3	6	7	8	10	11	12	14	15	16
4	5	6	7	9	10	11	13	14	15

## Global matrix structure

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 

```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

# Finite Element Method VI

## Multiple elements - continued

Gathering matrix

GM	1	2	3	4	5	6	7	8	9
1	1	2	3	5	6	7	9	10	11
2	2	3	4	6	7	8	10	11	12
3	6	7	8	10	11	12	14	15	16
4	5	6	7	9	10	11	13	14	15

## Global matrix structure

- The global matrix is now constructed from the element matrices and gathering matrix by
- ```

A = 0
for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
      + Ke(i, j)
    end for
  end for
end for
  
```

# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

### Global matrix structure

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 

```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 


```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

### Global matrix structure

Contribution of element 1 to matrix



# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 


```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

### Global matrix structure

Contribution of element 2 to matrix



# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 


```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

### Global matrix structure

Contribution of element 3 to matrix





# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 


```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

### Global matrix structure

Contribution of element 4 to matrix



# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 


```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

### Global matrix structure

Contribution of element 5 to matrix



# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 


```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

### Global matrix structure

Contribution of element 6 to matrix



# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 


```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

### Global matrix structure

Contribution of element 7 to matrix



# Finite Element Method VI

## Multiple elements - continued


### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 
  - for  $e = 1$  to NoE
  - for  $i = 1$  to 4
  - for  $j = 1$  to 4
  - $$A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j)) + K^e(i, j)$$
  - end for
  - end for
  - end for

### Global matrix structure

Contribution of element 8 to matrix



# Finite Element Method VI

## Multiple elements - continued


### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 
  - for  $e = 1$  to NoE
  - for  $i = 1$  to 4
  - for  $j = 1$  to 4
  - $$A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j)) + K^e(i, j)$$
  - end for
  - end for
  - end for

### Global matrix structure

Contribution of element 9 to matrix



# Finite Element Method VI

## Multiple elements - continued

### Gathering matrix

| GM | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|----|---|---|---|----|----|----|----|----|----|
| 1  | 1 | 2 | 3 | 5  | 6  | 7  | 9  | 10 | 11 |
| 2  | 2 | 3 | 4 | 6  | 7  | 8  | 10 | 11 | 12 |
| 3  | 6 | 7 | 8 | 10 | 11 | 12 | 14 | 15 | 16 |
| 4  | 5 | 6 | 7 | 9  | 10 | 11 | 13 | 14 | 15 |

- The global matrix is now constructed from the element matrices and gathering matrix by
- $A = 0$ 


```

for e = 1 to NoE
  for i = 1 to 4
    for j = 1 to 4
      A(GM(e, i), GM(e, j)) = A(GM(e, i), GM(e, j))
        + Ke(i, j)
    end for
  end for
end for

```

### Global matrix structure

Sparsity pattern FEM Matrix



# Summary

## What you should have learned

### Basic idea

- The three basic methods – FDM, FVM and FEM – to discretize an elliptic problem on a square domain
- How a mapping allows you to convert problems on a non-square domain to a square domain.
- That a mixed formulation should be discretized on a staggered grid
- That all methods lead to a symmetric linear, algebraic system.



# Summary

## What you should have learned

### Basic idea

- The three basic methods – **FDM, FVM and FEM** – to discretize an elliptic problem on a square domain
- How a **mapping** allows you to convert problems on a non-square domain to a square domain.
- That a mixed formulation should be discretized on a **staggered grid**
- That all methods lead to a symmetric linear, algebraic system.

# Summary

## What you should have learned

### Basic idea

- The three basic methods – **FDM, FVM and FEM** – to discretize an elliptic problem on a square domain
- How a **mapping** allows you to convert problems on a non-square domain to a square domain.
- That a mixed formulation should be discretized on a **staggered grid**
- That all methods lead to a symmetric linear, algebraic system.

# Summary

## What you should have learned

### Basic idea

- The three basic methods – **FDM, FVM and FEM** – to discretize an elliptic problem on a square domain
- How a **mapping** allows you to convert problems on a non-square domain to a square domain.
- That a mixed formulation should be discretized on a **staggered grid**
- That all methods lead to a symmetric linear, algebraic system.

# Summary

## What you should have learned

### Basic idea

- The three basic methods – **FDM, FVM and FEM** – to discretize an elliptic problem on a square domain
- How a **mapping** allows you to convert problems on a non-square domain to a square domain.
- That a mixed formulation should be discretized on a **staggered grid**
- That all methods lead to a symmetric linear, algebraic system.