

Uncountability

Vast is still finite. Now we count things that aren't — and discover that some infinities are bigger than others.

[← ICS 06 · The Machine](#)

MEANING

What we want to talk about is **not** countable.

f1	1	1	0	1	1	0	0	0
f2	1	1	1	1	0	1	1	1
f3	0	1	1	1	0	1	1	1
f4	1	1	0	1	1	0	0	1
f5	0	1	1	0	0	0	0	1
f6	1	1	1	0	1	1	0	0
f7	0	1	1	0	1	1	1	0
f8	0	1	1	1	0	1	1	1
flip	0	0	0	0	1	0	0	0

differs from every row – so it is on no row

the diagonal

Consider all infinite sequences of bits:
0110100011... forever. Each one is a complete answer to an endless list of yes/no questions — a behaviour, a function, a real number, a fate.

Claim: no list can contain them all. Not a long list. Not an infinite list. No list.

Cantor's proof is three lines and it is the most consequential argument of the 20th century.
We run it twice, slowly.

THE OBJECTS

A subset of the numbers is one yes-or-no answer per number.

	1	2	3	4	5	6	7	8	...
{ }	0	0	0	0	0	0	0	0	...
all	1	1	1	1	1	1	1	1	...
evens	0	1	0	1	0	1	0	1	...
primes	0	1	1	0	1	0	1	0	...
{3}	0	0	1	0	0	0	0	0	...
no name	1	1	0	1	0	1	1	0	...

and almost every subset is this one

subsets as answer sheets · 1 = in

Take any collection of counting numbers — the evens, the primes, just {3}, none, all. Each is a subset. To pin one down, walk 1, 2, 3, ... and answer in or out, forever.

That answer sheet is an infinite bit string, and every infinite bit string is one. Together they make the power set, $2^{\mathbb{N}}$ — two choices, made $\mathbb{N}$ times.

A handful have names. Almost all are an endless coin-flip, with nothing shorter to say than the flips.

Hand me a list of every subset.
I will build one that is **not on it**.

	1	2	3	4	5	6
S1						
S2						
S3						
S4						
S5						
S6						
D						

D disagrees with row n about n

 in ·  out

Suppose the subsets could be listed: S1, S2, S3, ..., every one somewhere on the list.

Go down the **diagonal** and ask each row about *its own number*: is 1 in S1? Is 2 in S2? Is n in Sn?

Build D by answering the opposite every time: $D = \{ n : n \text{ is not in } S_n \}$.

D is not S1: they disagree about 1. Not S2: they disagree about 2. Not any Sn. So the list left something out, and it was any list at all.

The same move, on the numbers between 0 and 1.

r1 = 0.	1	4	1	5	9	2	...
r2 = 0.	7	1	8	2	8	1	...
r3 = 0.	2	5	5	5	5	5	...
r4 = 0.	3	3	3	3	3	3	...
r5 = 0.	0	0	0	0	5	0	...
r6 = 0.	4	1	4	2	1	3	...
x = 0.	5	5	4	5	4	5	...

x differs from row n in place n

any list · and the number it missed

Suppose you could list them: r1, r2, r3, ..., each an endless decimal. Take the first digit of the first, the second of the second, the n-th of the n-th. The diagonal again.


Build a new number x by changing every one of those digits. Then x differs from rn in the n-th place — so x is on no row.

THE ONE PLACE IT NEEDS CARE

Change each digit to 5, or to 4 if it was 5. That avoids the trailing nines: 0.4999... and 0.5000... are the same number written twice, and a proof that landed on that would prove nothing.

INTUITION

Between any two numbers, however close, there is **another**. But do not let it do the work.



and again, forever

halve, and halve again

Pick two reals as close together as you like. Their midpoint lies strictly between them. Do it again, forever. There is no *next* real number, and a list is nothing but firsts and nexts.

Density is the feeling, not the reason. The fractions are dense in the same way — and they can still be listed, by walking a grid of numerators and denominators corner by corner. Week 5's exercise.

Only the diagonal tells the two cases apart. When an intuition and a proof agree, check which of them you are actually relying on.

THEOREM

The diagonal, **once and for all**.


CANTOR · 1891

For every set S , the set of all subsets of S is strictly larger than S . In particular, the real numbers cannot be listed.

Where self-reference enters. The proof builds an object from the list that asks of each row: "do you contain yourself here?" — and then answers the opposite. The list is used against itself.

You have just seen it twice. Power set, real numbers, bit sequences: three costumes, one cardinality. Russell turned the same trick on set theory in 1901: the set of all sets that do not contain themselves.

There are incomparably more meanings than notations.



Programs: countable. **Behaviours:** uncountable. So almost every behaviour has no program.

Proofs: countable. **Truths** about numbers: uncountable. So almost every truth has no proof.

Sentences: countable. **Distinctions** the world admits: uncountable. So almost everything is unsaid.

"Almost every" here is exact: the expressible is a vanishing sliver, of measure zero, inside the meaningful.

Scarcity of notation is a job description.

WHAT IT FORBIDS

A final language. No vocabulary — mathematical, legal, musical, or neural — will ever cover the space of meanings.

WHAT IT OPENS

An inexhaustible supply of things worth naming. Every new notation captures meaning that was previously unreachable — and there is always more left. Calculus, double-entry bookkeeping, staff notation, the periodic table, DNA sequencing, type systems: each was a raid on the uncountable.

Chaitin's version: almost all numbers are random, that is, incompressible, that is, nameless. Not a wall — unclaimed territory, and claiming it is what every discipline calls progress. This is the first time the title of the class shows through.

Recommended exercises.

- 1 Read the Meaning chapter from *What We Want to Talk About Is Not to Syntax and Semantics*.
- 2 Write six subsets of $\{1, \dots, 6\}$ as answer sheets and build the diagonal D by hand. Check it against every row.
- 3 Run the decimal version on six decimals of your own. Why do the fractions escape the argument?
- 4 How many programs of at most 1,000 characters are there over 100 symbols? How many behaviours on inputs 1 to 1,000? Which is bigger, and by how much?
- 5 Find one new notation from your own life — a recipe format, chess notation, a knitting pattern — and say what it made sayable.

NEXT WEEK

Self-reference I: a compiler defines the meaning of the language it is written in; the fixed point, and what Ken Thompson says it cannot prove; and Gödel's two theorems, with the same diagonal.

Listen, then read.

LISTEN

Bruckner · Symphony No. 7 — Karajan, Vienna Philharmonic, 1989. The Adagio, written as Wagner lay dying: endlessness that comes in sizes. Bruckner's time does not tick, it expands, and the climax arrives the way the diagonal does — from outside every row.

READ

Everything and More: A Compact History of ∞ — Wallace, 2003: the infinite by a novelist who did the mathematics. And, more technical, Cantor's diagonal argument — 1891, four pages; the article has the construction and the reference to the original.