

# Class 11 Lab Session

Chloe Wong (PID: A16893383)

2024-11-07

## Section 1. Proportion of G/G in a population

Downloaded a CSV file from Ensembl < [https://useast.ensembl.org/Homo\\_sapiens/Variation/Sample?db=core;r=17:39894595-39895595;v=rs8067378;vdb=variation;vf=959672880#373531\\_tablePanel](https://useast.ensembl.org/Homo_sapiens/Variation/Sample?db=core;r=17:39894595-39895595;v=rs8067378;vdb=variation;vf=959672880#373531_tablePanel) >

Here we read this CSV file

```
mxl <- read.csv("373531-SampleGenotypes-Homo_sapiens_Variation_Sample_rs8067378.csv")
head(mxl)
```

```
## Sample..Male.Female.Unknown. Genotype..forward.strand. Population.s. Father
## 1 NA19648 (F) A|A ALL, AMR, MXL -
## 2 NA19649 (M) G|G ALL, AMR, MXL -
## 3 NA19651 (F) A|A ALL, AMR, MXL -
## 4 NA19652 (M) G|G ALL, AMR, MXL -
## 5 NA19654 (F) G|G ALL, AMR, MXL -
## 6 NA19655 (M) A|G ALL, AMR, MXL -
## Mother
## 1 -
## 2 -
## 3 -
## 4 -
## 5 -
## 6 -
```

```
table(mxl$Genotype..forward.strand.)
```

```
##
## A|A A|G G|A G|G
## 22 21 12 9
```

```
table(mxl$Genotype..forward.strand.) / nrow(mxl) * 100
```

```
##
## A|A A|G G|A G|G
## 34.3750 32.8125 18.7500 14.0625
```

## Section 4: Population Scale Analysis

One sample is obviously not enough to know what is happening in a population. You are interested in assessing genetic differences on a population scale.

How many samples do we have?

```
expr <- read.table("rs8067378_ENSG00000172057.6.txt")
head(expr)
```

```
##      sample geno      exp
## 1 HG00367   A/G 28.96038
## 2 NA20768   A/G 20.24449
## 3 HG00361   A/A 31.32628
## 4 HG00135   A/A 34.11169
## 5 NA18870   G/G 18.25141
## 6 NA11993   A/A 32.89721
```

```
nrow(expr)
```

```
## [1] 462
```


```
table(expr$geno)
```

```
##
## A/A A/G G/G
## 108 233 121
```

```
library(ggplot2)
```

Lets make a boxplot

```
plot <- ggplot(expr) + aes(x=geno, exp, fill=geno) +
  geom_boxplot(notch=TRUE)
plot
```



Q13: Read this file into R and determine the sample size for each genotype and their corresponding median expression levels for each of these genotypes.

(Code above for sample sizes)

The sample size for A/A is 108. The sample size for A/G is 233. The sample size for G/G is 121. The median expression for A/A is 30.779. The median for A/G is 25.640. The median for G/G is 20.004.

```
summary(expr)
```

```
##      sample          geno          exp
## Length:462      Length:462      Min.   : 6.675
## Class :character Class :character 1st Qu.:20.004
## Mode  :character Mode  :character Median :25.116
##                                     Mean  :25.640
##                                     3rd Qu.:30.779
##                                     Max.   :51.518
```

Q14: Generate a boxplot with a box per genotype, what could you infer from the relative expression value between A/A and G/G displayed in this plot? Does the SNP effect the expression of ORMDL3?

In the boxplot, the expression value of A/A is significantly higher than the expression value of G/G. The SNP affects the expression of ORMDL3 as the expression is seen to be approximately 10% greater in A/A compared to G/G.