

Physical Examination in Sleep Medicine

Kryger Chapter 59



Overview

- **Certain sleep disorders can be attributed to physiological factors**
- **Physicians perform a physical assessment of the patient to determine if these physiological factors contribute or cause the sleep disorder**
- **Sleep disorders with physiological components:**
 - **Sleep Apnea**
 - **Narcolepsy**
 - **Restless Leg Syndrome**
 - **Parasomnias**


Sleep Apnea

- Has multiple anatomic risk factors
 - Obesity and increased neck circumference are main ones
 - Other contributing factors for OSA:
 - Reduced oropharyngeal airspace



Sleep Apnea

- **Retrognathia:** Condition in which either or both jaws recede with respect to the frontal plane of the forehead
- **Micrognathia:** Unusual smallness of the jaws, especially the lower jaw



Sleep Apnea


- **CSA contributing factors:**
 - **Impaired respiratory effort**
 - **Heart failure**
 - **CNS disease**
 - **Neuromuscular disease**

Sleep Apnea



- **Overall inspection:**
 - **Obesity significant risk factor**
 - **Increases OSA likelihood 10-fold, particularly central type of obesity**
 - **May subject patients to risk of worsening obesity due to OSA effects, such as sleep deprivation, hypersomnia, and disrupted metabolism**


Sleep Apnea



- Also associated with hypothyroidism and acromegaly (abnormal enlargement of limbs)
- Oropharyngeal airway myopathy, edema, and obesity contribute to upper airway collapse and obstruction

Sleep Apnea

- Goiter another contributing factor to OSA



Sleep Apnea

- Polycystic ovarian syndrome
 - Associated with metabolic syndrome
 - Greater risk for:
 - Obesity
 - Impaired glucose tolerance
 - Type 2 diabetes
 - Cardiovascular disease
 - OSA

Sleep Apnea

- **Craniofacial features:**
 - **Usually there are changes to soft palate and uvula, to volume and position of the tongue, and jaw position in OSA patients**
 - **If tongue cannot lay flat due to jaw position, more likely to block airway during sleep**
 - **Down syndrome patients more likely to have OSA due to craniofacial features**



Sleep Apnea

- Retrognathia and micrognathia
 - Crowded teeth
 - Scalloped tongue

Sleep Apnea

- Nasal factors:
 - Look at anatomic abnormalities that may contribute to nasal obstruction
 - Congenital
 - Traumatic
 - Infectious
 - Neoplastic



Sleep Apnea



- **Neck Circumference:**
 - Increased neck circumference risk factor for OSA
 - Greater than 48 cm (19.2 in) has 20-fold increased risk for OSA

Sleep Apnea

- Examination of the Pharynx
 - Two classification systems to determine relation of tongue to pharynx
 - Mallampati
 - Four classes:
 - Class I: Soft palate, fauces, uvula, and posterior and anterior pillars are visible
 - Class II: Soft palate, fauces, and uvula visible
 - Class III: Soft palate, fauces, and only base of uvula visible
 - Class IV: Soft palate not visible


Sleep Apnea

- Mallampati



Sleep Apnea

- Friedman



Sleep Apnea

- Examination of the Tonsils
 - Enlarged tonsils and adenoids are major cause of airway obstruction and OSA in children
 - Size graded on a scale of 1 to 4

Sleep Apnea

- Examination of the Tonsils



0

Surgically removed tonsils



1

Tonsils hidden within tonsil pillars



2

Tonsils extending to the pillars



3

Tonsils are beyond the pillars



4

Tonsils extend to midline

Sleep Apnea

- **Neurologic Examination**
 - **Neuromuscular disease may cause OSA**
 - **Muscular dystrophies**
 - **Muscle atrophy**
 - **Multiple sclerosis**
- **Cardiopulmonary Examination**
 - **CHF = High likelihood for CSA**
 - **Peripheral edema a common finding in obesity hypoventilation syndrome and some OSA patients**
 - **COPD and asthma also seen in association with OSA**

Narcolepsy

- In general, patients tend to be obese, with increased predisposition to type 2 diabetes and lower basal metabolism
- Seen in conditions such as:
 - CNS tumors
 - Head trauma
 - Multiple sclerosis
 - Encephalitis
 - Neurodegeneration

Restless Leg Syndrome

- Prevalence in Type 2 diabetes is 17.7%
 - May be higher in those with hereditary neuropathy
- Occurs in 1/3 of those with polyneuropathy
- Reduced iron stores can cause RLS
- Neurologic examination reveals:
 - Sensory loss
 - Paresthesias (tingling sensation) and dyesthesias (distortion of any sense, especially touch)
- Pain may be a significant symptom


Parasomnias

- Sleep-Related Eating Disorder
 - BMI usually abnormally high in 60% of patients
- RBD
 - Frequently at risk for Parkinson's
 - Majority of patients present with hyposmia (impaired smell)

Bruxism

- Grinding of teeth
- Can lead to abnormal wear of teeth, periodontal tissue damage, or jaw pain
- Other symptoms include facial muscle and tooth pain and headache
- Left untreated, leads to things like recession and inflammation of gums and TMJ

Bruxism





Insomnia

- **Seen often with various disorders:**
 - **Endocrine**
 - **Mood**
 - **Anxiety**
 - **Rheumatologic**
 - **Pain**
 - **Graves disease (autoimmune disorder affecting thyroid)**