

A BEALE–KATO–MAJDA CRITERION FOR RIGHT-INVARIANT H^s METRICS ON $\text{SDiff}(\mathbb{T}^3)$

ADILKHAN SALKIMBAYEV

ABSTRACT. Fix the bracket of $\text{SDiff}(\mathbb{T}^3)$ and let the inertia operator range over the isotropic Fourier multipliers $a(|k|) = |k|^{2s}$, so that $s = 0$ is the incompressible Euler equation. We prove a continuation criterion for $s > 0$: with m the momentum and $w = \text{curl } m$, a solution persists as long as $\int_0^T \|w\|_{L^p} dt < \infty$ for one exponent $p > 3/(2s)$. Two ingredients. The first is a reduction: w obeys the *unmodified* vortex-stretching equation for every s , so that all the s -dependence sits in the recovery of the velocity from w , a Fourier multiplier of order $-1 - 2s$. The second is that for $s > 0$ the resulting generalised Biot–Savart law is smoothing, its kernel locally integrable, and $\|\nabla u\|_{L^\infty} \lesssim \|w\|_{L^p}$ with no logarithm. The classical Beale–Kato–Majda criterion is the endpoint $s \rightarrow 0^+$, where the admissible p escapes to infinity and the logarithm of Brezis–Gallouët–Wainger becomes unavoidable. Tao observed that no such criterion was available for generalised Euler equations and did not attempt one. We also record what the criterion does not do: the conserved energy is $\|w\|_{\dot{H}^{-1-s}}^2$, which dominates $\|\nabla u\|_{L^\infty}$ exactly when $s \geq 5/2$, recovering the geodesic completeness threshold of Bruveris–Vialard from the analytic side and confirming that the criterion alone moves nothing.

1. INTRODUCTION

Arnold’s theorem realises the incompressible Euler equation as the geodesic flow of the right-invariant L^2 metric on the group of volume-preserving diffeomorphisms [1], in the rigorous form given by Ebin and Marsden [14]. Fixing that group and varying the metric gives a one-parameter family

$$a(|k|) = |k|^{2s}, \quad s \in \mathbb{R}, \tag{1}$$

of right-invariant metrics on $\text{SDiff}(\mathbb{T}^3)$, with Euler at $s = 0$. The family is the three-dimensional, volume-preserving analogue of the H^s metrics on $\text{Diff}(S^1)$, where the behaviour is understood at both ends: shocks at $s = 0$, wave breaking at $s = 1$, and geodesic completeness for $s \geq 3/2$ [16, 3]. In three dimensions only the smoothing end is settled. For $s > \frac{1}{2} \dim M + 1$, that is $s > 5/2$, the geodesic flow of a strong right-invariant H^s metric is complete, on the volume-preserving subgroup as well as on the full group [9, Rems. 7.3 and 7.5, Cor. 7.6]; below $5/2$ nothing is known in either direction, and $s = 0$ is the Euler problem itself. How far that threshold can be displaced is asked in [25, Question 7.2], which this note was written to make approachable and does not answer.

Any attempt to move that threshold has to begin with a continuation criterion: a statement that a solution persists for as long as some quantity remains integrable in time. For $s = 0$ this is the theorem of Beale, Kato and Majda [4], which replaces $\|\nabla u\|_{L^\infty}$ by $\|\omega\|_{L^\infty}$ at the cost of a logarithm. For the family (1) no such criterion appears to have been written. Tao, considering the wider class of generalised Euler equations in which the vector potential operator is an arbitrary operator of order -2 , records the gap explicitly: analogues of at least some of the known refinements should be establishable, the generalised Biot–Savart law obeying estimates

very similar to the true one, but he does not attempt it [28, Rem. 1.10]. This note supplies the criterion for (1) with $s > 0$.

What is proved. Four things.

The reduction (Proposition 2.2). The field $w := \operatorname{curl} m$, m the momentum, satisfies

$$\partial_t w + (u \cdot \nabla) w = (w \cdot \nabla) u, \quad \operatorname{div} w = 0,$$

the vortex-stretching equation of the *unmodified* Euler equation, for every s . It is an identity of vector calculus valid for any m whatever and any divergence-free u ; it does not use $m = Au$. So the whole s -dependence is carried by the constitutive law $w \mapsto u$, which by Lemma 2.4 is a Fourier multiplier of order $-1 - 2s$, whence $\nabla u = T_s w$ with T_s of order $-2s$.

The kernel estimate (Lemma 3.1). For $s > 0$ the operator T_s is smoothing, its convolution kernel is bounded by $|x|^{2s-3}$ and hence lies in $L^{p'}$ exactly when $p > 3/(2s)$, so that $\|\nabla u\|_{L^\infty} \lesssim_{s,p} \|w\|_{L^p}$ with no logarithm.

The a priori estimate (Lemma 4.1, Proposition 4.3 and Theorem 6.2). For every $s > 0$ and every $r > \frac{5}{2}$, with no threshold in between,

$$\frac{d}{dt} \|w\|_{H^r} \lesssim (\|\nabla u\|_{L^\infty} + \|w\|_{L^\infty}) \|w\|_{H^r},$$

both terms being controlled by $\|\nabla u\|_{L^\infty}$ alone, so that finiteness of $\int_0^T \|\nabla u\|_{L^\infty}$ forbids breakdown at T .

The local theory (Section 5). Existence, uniqueness and the blowup alternative, none of them assumed. Combining the four gives the criterion, Theorem 6.3.

The whole degeneracy sits in p . The admissible exponent $p > 3/(2s)$ escapes to ∞ as $s \rightarrow 0^+$, and at $s = 0$ no finite p repairs the estimate: one must buy it back with the logarithm of Brezis–Gallouët–Wainger, so that the classical criterion is the endpoint of this family rather than an instance of it. The a priori estimate, by contrast, is uniform in $s \geq 0$ and needs only $r > \frac{5}{2}$.

What is not proved, and what this does not do. Nothing about the local theory is assumed. Existence is proved by Galerkin truncation (Theorem 5.3), uniqueness by an L^2 argument (Proposition 4.5), and the blowup alternative follows (Proposition 6.1). Solutions are taken in the class $\mathcal{X}_T^r$ of Definition 5.1, bounded in H^r and continuous in every $H^{r'}$ with $r' < r$, which is what the construction delivers and what every estimate here uses; such solutions lie in H^r at every time and are weakly continuous into it (Lemma 5.2). They are also strongly continuous into H^r , by the argument of Bona and Smith, which we omit as unused rather than unavailable. The commutator and product estimates of Kato–Ponce and Moser type are quoted, not proved, as are the Bernstein inequalities and the kernel asymptotics of [27]. Section 7 records the negative half: the conserved energy is $\|w\|_{\dot{H}^{-1-s}}^2$ and a two-line count shows that it dominates $\|\nabla u\|_{L^\infty}$ precisely when $s \geq 5/2$, so the criterion recovers the completeness threshold of [9] and moves it nowhere. That is the outcome to insist on: a criterion improving the threshold by itself, continued to $s = 0$, would prove global regularity for three-dimensional Euler.

Verification. The identities of Section 2 are checked spectrally on a fixed 24^3 grid with band-limited data, to relative errors between 10^{-16} and 10^{-18} , by `bkm_check.py`. The kernel claim underlying Lemma 3.1 is checked separately by `kernel_check.py`; see Remark 3.2. The remaining estimates, in Sections 4–6, are not numerical.

2. SETUP AND THE REDUCTION

Let $\mathbb{T}^3 = (\mathbb{R}/2\pi\mathbb{Z})^3$ and let $\mathfrak{g}$ be the space of smooth, divergence-free, mean-zero vector fields on $\mathbb{T}^3$, with the Jacobi–Lie bracket $[u, v] = (u \cdot \nabla)v - (v \cdot \nabla)u$ and the L^2 pairing. For $s \in \mathbb{R}$ let $A = A_s$ be the Fourier multiplier with symbol $a(|k|) = |k|^{2s}$, symmetric and positive definite on $\mathfrak{g}$ because the mean mode is excluded. Write $m = Au$ and let P be the Leray projection. All fields below are mean-zero, so that $\|\cdot\|_{H^\sigma}$ and $\|\cdot\|_{\dot{H}^\sigma}$ are equivalent, and we write $\Lambda = (1 - \Delta)^{1/2}$. The Euler–Arnold equation of the metric $\langle A\cdot, \cdot \rangle$ is

$$\dot{m} = \text{ad}_u^* m, \quad u = A^{-1}m. \quad (2)$$

Lemma 2.1. *For $u \in \mathfrak{g}$ and any vector field m ,*

$$\text{ad}_u^* m = -P[(u \cdot \nabla)m + (\nabla u)^\top m], \quad ((\nabla u)^\top m)_i = m_j \partial_i u^j.$$

Proof. By definition $\langle \text{ad}_u^* m, v \rangle = \langle m, [u, v] \rangle$ for $v \in \mathfrak{g}$, and

$$\langle m, [u, v] \rangle = \int m_i (u^j \partial_j v^i - v^j \partial_j u^i) = - \int (u^j \partial_j m_i) v^i - \int m_j (\partial_i u^j) v^i,$$

integrating the first term by parts and using $\text{div } u = 0$, and relabelling indices in the second. As this holds for all divergence-free v , the formula follows after applying P . $\square$

So (2) is the system

$$\partial_t m + (u \cdot \nabla)m + (\nabla u)^\top m = -\nabla p, \quad u = A^{-1}m, \quad \text{div } u = 0. \quad (3)$$

Proposition 2.2 (the reduction). *Let u be divergence-free and let m be any vector field. With $w = \text{curl } m$,*

$$\text{curl}[(u \cdot \nabla)m + (\nabla u)^\top m] = (u \cdot \nabla)w - (w \cdot \nabla)u.$$

Consequently a solution of (3) satisfies

$$\partial_t w + (u \cdot \nabla)w = (w \cdot \nabla)u, \quad \text{div } w = 0, \quad (4)$$

for every s : the vortex-stretching equation of the unmodified Euler equation.

Proof. Regard m as a 1-form; in Cartesian coordinates $(\mathcal{L}_u m)_i = u^j \partial_j m_i + m_j \partial_i u^j$, the bracketed expression. Since d commutes with $\mathcal{L}_u$ we have $\mathcal{L}_u(dm) = d(\mathcal{L}_u m)$. Identify the 2-form dm with the vector field $w = \text{curl } m$ by $dm = \iota_w \text{vol}$. Then

$$\mathcal{L}_u(\iota_w \text{vol}) = \iota_{[u, w]} \text{vol} + \iota_w \mathcal{L}_u \text{vol} = \iota_{[u, w]} \text{vol},$$

since $\mathcal{L}_u \text{vol} = (\text{div } u) \text{vol} = 0$, and $[u, w] = (u \cdot \nabla)w - (w \cdot \nabla)u$. Taking curl of (3) kills the gradient and gives (4). $\square$

Remark 2.3. The force of Proposition 2.2 is its quantifier: the identity holds for arbitrary m , so the transported quantity obeys Euler’s own equation whichever inertia operator is used, and nothing is lost or gained there. What varies with s is only the constitutive law recovering u from w . This is why the family is tractable by Euler’s own methods, and why the criterion below is a statement about a kernel rather than about a dynamics.

Lemma 2.4 (the orders). *On mean-zero divergence-free fields, $\hat{m}(k) = i|k|^{-2} k \times \hat{w}(k)$ and*

$$\hat{u}(k) = \frac{i}{|k|^{2+2s}} k \times \hat{w}(k), \quad \widehat{\partial_j u^i}(k) = -\frac{k_j (k \times \hat{w}(k))_i}{|k|^{2+2s}}.$$

Thus $u = S_s w$ and $\nabla u = T_s w$ for Fourier multipliers whose symbols are smooth away from the origin and homogeneous of degrees $-1 - 2s$ and $-2s$ respectively. In particular, for every σ ,

$$\|u\|_{H^\sigma} \lesssim \|w\|_{H^{\sigma-1-2s}}, \quad \|\nabla u\|_{H^\sigma} \lesssim \|w\|_{H^{\sigma-2s}}. \quad (5)$$

Proof. Since $\hat{m}(k) \perp k$, $k \times \hat{w} = ik \times (k \times \hat{m}) = i(k(k \cdot \hat{m}) - |k|^2 \hat{m}) = -i|k|^2 \hat{m}$, which inverts to the stated formula; then $\hat{u} = a(|k|)^{-1} \hat{m} = |k|^{-2s} \hat{m}$ and $\widehat{\partial_j u^i} = ik_j \hat{u}^i$. The symbol of T_s is a homogeneous polynomial of degree 2 over $|k|^{2+2s}$. $\square$

Lemma 2.5 (the conserved quantity). *For $u \in \mathfrak{g}$ and $w = \text{curl } Au$,*

$$E = \langle Au, u \rangle = \|u\|_{\dot{H}^s}^2 = \|w\|_{\dot{H}^{-1-s}}^2,$$

and E is conserved along (2).

Proof. $|\hat{w}(k)| = |k| |\hat{m}(k)|$ because $\hat{m}(k) \perp k$, and $\hat{m} = |k|^{2s} \hat{u}$, so $|k|^{-2-2s} |\hat{w}|^2 = |k|^{-2s} |\hat{m}|^2 = |k|^{2s} |\hat{u}|^2$; sum over $k \neq 0$. Conservation is the Lie–Poisson cancellation: $\frac{d}{dt} \langle m, A^{-1} m \rangle = 2 \langle A^{-1} m, \dot{m} \rangle = 2 \langle u, \text{ad}_u^* m \rangle = 2 \langle m, [u, u] \rangle = 0$. $\square$

3. THE KERNEL ESTIMATE

Lemma 3.1. *Let $s > 0$ and let T_s be as in Lemma 2.4. Let $p \in [1, \infty]$ satisfy $p > 3/(2s)$. Then for mean-zero $w \in L^p(\mathbb{T}^3)$,*

$$\|T_s w\|_{L^\infty} \leq C(s, p) \|w\|_{L^p}.$$

Proof. The symbol of T_s is smooth on $\mathbb{R}^3 \setminus \{0\}$ and homogeneous of degree $-2s$. The convolution kernel on $\mathbb{T}^3$ of such a multiplier agrees, modulo a function smooth on all of $\mathbb{T}^3$, with the periodisation of the kernel of the corresponding multiplier on $\mathbb{R}^3$, which for degree $-2s$ with $0 < 2s < 3$ is homogeneous of degree $2s - 3$ [27, Ch. III, V]. Hence the periodic kernel κ_s obeys $|\kappa_s(x)| \leq C|x|^{2s-3}$ for $0 < |x| \leq \pi$ and is bounded elsewhere. For $2s > 3$ the kernel is bounded and continuous, and for $2s = 3$ it is logarithmic.

For $0 < 2s < 3$ we have $\kappa_s \in L^{p'}(\mathbb{T}^3)$ if and only if $(3 - 2s)p' < 3$, that is $p' < 3/(3 - 2s)$, that is $p > 3/(2s)$; Young's inequality $\|\kappa_s * w\|_{L^\infty} \leq \|\kappa_s\|_{L^{p'}} \|w\|_{L^p}$ gives the claim. For $2s \geq 3$ the kernel lies in every L^q , $q < \infty$, so every $p > 1$ is admissible, and for $2s > 3$ boundedness admits $p = 1$. $\square$

Remark 3.2 (numerical check). The two quantitative claims in the proof are checked by `kernel_check.py`, for the model symbol $|k|^{-2s}$ and for $k_1 k_2 / |k|^{2+2s}$, which is the shape of an entry of T_s . The integrability threshold is confirmed sharply: for $s \in \{0.25, 0.5, 1, 1.25\}$ the discrete $L^{p'}$ norm of the kernel converges under grid refinement at $p' = 0.8p'_c$ and grows at $p' = 1.2p'_c$, with $p'_c = 3/(3 - 2s)$ in each case. The pointwise decay is checked by summing the Fourier series along a ray with a smooth cutoff at $|k| = K$; the fitted exponent approaches $2s - 3$ monotonically in K , the gap at $K = 64$ being 0.34, 0.15, 0.066, 0.027 and 0.006 for $s = 0.25, \dots, 1.25$. The slower convergence at small s is expected, the kernel being more singular there and needing more modes to resolve.

The range $p > 3/(2s)$ cannot be enlarged, and in particular the critical exponent itself is genuinely excluded rather than lost to the method.

Proposition 3.3 (sharpness of the range). *Let $0 < 2s < 3$ and let $1 \leq p \leq 3/(2s)$. Then there is no constant C with*

$$\|T_s w\|_{L^\infty} \leq C \|w\|_{L^p}$$

for all mean-zero divergence-free $w \in C^\infty(\mathbb{T}^3)$. Consequently the hypothesis of Theorem 6.3 fails to be implied by boundedness of $\|w\|_{L^p}$ for any $p \leq 3/(2s)$, and the admissible range of that theorem is optimal within the Lebesgue scale.

Proof. Identify the ball $\{|x| < 1\}$ with a subset of $\mathbb{T}^3$ and let $\mathcal{A} := \{1 < |x| < 2\}$.

Step 1: a test field with non-vanishing response at the origin. We claim there is a divergence-free $\varphi \in C_c^\infty(\mathcal{A}; \mathbb{R}^3)$ and indices i, j with $(T_s \varphi)_{ij}(0) \neq 0$, where T_s is taken on $\mathbb{R}^3$. If not, then $\int \kappa_s^{ij,b}(-y) \varphi^b(y) dy = 0$ for all such φ and all i, j , where κ_s is the kernel of T_s on $\mathbb{R}^3$. Divergence-free compactly supported fields in $\mathcal{A}$ annihilate exactly the gradients, so this says $P \kappa_s^{ij}(-\cdot) = 0$ on $\mathcal{A}$; as κ_s is homogeneous it then vanishes on $\mathbb{R}^3 \setminus \{0\}$, so $T_s \circ P = 0$, which is false because the symbol of T_s is injective on $k^\perp$. Fix such a φ and write $\gamma := (T_s \varphi)_{ij}(0) \neq 0$.

Step 2: the construction. For $n \geq 1$ put $\varphi_n(x) := \varphi(2^n x)$, supported in the annulus $\{2^{-n} < |x| < 2^{-n+1}\}$; these are pairwise disjoint and contained in $\{|x| < 1\}$. Each φ_n is divergence-free and has mean zero, being a curl. Since κ_s is homogeneous of degree $2s - 3$,

$$(\kappa_s * \varphi_n)(0) = 2^{-2sn} (\kappa_s * \varphi)(0), \quad \text{so} \quad (\kappa_s * \varphi_n)_{ij}(0) = 2^{-2sn} \gamma.$$

Assume first $p > 1$; the interval $(1, 3/(2s)]$ is non-empty because $2s < 3$, and the case $p = 1$ follows from any member of it, since $\|w\|_{L^1} \leq |\mathbb{T}^3|^{1/p'} \|w\|_{L^p}$ would propagate an L^1 bound to every larger exponent. Fix β with $1/p < \beta \leq 1$, possible as $p > 1$, and set $c_n := 2^{3n/p} n^{-\beta}$ and

$$w_N := \sum_{n=1}^N c_n \varphi_n.$$

Step 3: the two sides. The supports are disjoint and $\|\varphi_n\|_{L^p}^p = 2^{-3n} \|\varphi\|_{L^p}^p$, so

$$\|w_N\|_{L^p}^p = \|\varphi\|_{L^p}^p \sum_{n=1}^N c_n^p 2^{-3n} = \|\varphi\|_{L^p}^p \sum_{n=1}^N n^{-\beta p} \leq \|\varphi\|_{L^p}^p \sum_{n \geq 1} n^{-\beta p} < \infty,$$

uniformly in N , since $\beta p > 1$. On the other side, by Lemma 3.1's decomposition the periodic kernel is $\kappa_s + h$ with h bounded on $\mathbb{T}^3$, so

$$|(T_s w_N)_{ij}(0)| \geq |\gamma| \sum_{n=1}^N c_n 2^{-2sn} - \|h\|_{L^\infty} \|w_N\|_{L^1} \geq |\gamma| \sum_{n=1}^N n^{-\beta} 2^{n(3/p-2s)} - C',$$

with C' independent of N because $\|w_N\|_{L^1} \lesssim \|w_N\|_{L^p}$ on a finite measure space. All terms of the sum are non-negative, so no cancellation can occur. If $p < 3/(2s)$ the exponent $3/p - 2s$ is positive and the sum grows geometrically; if $p = 3/(2s)$ it is zero and the sum is $\sum_{n \leq N} n^{-\beta}$, divergent because $\beta \leq 1$. In both cases $\|T_s w_N\|_{L^\infty} \rightarrow \infty$ while $\|w_N\|_{L^p}$ stays bounded. $\square$

Remark 3.4 (numerical check). Step 1 above is non-constructive, and it is the only ingredient of the proposition that is neither quoted nor arithmetic. `sharp_check.py` exhibits a φ and computes γ for the explicit annulus bump $\text{curl}[\chi(|x|)(1 + \frac{2}{5}x_1, \frac{7}{10} - \frac{3}{10}x_2, \frac{1}{5} + \frac{1}{2}x_3)]$, χ the standard bump on $(1, 2)$: the largest entry of $(T_s \varphi)(0)$ is $-4.740 \cdot 10^{-4}$, $-7.609 \cdot 10^{-4}$, $-8.325 \cdot 10^{-4}$ and $-7.501 \cdot 10^{-4}$ at $s = \frac{1}{4}, \frac{1}{2}, 1, \frac{5}{4}$, stable to six or seven significant figures across grids 96^3 to 192^3 , so it is a number and not a discretisation artefact. The two sums of Step 3 are tabulated there as well. The homogeneity of κ_s used in Step 2 is the content of Remark 3.2.

Remark 3.5 (the range is exactly the subcritical one). Proposition 3.3 and Lemma 3.1 together say that T_s is bounded from L^p to L^∞ precisely when $p > 3/(2s)$. By the dilation count of Section 7 the norm $\|w\|_{L^p}$ is subcritical for (4) precisely when $p > d/(2s)$, which in three dimensions is the same condition. So the criterion is available in exactly the scale-subcritical Lebesgue norms and in no others, the critical exponent $p = 3/(2s)$ included in the failure. That the two ranges coincide is not a coincidence — both are the statement that $\|\nabla u\|_{L^\infty} \sim \lambda^{a-2s}$ is dominated by $\|w\|_{L^p} \sim \lambda^{a-3/p}$ — but it does say that no sharpening of Lemma 3.1 can extend

Theorem 6.3 to a critical or supercritical quantity, and that the single exponent p is the only freedom the criterion has.

Remark 3.6 (why $s = 0$ differs in kind). At $s = 0$ the symbol of T_0 is homogeneous of degree 0 and T_0 is a Calderón–Zygmund operator, unbounded from L^∞ to L^∞ . Lemma 3.1 then fails for every p , and the classical route substitutes the logarithmic Sobolev inequality of Brezis–Gallouët–Wainger [5, 6], in the form $\|\nabla u\|_{L^\infty} \lesssim \|\omega\|_{L^\infty}(1 + \log(e + \|u\|_{H^r})) + \|\omega\|_{L^2}$, which is precisely the logarithm in [4].

4. THE A PRIORI ESTIMATE

Throughout this section $s > 0$ and w is a smooth solution of (4) on $[0, T)$ with $u = S_s w$ as in Lemma 2.4. We use two standard estimates. The first is the Kato–Ponce commutator estimate [17]: for $\sigma > 0$,

$$\|[\Lambda^\sigma, f]g\|_{L^2} \lesssim \|\nabla f\|_{L^\infty} \|g\|_{H^{\sigma-1}} + \|f\|_{H^\sigma} \|g\|_{L^\infty}. \quad (6)$$

The second is the Moser product estimate [18, Ch. 3]: $\|fg\|_{H^\sigma} \lesssim \|f\|_{L^\infty} \|g\|_{H^\sigma} + \|f\|_{H^\sigma} \|g\|_{L^\infty}$.

Lemma 4.1 (commutator with a divergence-free drift). *Let $r > 0$ and let u be divergence-free. Then*

$$\|[\Lambda^r, u \cdot \nabla]w\|_{L^2} \lesssim \|\nabla u\|_{L^\infty} \|w\|_{H^r} + \|u\|_{H^{r+1}} \|w\|_{L^\infty}.$$

Proof. Write $[\Lambda^r, u]w$ for the array with entries $[\Lambda^r, u^j]w$. Since $\operatorname{div} u = 0$ we have $\partial_j(u^j w) = u^j \partial_j w$, and therefore

$$\operatorname{div}([\Lambda^r, u]w) = \sum_j \partial_j(\Lambda^r(u^j w) - u^j \Lambda^r w) = \Lambda^r(u \cdot \nabla w) - u \cdot \nabla \Lambda^r w = [\Lambda^r, u \cdot \nabla]w.$$

The commutator on the left carries no derivative of w ; that is the whole point. Since $\|\operatorname{div} F\|_{L^2} \leq \|\Lambda F\|_{L^2}$,

$$\|[\Lambda^r, u \cdot \nabla]w\|_{L^2} \leq \|\Lambda[\Lambda^r, u]w\|_{L^2} \leq \|[\Lambda^{r+1}, u]w\|_{L^2} + \|[\Lambda, u]\Lambda^r w\|_{L^2},$$

using the identity $\Lambda[\Lambda^r, u]w = [\Lambda^{r+1}, u]w - [\Lambda, u]\Lambda^r w$, which is immediate on expanding both sides. For the first term, (6) at order $r + 1$ gives $\|\nabla u\|_{L^\infty} \|w\|_{H^r} + \|u\|_{H^{r+1}} \|w\|_{L^\infty}$. For the second, Calderón’s commutator theorem [10] gives $\|[\Lambda, u]h\|_{L^2} \lesssim \|\nabla u\|_{L^\infty} \|h\|_{L^2}$, applied with $h = \Lambda^r w$. $\square$

Lemma 4.2 (transport bound). *For every $q \in [1, \infty]$ and $t < T$,*

$$\|w(t)\|_{L^q} \leq \|w_0\|_{L^q} \exp\left(\int_0^t \|\nabla u(\tau)\|_{L^\infty} d\tau\right).$$

Proof. Along the flow of u , (4) gives $\frac{d}{dt}|w| \leq |\nabla u| |w|$ pointwise, and the flow preserves Lebesgue measure because $\operatorname{div} u = 0$; integrate. $\square$

Proposition 4.3 (a priori estimate). *Let $s > 0$ and $r > \frac{5}{2}$. Then, with $C = C(r, s)$,*

$$\frac{d}{dt} \|w\|_{H^r} \leq C(\|\nabla u\|_{L^\infty} + \|w\|_{L^\infty}) \|w\|_{H^r}.$$

Proof. Pair Λ^r of (4) with $\Lambda^r w$:

$$\frac{1}{2} \frac{d}{dt} \|w\|_{H^r}^2 = -\langle [\Lambda^r, u \cdot \nabla]w, \Lambda^r w \rangle - \langle u \cdot \nabla \Lambda^r w, \Lambda^r w \rangle + \langle \Lambda^r(w \cdot \nabla u), \Lambda^r w \rangle.$$

The middle term vanishes because $\operatorname{div} u = 0$.

Stretching term. By the Moser estimate and then (5),

$$\|\Lambda^r(w \cdot \nabla u)\|_{L^2} \lesssim \|w\|_{L^\infty} \|\nabla u\|_{H^r} + \|w\|_{H^r} \|\nabla u\|_{L^\infty} \lesssim (\|w\|_{L^\infty} + \|\nabla u\|_{L^\infty}) \|w\|_{H^r},$$

using $\|\nabla u\|_{H^r} \lesssim \|w\|_{H^{r-2s}} \leq \|w\|_{H^r}$ for $s > 0$.

Commutator term. By Lemma 4.1 and then (5),

$$\|[\Lambda^r, u \cdot \nabla]w\|_{L^2} \lesssim \|\nabla u\|_{L^\infty} \|w\|_{H^r} + \|u\|_{H^{r+1}} \|w\|_{L^\infty} \lesssim (\|\nabla u\|_{L^\infty} + \|w\|_{L^\infty}) \|w\|_{H^r},$$

since $\|u\|_{H^{r+1}} \lesssim \|w\|_{H^{r-2s}} \leq \|w\|_{H^r}$, again by (5) and $s > 0$.

Both contributions are thus at most $C(\|\nabla u\|_{L^\infty} + \|w\|_{L^\infty}) \|w\|_{H^r}^2$; dividing by $\|w\|_{H^r}$ gives the statement. $\square$

Remark 4.4 (no threshold in r). The estimate holds for every $r > \frac{5}{2}$, with no threshold depending on s , and its proof uses no interpolation. What buys this is the divergence form in Lemma 4.1: writing the commutator as $\text{div}([\Lambda^r, u]w)$ moves the derivative off w and onto u , and by (5) u has $1 + 2s$ derivatives to spare, of which the manoeuvre spends one. Applying (6) directly instead leaves $\|u\|_{H^r} \|\nabla w\|_{L^\infty}$, whose second factor has then to be interpolated against $\|w\|_{H^r}$; that route closes only for r above an s -dependent threshold which diverges as $s \rightarrow 0^+$, and is an artefact of the manoeuvre rather than of the equation. At $s = 0$ nothing is left over, the two routes coincide, and the estimate above is the classical one for the Euler equation.

Uniqueness is not assumed; it follows from the smoothing of S_s and T_s by an L^2 argument.

Proposition 4.5 (uniqueness). *Let $s > 0$, $r > \frac{5}{2}$ and $T > 0$. If $w_1, w_2 \in C([0, T]; H^r)$ both solve (4) with $w_1(0) = w_2(0)$, then $w_1 \equiv w_2$.*

Proof. Put $\delta = w_1 - w_2$ and $v = u_1 - u_2 = S_s \delta$. Subtracting the two copies of (4),

$$\partial_t \delta + (u_1 \cdot \nabla) \delta + (v \cdot \nabla) w_2 = (\delta \cdot \nabla) u_1 + (w_2 \cdot \nabla) v.$$

Pair with δ in L^2 . The term $\langle (u_1 \cdot \nabla) \delta, \delta \rangle$ vanishes since $\text{div } u_1 = 0$. Because $s > 0$, the symbols $|k|^{-1-2s}$ and $|k|^{-2s}$ of S_s and T_s are bounded by 1 on $|k| \geq 1$, so $\|v\|_{L^2} \leq \|\delta\|_{L^2}$ and $\|\nabla v\|_{L^2} = \|T_s \delta\|_{L^2} \leq \|\delta\|_{L^2}$. Hence

$$|\langle (v \cdot \nabla) w_2, \delta \rangle| \leq \|\nabla w_2\|_{L^\infty} \|\delta\|_{L^2}^2, \quad |\langle (\delta \cdot \nabla) u_1, \delta \rangle| \leq \|\nabla u_1\|_{L^\infty} \|\delta\|_{L^2}^2,$$

$$|\langle (w_2 \cdot \nabla) v, \delta \rangle| \leq \|w_2\|_{L^\infty} \|\nabla v\|_{L^2} \|\delta\|_{L^2} \leq \|w_2\|_{L^\infty} \|\delta\|_{L^2}^2.$$

Since $r > \frac{5}{2}$, the three coefficients are bounded on $[0, T]$ by $C(\|w_1\|_{C([0, T]; H^r)} + \|w_2\|_{C([0, T]; H^r)})$. Grönwall with $\delta(0) = 0$ gives $\delta \equiv 0$. $\square$

5. LOCAL EXISTENCE

It remains to produce a solution. We do so by Galerkin truncation, which on $\mathbb{T}^3$ is cleaner than mollification: the Fourier projection is self-adjoint, idempotent, and commutes with every operator appearing in the problem. Solutions are taken in the class

Definition 5.1. For $T > 0$ and $r > \frac{5}{2}$ set

$$\mathcal{X}_T^r := L^\infty([0, T]; H^r) \cap \bigcap_{\frac{5}{2} < r' < r} C([0, T]; H^{r'}),$$

and call $w \in \mathcal{X}_T^r$ a *solution* of (4) if the equation holds in $C([0, T]; H^{r'-1})$ for every such r' .

Every statement above holds in this class: Propositions 4.3 and 4.5 and Lemma 4.2 use only the size of $\|w(t)\|_{H^r}$, not continuity in H^r . Membership of H^r at *every* time, rather than almost every time, is not free, and is what the continuation argument of Section 6 needs; it follows from the class.

Lemma 5.2 (weak continuity). *Let $w \in \mathcal{X}_T^r$ and let M be the essential supremum of $\|w(t)\|_{H^r}$ on $[0, T]$. Then $w(t) \in H^r$ for every $t \in [0, T]$, with $\sup_{[0, T]} \|w(t)\|_{H^r} = M$, and $w : [0, T] \rightarrow H^r$ is weakly continuous.*

Proof. Fix t and choose $t_n \rightarrow t$ with $\|w(t_n)\|_{H^r} \leq M$; this is possible since the set of such times has full measure. Fix $r' \in (\frac{5}{2}, r)$. The sequence $(w(t_n))$ is bounded in H^r , so a subsequence converges weakly in H^r to some z , and weak convergence in H^r implies weak convergence in $H^{r'}$. As $w \in C([0, T]; H^{r'})$ we also have $w(t_n) \rightarrow w(t)$ strongly, hence weakly, in $H^{r'}$. Therefore $z = w(t)$, so $w(t) \in H^r$, and weak lower semicontinuity of the norm gives $\|w(t)\|_{H^r} \leq M$. The reverse inequality is immediate, so the supremum equals M .

For weak continuity, let ψ be a trigonometric polynomial. Then $\langle \Lambda^r w(t), \Lambda^r \psi \rangle = \langle w(t), \Lambda^{2r} \psi \rangle$ is continuous in t because $w \in C([0, T]; L^2)$. Such ψ are dense in H^r and $\|w(t)\|_{H^r} \leq M$ uniformly, so $t \mapsto w(t)$ is weakly continuous into H^r . $\square$

A solution in $\mathcal{X}_T^r$ is in fact *strongly* continuous into H^r . That is the argument of Bona and Smith [8], which applies here unchanged — the equation (4) is reversible, $-w(-t)$ solving it whenever $w(t)$ does, which is what the argument uses to pass from continuity of the norm at $t = 0$ to continuity at every time. We omit it because no statement in this paper invokes it: Lemma 5.2 supplies everything used below.

Theorem 5.3 (local existence). *Let $s > 0$ and $r > \frac{5}{2}$. There is $C = C(r, s)$ such that every mean-zero divergence-free $w_0 \in H^r(\mathbb{T}^3)$, $w_0 \neq 0$, admits a solution $w \in \mathcal{X}_T^r$ of (4) with $w(0) = w_0$ on $[0, T]$, where*

$$T = \tau(\|w_0\|_{H^r}), \quad \tau(\rho) := \frac{1}{2C\rho},$$

and moreover $\sup_{[0, T]} \|w\|_{H^r} \leq 2\|w_0\|_{H^r}$. The function τ is non-increasing.

Proof. Write $B(w) := -[u, w] = -(u \cdot \nabla)w + (w \cdot \nabla)u$ with $u = S_s w$, so that (4) reads $\partial_t w = B(w)$, and put $\rho := \|w_0\|_{H^r}$.

Step 1: the truncated system. For $N \in \mathbb{N}$ let P_N be the Fourier projection onto $\{0 < |k| \leq N\}$ and $V_N := P_N \mathfrak{g}$, a finite-dimensional space of mean-zero divergence-free trigonometric fields. Consider

$$\partial_t w^N = P_N B(w^N), \quad w^N(0) = P_N w_0. \quad (7)$$

The right-hand side maps V_N to V_N : $B(w)$ is a Lie bracket of divergence-free fields, hence divergence-free, and has zero mean because $\int [u, w] = \int ((\operatorname{div} w)u - (\operatorname{div} u)w) = 0$. It is a quadratic polynomial map of the finite-dimensional space V_N , so Picard–Lindelöf gives a unique maximal solution on some $[0, T_N)$.

Step 2: a bound uniform in N . Since P_N is self-adjoint, commutes with Λ^r , and fixes w^N ,

$$\frac{1}{2} \frac{d}{dt} \|w^N\|_{H^r}^2 = \langle \Lambda^r P_N B(w^N), \Lambda^r w^N \rangle = \langle \Lambda^r B(w^N), \Lambda^r P_N w^N \rangle = \langle \Lambda^r B(w^N), \Lambda^r w^N \rangle,$$

so the truncation is invisible and the proof of Proposition 4.3 applies to w^N verbatim, with the same constant. Each term of the bracket there is $\lesssim \|w^N\|_{H^r}$: indeed $\|w\|_{L^\infty} \lesssim \|w\|_{H^r}$ since $r > \frac{3}{2}$; $\|\nabla u\|_{L^\infty} = \|T_s w\|_{L^\infty} \lesssim \|T_s w\|_{H^2} \lesssim \|w\|_{H^{2-2s}} \leq \|w\|_{H^r}$ by (5), using $s > 0$ and $r > 2$. Hence

$$\frac{d}{dt} \|w^N\|_{H^r} \leq C \|w^N\|_{H^r}^2. \quad (8)$$

As $\|w^N(0)\|_{H^r} \leq \rho$, integrating (8) gives $\|w^N(t)\|_{H^r} \leq \rho/(1 - C\rho t) \leq 2\rho$ for $t \leq T = 1/(2C\rho)$, uniformly in N . All norms on V_N are equivalent, so this bound also forbids the ODE from leaving every ball, whence $T_N > T$ and w^N exists on $[0, T]$.

Step 3: Cauchy in L^2 . Let $M > N$ and $\delta := w^M - w^N$, so $P_M \delta = \delta$ and

$$\partial_t \delta = P_M [B(w^M) - B(w^N)] - (P_M - P_N)B(w^N).$$

Pair with δ and use $\langle P_M F, \delta \rangle = \langle F, \delta \rangle$. The first term is bounded exactly as in the proof of Proposition 4.5 — that argument used only that both fields lie in H^r with $r > \frac{5}{2}$ — by $C(\|w^M\|_{H^r} + \|w^N\|_{H^r})\|\delta\|_{L^2}^2 \leq C\rho\|\delta\|_{L^2}^2$. For the second, H^{r-1} is an algebra because $r-1 > \frac{3}{2}$, so by (5) $\|B(w^N)\|_{H^{r-1}} \lesssim \|u^N\|_{H^r}\|w^N\|_{H^r} \lesssim \rho^2$, while $P_M - P_N$ is supported in $|k| > N$, whence $\|(P_M - P_N)B(w^N)\|_{L^2} \leq N^{-(r-1)}\|B(w^N)\|_{H^{r-1}} \lesssim \rho^2 N^{-(r-1)}$. Therefore

$$\frac{d}{dt}\|\delta\|_{L^2} \leq C\rho\|\delta\|_{L^2} + C\rho^2 N^{-(r-1)},$$

and $\|\delta(0)\|_{L^2} = \|(P_M - P_N)w_0\|_{L^2} \leq N^{-r}\rho$, so Grönwall gives $\sup_{[0,T]}\|\delta\|_{L^2} \leq C(\rho, T)N^{-(r-1)}$ uniformly in M . Thus (w^N) is Cauchy in $C([0, T]; L^2)$.

Step 4: passage to the limit. Let w be the limit in $C([0, T]; L^2)$. For $r' < r$, interpolation and Step 2 give $\|w^N - w^M\|_{H^{r'}} \leq \|w^N - w^M\|_{L^2}^{1-r'/r} \|w^N - w^M\|_{H^r}^{r'/r}$, so (w^N) is Cauchy in $C([0, T]; H^{r'})$ and $w \in C([0, T]; H^{r'})$. Weak-* compactness with lower semicontinuity of the norm gives $w \in L^\infty([0, T]; H^r)$ with $\sup_{[0,T]}\|w\|_{H^r} \leq 2\rho$. Fix $r' \in (\frac{5}{2}, r)$. Then $H^{r'-1}$ is an algebra and, by (5), B is locally Lipschitz from $H^{r'}$ to $H^{r'-1}$, so $B(w^N) \rightarrow B(w)$ in $C([0, T]; H^{r'-1})$. Also $P_N \rightarrow I$ strongly on $H^{r'-1}$, hence uniformly on the compact set $\{B(w(t)) : t \in [0, T]\}$. Passing to the limit in (7) gives $\partial_t w = B(w)$ in $C([0, T]; H^{r'-1})$. Hence $w \in \mathcal{X}_T^r$ solves (4) with $w(0) = w_0$. $\square$

Remark 5.4. The scheme is the classical one [18, Ch. 3]. What lets it run without modification is that S_s and T_s are *smoother* than their Euler counterparts by $2s$ derivatives, so every estimate above has room to spare; nothing in the argument is sharp in s , and none of it would survive $s < 0$, where (5) reverses.

6. THE CRITERION

Proposition 6.1 (maximal solution and the blowup alternative). *Let $s > 0$ and $r > \frac{5}{2}$. Each mean-zero divergence-free $w_0 \in H^r$ admits a unique maximal solution on some $[0, T_*)$, $T_* \in (0, \infty]$, and if $T_* < \infty$ then*

$$\sup_{t < T_*} \|w(t)\|_{H^r} = \infty.$$

Proof. By Proposition 4.5 any two solutions with the same datum agree on the intersection of their intervals, so the union of all solutions is a well-defined solution on a maximal $[0, T_*)$. Suppose $T_* < \infty$ and $M := \sup_{t < T_*} \|w(t)\|_{H^r} < \infty$. Pick $t_0 < T_*$ with $T_* - t_0 < \tau(M)$, which is possible as $\tau(M) > 0$. By Lemma 5.2 the datum $w(t_0)$ lies in H^r — at that particular time, not merely almost everywhere — with $\|w(t_0)\|_{H^r} \leq M$, so Theorem 5.3 and the monotonicity of τ give a solution on $[t_0, t_0 + \tau(M)]$, and by Proposition 4.5 it extends w past T_* . This contradicts maximality. $\square$

Theorem 6.2 (continuation in $\|\nabla u\|_{L^\infty}$). *Let $s > 0$ and $r > \frac{5}{2}$. If $T < \infty$ and*

$$\int_0^T \|\nabla u(t)\|_{L^\infty} dt < \infty,$$

then $\sup_{t < \min(T, T_)} \|w(t)\|_{H^r} < \infty$; in particular $T < T_*$.*

Proof. Write $I := \int_0^T \|\nabla u\|_{L^\infty} dt < \infty$. By Lemma 4.2, $\|w(t)\|_{L^2}$ and $\|w(t)\|_{L^\infty}$ are bounded on $[0, \min(T, T_*)]$ by $\|w_0\|_{L^2} e^I$ and $\|w_0\|_{L^\infty} e^I$. Hence the bracket in Proposition 4.3 is bounded by $\|\nabla u(t)\|_{L^\infty} + M$ for a constant M depending only on $\|w_0\|_{L^2}$, $\|w_0\|_{L^\infty}$, I , r and s . Grönwall gives

$$\|w(t)\|_{H^r} \leq \|w_0\|_{H^r} e^{C(I+MT)},$$

which is finite. By the blowup alternative of Proposition 6.1 this is incompatible with $T_* \leq T$. $\square$

Theorem 6.3 (the criterion). *Let $s > 0$, $r > \frac{5}{2}$, and let $p \in [1, \infty]$ satisfy $p > 3/(2s)$. If $T < \infty$ and*

$$\int_0^T \|w(t)\|_{L^p} dt < \infty,$$

then the solution does not break down at or before T .

Proof. By Lemma 3.1, $\|\nabla u(t)\|_{L^\infty} \leq C(s, p)\|w(t)\|_{L^p}$ for each t , so

$$\int_0^T \|\nabla u\|_{L^\infty} dt \leq C(s, p) \int_0^T \|w\|_{L^p} dt < \infty,$$

and Theorem 6.2 applies. $\square$

Remark 6.4. Only p is constrained by s . As $s \rightarrow 0^+$ the admissible exponent $p > 3/(2s)$ escapes to ∞ and no finite p survives the limit, which is the correct behaviour: at $s = 0$ the estimate of Lemma 3.1 is false for every p (Remark 3.6) and the logarithm is unavoidable. The regularity requirement is $r > \frac{5}{2}$ throughout, independently of s , by Remark 4.4.

7. WHAT THE CRITERION DOES NOT DO

It recovers $5/2$ and moves nothing. By Lemma 2.5 the conserved quantity is $\|w\|_{\dot{H}^{-1-s}}$. By Lemma 2.4 and Sobolev embedding, $\|\nabla u\|_{L^\infty} = \|T_s w\|_{L^\infty} \lesssim \|w\|_{\dot{H}^{3/2-2s+\varepsilon}}$ for every $\varepsilon > 0$, and the conserved norm dominates this exactly when

$$\frac{3}{2} - 2s \leq -1 - s \iff s \geq \frac{5}{2},$$

the geodesic completeness threshold of [9] and, as $s > \frac{1}{2} \dim M + 1$, the same count that returns $3/2$ on $\text{Diff}(S^1)$ [16, 3]. So Theorem 6.3 is consistent with the known completeness result and improves on it nowhere.

That is the outcome to insist on. A continuation criterion improving the threshold by itself would, continued to $s = 0$, prove global regularity for the three-dimensional Euler equation. What the criterion does is make the problem definite: bound $\|w\|_{L^p}$ for a single $p > 3/(2s)$, given only that $\|w\|_{\dot{H}^{-1-s}}$ is conserved and that w obeys (4). Since $-1 - s < 0$ and negative-regularity spaces do not embed in L^p for $p \geq 2$, the conservation law cannot supply the hypothesis directly at any $s < 5/2$, and the gain must come from the equation.

The threshold is the critical exponent, not an artefact. It is worth saying why $5/2$ is where it is, because the answer removes a hope. Equation (4) admits the dilations $w_\lambda(x, t) = \lambda^a w(\lambda x, \lambda^b t)$. As S_s is homogeneous of degree $-1 - 2s$ we get $u_\lambda = \lambda^{a-1-2s} u(\lambda x, \lambda^b t)$, so both nonlinear terms carry λ^{2a-2s} while $\partial_t w$ carries λ^{a+b} , and the equation is invariant precisely when $b = a - 2s$. In dimension d one has $\|\lambda^a f(\lambda \cdot)\|_{\dot{H}^\sigma} = \lambda^{a-d/2+\sigma} \|f\|_{\dot{H}^\sigma}$, so by Lemma 2.5 the conserved energy $\|w\|_{\dot{H}^{-1-s}}^2$ is scale-invariant exactly at $a = \frac{d}{2} + 1 + s$, and then

$$b = \frac{d}{2} + 1 - s.$$

The energy-invariant dilation therefore concentrates in finite time when $s < \frac{d}{2} + 1$ and spreads when $s > \frac{d}{2} + 1$. That is,

$$s^* = \frac{d}{2} + 1$$

is the scaling-critical exponent. In $d = 3$ this is $5/2$, the threshold of [9]; in $d = 1$ it is $3/2$, which is the *sharp* threshold on $\text{Diff}(S^1)$ [16, 3]. So on the circle, where the answer is known, completeness holds exactly in the subcritical range and breakdown is known, and conjectured throughout, exactly in the supercritical range.

This is corroborated from an unexpected direction, and we should record that the family (1) is not new. Olson and Titi [20] study, on the periodic torus, the two-parameter system

$$\partial_t v + \nu(-\Delta)^{\theta_2} v - u \times (\nabla \times v) + \nabla p = f, \quad v = u + (-\alpha^2 \Delta)^{\theta_1} u,$$

whose inviscid case is (2) with $s = \theta_1$ — as they note, it is variational with Hamiltonian the H^{θ_1} norm — and which is the Navier–Stokes equation at $\theta_1 = 0, \theta_2 = 1$ and the Navier–Stokes- α model at $\theta_1 = \theta_2 = 1$. Their main theorem is global existence, uniqueness and regularity when

$$\nu > 0 \text{ and } 2\theta_1 + 4\theta_2 \geq 5, \quad \text{or} \quad \nu = 0 \text{ and } 2\theta_1 \geq 5.$$

The second condition is $s \geq \frac{5}{2}$, an analytic proof of the inviscid half of [9] predating it, and the first is the same line with a viscous dial: repeating the computation above for that system, balancing the dissipation against ∂_t forces $b = 2\theta_2$ and against the nonlinearity $a = 2\theta_2 - 1$, so that $E = \|u\|_{\dot{H}^{\theta_1}}^2$ is scale-invariant exactly when $\theta_1 + 2\theta_2 = \frac{5}{2}$. Their hypothesis is therefore precisely scale-subcriticality, and their theorem says that the whole closed subcritical region is regular. Two consequences for what follows. The identification of $\frac{5}{2}$ as criticality rather than as an artefact is confirmed independently, at both ends of a line rather than at one point. And Question 7.2 of Paper I is harder than the phrase “nothing is known below $\frac{5}{2}$ ” suggests: what it asks for is a *strictly supercritical* global regularity theorem, the subcritical side having been settled in 2007.

Two consequences, neither of them the obvious one. First, the coincidence of the count of the previous subsection with $\frac{d}{2} + 1$ is no accident of the method — but it is a statement about the *energy*, not about the problem. The energy is scale-invariant exactly at s^* , so below s^* it is blind to concentration and no argument resting on it alone can reach there. The same count puts the criterion of Theorem 6.3 on the other side of the line: $\|\lambda^a w(\lambda \cdot)\|_{L^p} = \lambda^{a-d/p} \|w\|_{L^p}$ against $\|\nabla u\|_{L^\infty} \sim \lambda^{a-2s}$, so L^p is subcritical exactly when $p > d/(2s)$ — which is precisely the range Lemma 3.1 admits, $3/(2s)$ in three dimensions and $1/s$ in two. The criterion is therefore stated in a subcritical quantity, and the question is whether any such quantity can be controlled. Both counts are checked in `scaling_check.py`.

In two dimensions one can be, and the whole supercritical range falls.

Proposition 7.1 (the two-dimensional case). *Let $d = 2$ and $s > 0$. The geodesic flow of the metric (1) on $\text{SDiff}(\mathbb{T}^2)$ is globally regular: for every $r > \frac{5}{2}$ and every mean-zero divergence-free $w_0 \in H^r(\mathbb{T}^2)$ the solution exists for all time.*

Proof. Sections 2–6 go through in $d = 2$ with two changes. First, in the proof of Proposition 2.2 the field $w = \text{curl } m$ is a scalar and dm a 2-form, so $\mathcal{L}_u(w \text{ vol}) = (u \cdot \nabla w) \text{ vol}$ for divergence-free u , and the reduction reads

$$\partial_t w + (u \cdot \nabla) w = 0$$

with *no* stretching term — there being no way to form $(w \cdot \nabla)u$ out of a scalar. The transport is by a measure-preserving flow, so $\|w(t)\|_{L^p} = \|w_0\|_{L^p}$ for every p and every t . Second, the

kernel in Lemma 3.1 is homogeneous of degree $2s - 2$ and lies in $L^{p'}(\mathbb{T}^2)$ iff $(2 - 2s)p' < 2$, that is iff $p > 1/s$. Fixing such a p ,

$$\int_0^T \|\nabla u\|_{L^\infty} dt \leq C(s, p) T \|w_0\|_{L^p} < \infty$$

for every T , and Theorem 6.2, whose proof is unchanged, forbids breakdown at T . As T was arbitrary the solution is global. $\square$

Remark 7.2. The argument is elementary once the reduction is in hand and we claim no novelty for it; its role here is comparative, and it is worth three observations. It fails at $s = 0$, where Lemma 3.1 is unavailable for every p and one is returned to the classical two-dimensional theory and its logarithm. Since $s^* = 2$ in two dimensions, it covers the whole supercritical range $[0, 2)$: on $\text{SDiff}(\mathbb{T}^2)$ volume preservation buys every order below criticality, so scale-criticality bounds what the energy can do and not what the equation does. And the mechanism is exactly the one absent in three dimensions — the subcritical L^p norms are conserved there because nothing stretches. The identity underlying the reduction is checked in `dim2_check.py`.

It is tempting, separately, to read the one-dimensional case as evidence that the threshold sits *at* criticality — on $\text{Diff}(S^1)$ it does, exactly — and to expect the same in three dimensions. That reading is wrong. $\text{Diff}(S^1)$ is the full diffeomorphism group, and the comparison bearing on $\text{SDiff}(\mathbb{T}^3)$ is between Diff and SDiff in a *fixed* dimension; carried out in $d = 2$ by Proposition 7.1, against the threshold s^* which Diff attains in $d = 1$ [16, 3] and in $d \geq 3$ [7, 9], it goes the other way.

The two analogies thus point in opposite directions, and three dimensions is undetermined by either. What distinguishes it is that the stretching term neither vanishes, as in $d = 2$, nor acts unimpeded, as on Diff : it is present but depleted, by the alignment cancellation of Lemma 7.6, which is the three-dimensional residue of the two-dimensional vanishing. How much that depletion is worth admits a definite answer, in two parts, and neither is a reserve of unused strength.

Unconditionally it is worth nothing. Theorem 6.3 already rests on the bound $|\kappa_s(y)| \leq C|y|^{2s-3}$ of Lemma 3.1, whose exponent is sharp (Remark 3.2). The alignment factor multiplies that kernel by $|D_s| \leq \min\{C, C|\xi' - \xi|\}$, and absent a hypothesis on ξ the second bound is no improvement on the first, $|\xi' - \xi| \leq 2$ always: a direction field may turn through a right angle on the scale at which $|w|$ varies, and then nothing is depleted at all. Measured, the effect is starker than that argument suggests. For random band-limited divergence-free fields the ratio $|\alpha(x)|/\|S(x)\|_2$, at most 1 pointwise since ξ is a unit vector, attains 0.9999 at $s = \frac{1}{4}, \frac{1}{2}, 1, 2$ alike (`deplete_ratio.py`): the stretching rate saturates the trivial bound, uniformly in s .

Conditionally it is worth exactly $\beta/2$ in s , as computed above. The same script shows the mechanism switching on as one drives the direction field towards a constant: the ratio falls to 0.14 once the angular spread of ξ over the region carrying the vorticity is reduced to 0.02. That regime is the hypothesis of [12], and in the limit it is Proposition 7.1.

So the depletion is the conditional mechanism and nothing besides, and it is already accounted for in Theorem 6.3. What remains open is therefore not its size but whether some *other* structure controls a subcritical quantity in three dimensions — which by the count of the previous subsection is the question whether $s_c < \frac{5}{2}$, and at $s = 0$ is the global regularity of the Euler equation. We have nothing to offer on it, and prefer to say so in that form rather than as a direction to look in.

A logarithm below the endpoint, bought without the structure. The inclusive endpoint $s = \frac{5}{2}$ in [20] is a Brezis–Gallouët statement. There the energy controls ∇u in $H^{3/2}$, which

misses L^∞ by a logarithm, and Grönwall absorbs the logarithm. Pushed through Osgood's lemma, the same argument reaches a whole logarithm below $\frac{5}{2}$. We record this because it bears on what the structure may be credited with. It needs one estimate that Proposition 4.3 does not supply. The bracket there contains $\|w\|_{L^\infty}$, which the energy does not control and which by Lemma 4.2 may grow like $\exp \int \|\nabla u\|_{L^\infty}$, and any logarithmic closure is lost to it. In the momentum form (3) that term does not appear.

For this subsection only, let the inertia symbol be $a(|k|)$ with $a(r) = r^{2s}\ell(r)$. Here $\ell > 0$ is smooth, $|r^j \ell^{(j)}(r)| \leq C_j \ell(r)$, and $r^{-\varepsilon} \lesssim \ell(r) \lesssim r^\varepsilon$ for every $\varepsilon > 0$; $\ell \equiv 1$ is (1). Let $\mathcal{Q}_j$ be the Littlewood–Paley block at frequency 2^j and $\mathcal{S}_j = \sum_{i < j} \mathcal{Q}_i$.

Lemma 7.3 (the momentum form). *Let $s > \frac{1}{2}$ and $\sigma > 0$. For a smooth solution of (3),*

$$\frac{d}{dt} \|m\|_{H^\sigma} \leq C \|\nabla u\|_{L^\infty} \|m\|_{H^\sigma}, \quad C = C(\sigma, s, \ell).$$

Proof. Pair Λ^σ of (3) with $\Lambda^\sigma m$. The pressure drops out because $\text{div } m = 0$, and $\langle u \cdot \nabla \Lambda^\sigma m, \Lambda^\sigma m \rangle = 0$. What is left is

$$\frac{1}{2} \frac{d}{dt} \|m\|_{H^\sigma}^2 = -\langle [\Lambda^\sigma, u \cdot \nabla] m, \Lambda^\sigma m \rangle - \langle \Lambda^\sigma ((\nabla u)^\top m), \Lambda^\sigma m \rangle.$$

Decompose the products by Bony's paraproduct. The standard estimates [2, Lemma 2.100, Thms. 2.82 and 2.85] bound three kinds of piece in L^2 by $C \|\nabla u\|_{L^\infty} \|m\|_{H^\sigma}$: every paraproduct in which u or ∇u is the lower-frequency factor, the commutator of Λ^σ with $T_u \cdot \nabla$, and every remainder, the last using $\sigma > 0$. Three pieces remain, those in which m or a derivative of it is the lower-frequency factor: $T_{\nabla m} u$ and $T_{\nabla \Lambda^\sigma m} u$ from the two halves of the commutator, and $T_m \nabla u$ from the second term. These are where Proposition 4.3 spends $\|w\|_{L^\infty}$. Instead, Bernstein's inequality on each block gives $\|\mathcal{Q}_i m\|_{L^\infty} \lesssim a(2^i) \|\mathcal{Q}_i u\|_{L^\infty} \lesssim a(2^i) 2^{-i} \|\nabla u\|_{L^\infty}$, and therefore

$$\|\mathcal{S}_j \nabla m\|_{L^\infty} \lesssim \sum_{i < j} a(2^i) \|\nabla u\|_{L^\infty} \lesssim a(2^j) \|\nabla u\|_{L^\infty}, \quad \|\mathcal{S}_j m\|_{L^\infty} \lesssim \sum_{i < j} \frac{a(2^i)}{2^i} \|\nabla u\|_{L^\infty} \lesssim \frac{a(2^j)}{2^j} \|\nabla u\|_{L^\infty}.$$

Both sums are dominated by their last term, because $a(r)$ and $a(r)/r$ grow at least like positive powers of r . For the second this is exactly $2s > 1$; at $s = \frac{1}{2}$ the sum is $\sum_{i < j} \ell(2^i)$, and one loses a logarithm, m being then a Riesz transform of ∇u . Consequently the frequency- 2^j block of $T_{\nabla m} u$ has L^2 norm at most $a(2^j) \|\nabla u\|_{L^\infty} \|\mathcal{Q}_j u\|_{L^2} \approx \|\nabla u\|_{L^\infty} \|\mathcal{Q}_j m\|_{L^2}$. The block of $T_m \nabla u$ is at most $a(2^j) 2^{-j} \|\nabla u\|_{L^\infty} \cdot 2^j \|\mathcal{Q}_j u\|_{L^2}$, the same. The block of $T_{\nabla \Lambda^\sigma m} u$ obeys the same bound with $2^{j\sigma} a(2^j)$ in place of $a(2^j)$. Summing in ℓ^2 with the weight $2^{j\sigma}$ bounds all three by $C \|\nabla u\|_{L^\infty} \|m\|_{H^\sigma}$. $\square$

Proposition 7.4 (a logarithm below $\frac{5}{2}$). *Let $a(r) = r^5/g(r)^2$, where $g \geq 1$ is smooth and nondecreasing with $|r^j g^{(j)}(r)| \leq C_j g(r)$, and put*

$$H(T) := \int_0^T g(e^\tau)^2 d\tau.$$

If $\int_0^\infty H(T)^{-1/2} dT = \infty$, then for every smooth datum the solution of (2) with this inertia operator is global and smooth. The hypothesis holds for $a(r) = r^5/\log(e+r)$ and for $a(r) = r^5/(\log(e+r)(\log \log(e^e+r))^2)$. It fails for $a(r) = r^5/\log(e+r)^{1+\varepsilon}$.

Proof. The local theory of Sections 4–6 uses the symbol only through (5). That bound holds here with $2s$ replaced by $5 - \varepsilon$ for any $\varepsilon > 0$, since $g(r) \lesssim r^\varepsilon$. So there is a unique maximal solution, with the blowup alternative of Proposition 6.1. Put $Z(t) = \int_0^t \|\nabla u\|_{L^\infty}$. By Lemma

7.3, $\|m(t)\|_{H^\sigma} \leq \|m_0\|_{H^\sigma} e^{CZ(t)}$ for each $\sigma > 0$. Since $\|w\|_{H^r} \lesssim \|m\|_{H^{r+1}}$, it suffices to show that Z stays finite on bounded intervals.

The energy $E = \langle Au, u \rangle$ is conserved by the Lie–Poisson cancellation in the proof of Lemma 2.5, which does not use the symbol. Up to the normalisation of the pairing, $E = \sum_k a(|k|) |\hat{u}(k)|^2$. Fix $\rho > 4$. For $N \geq 2$, Cauchy–Schwarz against E on $|k| \leq N$ and against $\|u\|_{H^\rho}$ beyond gives

$$\|\nabla u\|_{L^\infty} \leq \sum_{k \neq 0} |k| |\hat{u}(k)| \leq CE^{1/2} \left(\sum_{0 < |k| \leq N} \frac{g(|k|)^2}{|k|^3} \right)^{1/2} + CN^{5/2-\rho} \|u\|_{H^\rho}.$$

Comparing the lattice sum with $4\pi \int_1^N g(r)^2 dr/r$ bounds the first term by $CE^{1/2} (1 + H(\log N))^{1/2}$. Choose $\log N = \max\{\log 2, \log \|u\|_{H^\rho}/(\rho - \frac{5}{2})\}$. Since $\|u\|_{H^\rho} \lesssim \|m\|_{H^{\rho-4}}$, this gives $\log N \leq c_0 + CZ(t)$ and

$$Z'(t) \leq C_1 + C_2 E^{1/2} H(c_1 + c_2 Z(t))^{1/2} =: \Psi(Z(t)).$$

Ψ is positive, continuous and nondecreasing, and $\int^\infty dZ/\Psi(Z) = \infty$ is the hypothesis. By Osgood’s lemma [2, Lemma 3.4], Z is finite for all finite t .

For the examples: $g(e^\tau)^2 = \log(e + e^\tau)$ gives $H(T) \sim T^2/2$ and $\int^\infty dT/T = \infty$. The extra factor $(\log \log)^2$ gives $H(T) \sim \frac{1}{2} T^2 (\log T)^2$ and $\int^\infty dT/(T \log T) = \infty$. The power $1 + \varepsilon$ gives $H(T) \sim T^{2+\varepsilon}/(2 + \varepsilon)$ and a convergent integral. $\square$

At $g \equiv 1$ the argument is the endpoint of [20], with $Z' \lesssim 1 + Z^{1/2}$. At $g^2 = \log$ it gives Z exponential and $\|m\|$ double-exponential, the growth of the two-dimensional Euler equation.

Remark 7.5 (what it moves). First, nothing is used beyond the energy identity, Lemma 7.3 and the Brezis–Gallouët splitting [5, 6]. The bracket enters only through the conservation law, and incompressibility only through the transport term. On the full group the momentum equation gains the term $(\operatorname{div} u) m$, a product of the same kind as $(\nabla u)^\top m$, and $\langle u \cdot \nabla \Lambda^\sigma m, \Lambda^\sigma m \rangle = -\frac{1}{2} \langle (\operatorname{div} u) \Lambda^\sigma m, \Lambda^\sigma m \rangle$ is bounded by $\|\nabla u\|_{L^\infty} \|m\|_{H^\sigma}^2$. So Proposition 7.4 holds verbatim on $\operatorname{Diff}(\mathbb{T}^3)$. It is Sobolev embedding pushed to its logarithmic limit. It moves no power, and so it leaves Question 7.2 of Paper I, whose s_c is defined over the powers $|k|^{2s}$, exactly where it was. What it moves is the baseline. The region where completeness is bought without the structure is not $\{s \geq \frac{5}{2}\}$; it contains every symbol of Proposition 7.4, and a result crediting the structure has to get below that whole family.

Second, the reach is a whole logarithm, whereas the slightly supercritical two-dimensional Euler equations reach only a *double* logarithm [11, 15]. The difference is where criticality sits. There it is in L^∞ of the vorticity, where the velocity gradient loses a full logarithm. Here it is in an L^2 energy, where the Brezis–Gallouët inequality loses only its square root.

Third, the viscous end of the line of [20] admits the same extension. Take $\theta_2 = 1$, $\nu > 0$ and inertia $r/g(r)^2$. Their L^2 estimate for $q = \nabla \times v$ closes with the time-integrated dissipation in the exponent. Split q at a frequency chosen so that the dissipation absorbs the high part; what remains is an Osgood inequality, and regularity follows whenever $\int^\infty ds/(s g(s)^2) = \infty$. This is the exact counterpart of Yamazaki’s theorem for the Leray- α family [29], with the same budget of logarithm. We do not pursue it: it spends the dissipation, and so says nothing about the inviscid problem.

Lemma 7.3 is sanity-checked in `corner/lemma_m.py` on coherent wave packets at three scales. At $s = \frac{5}{2}$ the normalised pairing varies by less than a factor of 3 across the frequency ladder, while $\|w\|_{L^\infty}/\|\nabla u\|_{L^\infty}$ grows by a factor of 92. That excludes a genuinely needed $\|w\|_{L^\infty}$ term. Being a sample, it cannot exclude a logarithmic loss; only the proof does that.

Nothing above uses incompressibility. Beyond $\text{div } u = 0$ in the transport term and in Lemma 4.2, no property of the volume-preserving bracket is used. The mechanism that uses more is the geometric depletion of vortex stretching through the direction field $\xi = w/|w|$ [12]. It is worth saying precisely what that would and would not do here, because the obvious hope is misdirected.

What depletion repairs at $s = 0$ is a *borderline*. Constantin's representation of the stretching rate $\alpha = \xi \cdot (\nabla u)_{\text{sym}} \xi$ as a principal value with kernel $|y|^{-3}$ and a geometric factor vanishing as the directions align is exactly Calderón–Zygmund critical; Hölder continuity of ξ of order β bounds that factor by $|y|^\beta$ and renders the kernel integrable. For $s > 0$ there is no borderline to repair: Lemma 3.1 says the kernel is already $|y|^{2s-3}$ and already integrable, with no appeal to geometry whatever.

The cancellation itself does transfer, and for a reason that makes its scope clear.

Lemma 7.6 (alignment cancellation, for every s). *Let $s \geq 0$ and let $\sigma_{ij,k}^{(s)}(k) = -k_j \varepsilon_{iak} k_a |k|^{-2-2s}$ be the symbol of T_s from Lemma 2.4, in which i indexes the component, j the derivative and k the argument. For unit vectors ξ, ξ' put $D_s(k; \xi', \xi) := \xi_i \sigma_{ij,k}^{(s)}(k) \xi_j \xi'_k$, the factor which the representation of α integrates against $|w|$. Then*

$$D_s(k; \xi', \xi) = - \frac{(k \cdot \xi) \det(\xi, k, \xi')}{|k|^{2+2s}}.$$

In particular D_s is linear in ξ' , vanishes identically when $\xi' = \xi$, and $|D_s(k; \xi', \xi)| \leq |k|^{-2s} |\xi' - \xi|$.

Proof. Contract: $\xi_i \xi_j \xi'_k \sigma_{ij,k}^{(s)} = -(k \cdot \xi) \varepsilon_{iak} \xi_i k_a \xi'_k |k|^{-2-2s}$, and $\varepsilon_{iak} \xi_i k_a \xi'_k = \det(\xi, k, \xi')$. At $\xi' = \xi$ the determinant has a repeated argument. Linearity in ξ' is immediate, so $\det(\xi, k, \xi') = \det(\xi, k, \xi' - \xi)$, which Hadamard bounds by $|k| |\xi' - \xi|$; combined with $|k \cdot \xi| \leq |k|$ this gives the estimate. Since the contracted symbol vanishes identically at $\xi' = \xi$, so does the contracted kernel. $\square$

The s -dependence sits entirely in the scalar $|k|^{-2-2s}$ and cannot disturb the determinant, which comes from the ε tensor in $w = \text{curl } m$: the depletion is carried by the curl, not by the metric, and is in that sense indifferent to which inertia operator is used. It is checked against a brute-force contraction in `depletion_check.py`, where the vanishing at alignment is confirmed to machine zero for $s = 0, \frac{1}{4}, \frac{1}{2}, 1, 2$ against a symbol of size $O(1)$, so that the cancellation is genuine and not a vanishing tensor.

With ξ Hölder of order β we therefore get $|\xi(x+y) - \xi(x)| \lesssim |y|^\beta$ and a kernel $|y|^{2s+\beta-3}$ in place of $|y|^{2s-3}$: depletion acts exactly as though s were replaced by $s + \beta/2$. The admissible exponent in Theorem 6.3 would fall from $p > 3/(2s)$ to $p > 3/(2s + \beta)$, and the count of the previous subsection would close at $s + \beta \geq \frac{5}{2}$ rather than $s \geq \frac{5}{2}$. Since $\beta \leq 1$ for a direction field, the best conceivable version reaches $s \geq \frac{3}{2}$, and never $s = 0$; that it cannot reach Euler is the consistency one should demand, [12] being a conditional criterion and not a regularity theorem.

And conditional is the point. Hölder regularity of ξ is a property of the solution, not a consequence of the conservation law, and s_c is defined by an unconditional statement. So depletion is a candidate for a criterion sharper than Theorem 6.3 — a worthwhile thing, and one which by Lemma 4.2 and $\|w\|_{L^p} \lesssim \|w\|_{L^\infty}$ would feed straight into it — and not a candidate for displacing s_c . We know of no mechanism that would do the latter, and record that statement. We make no claims that it is a direction to look in.

The estimate reverses for $s < 0$. For $s < 0$ the multiplier T_s has positive order $2|s|$: it loses derivatives rather than gaining them, Lemma 3.1 is false for every p , and $\|\nabla u\|_{L^\infty}$ is controlled by no L^p norm of w whatever. This is not a defect of the argument but a statement about where

the family should break down: the stretching equation (4) is unchanged, while the constitutive law recovering u is strictly rougher than Euler's own. We regard it as the analytic shape of the expectation that finite-time breakdown, if it occurs in this family, occurs at negative order.

The exponential map changes type at $s = 0$. There is a second reason why nothing proved here for $s > 0$ reaches $s = 0$, and it is geometric rather than a matter of scaling. Along a geodesic $\eta(t) = \exp(tu_0)$, after left translation to the identity, the differential of the exponential map is governed by the operator $K_{u_0}(v) = A^{-1}P(\operatorname{curl} m_0 \times v)$, where $m_0 = Au_0$ [19]. Multiplication by the fixed smooth field $\operatorname{curl} m_0$ has order zero and A^{-1} has order $-2s$, so K_{u_0} is compact on $\mathbb{T}^3$ for every $s > 0$ and is not compact at $s = 0$. From the compactness Misiólek and Preston prove that the exponential map of a right-invariant H^r metric on the volume-preserving group of a manifold of dimension at least three is a nonlinear Fredholm map of index zero, for integer $r \geq 1$, the range in which they have a smooth exponential map [19, Lemma 6.3, Thm. 6.5, Rem. 6.4]. At $r = 0$ in three dimensions the property fails [13]. We do not claim it for non-integer $s \in (0, 1)$: Section 5 gives existence and uniqueness but not differentiable dependence on the data, and the Fredholm property is a statement about that derivative.

The failure at $s = 0$ is not a technicality. It is what allows Jacobi fields to concentrate near a single particle path. For the Euler equation itself, with $\omega_0 = \operatorname{curl} u_0$, if at some interior point x the ordinary differential equation

$$\frac{d}{dt}(\Lambda(t, x) \dot{y}) + \omega_0(x) \times \dot{y} = 0, \quad \Lambda = D\eta^\top D\eta,$$

has a nontrivial solution vanishing at two times, then the geodesic has a conjugate point [21]. A WKB analysis of the Jacobi equation locates further epiconjugate points, which typically fill intervals [22]. The equation sees only the Lagrangian stretching and the initial vorticity. For $s > 0$ the corresponding index form has a leading term of order $2s$ in the variation and a rotation term of order zero, so it is positive on sufficiently concentrated variations, and no local criterion of this kind exists. That is an order count, not a proof. On the dial, then, $s = 0$ is the one point at which conjugate points in three dimensions are a local phenomenon. The blowup dichotomy built on that local theory [23, 24] has no counterpart at any $s > 0$; what it says at $s = 0$ is recorded in [25, §7], and the constraints it places on any quantity deciding regularity are collected in [30].

8. ANISOTROPIC SYMBOLS

Isotropy has been used in one place only. Lemma 3.1 invokes the kernel asymptotics of [27] for a symbol smooth away from the origin and homogeneous of degree $-2s$, and homogeneity is the whole of what that argument needs; nothing else in Sections 2–6 looks at the symbol at all beyond the two gains (5). It is worth recording what this buys, because Paper I poses its Question 7.1 in the isotropic class while noting that the anisotropic translation-covariant multipliers — which lie between that class and the inhomogeneous operators of [28] — are open as well. The criterion covers them.

Definition 8.1. Let $s > 0$. An *inertia symbol of order $2s$* is a smooth map $\alpha : \mathbb{R}^3 \setminus \{0\} \rightarrow \operatorname{Sym}(\mathbb{R}^3)$ such that

- (i) $\alpha(\lambda k) = \lambda^{2s} \alpha(k)$ for $\lambda > 0$;
- (ii) $\alpha(-k) = \alpha(k)$;
- (iii) $\alpha(k)k = 0$ and $\alpha(k)k^\perp \subseteq k^\perp$;
- (iv) $\langle \alpha(k)v, v \rangle > 0$ for every $v \in k^\perp \setminus \{0\}$.

It is *scalar* if $\alpha(k) = a(k)P_k$ with P_k the orthogonal projection onto $k^\perp$, and *isotropic* if moreover $a(k) = |k|^{2s}$, which is (1). Write A for the associated Fourier multiplier on $\mathfrak{g}$; (ii) makes it real and (iii)–(iv) make it symmetric and positive definite there.

Lemma 8.2. *Let α be an inertia symbol of order $2s$ and let $c \leq C$ be the least and greatest eigenvalue of $\alpha(n)|_{n^\perp}$ over $n \in S^2$. Then $0 < c \leq C < \infty$ and*

$$c|k|^{2s}|v|^2 \leq \langle \alpha(k)v, v \rangle \leq C|k|^{2s}|v|^2, \quad v \in k^\perp.$$

Moreover $k \mapsto \alpha(k)^{-1}$, meaning the inverse of $\alpha(k)|_{k^\perp}$ extended by zero on $\mathbb{R}k$, is smooth on $\mathbb{R}^3 \setminus \{0\}$ and homogeneous of degree $-2s$, with eigenvalues in $[C^{-1}|k|^{-2s}, c^{-1}|k|^{-2s}]$ on $k^\perp$.

Proof. The function $(n, v) \mapsto \langle \alpha(n)v, v \rangle$ is continuous and strictly positive on the compact set $\{(n, v) : |n| = |v| = 1, v \perp n\}$, so $c > 0$ and $C < \infty$; homogeneity extends the bound to all $k \neq 0$. The assignment $k \mapsto \alpha(k)|_{k^\perp}$ is a smooth endomorphism of the smooth rank-two bundle $k^\perp$ over $\mathbb{R}^3 \setminus \{0\}$, invertible by (iv), so its inverse is smooth by Cramer's rule, and homogeneous of degree $-2s$ by (i). $\square$

Theorem 8.3. *Let $s > 0$ and let A be the multiplier of an inertia symbol α of order $2s$. Then, with $m = Au$ and $w = \text{curl } m$:*

- (i) Proposition 2.2 holds verbatim, w obeying (4);
- (ii) Lemma 2.4 holds with $\hat{u}(k) = \alpha(k)^{-1}\hat{m}(k)$, so that $u = Sw$ and $\nabla u = Tw$ for multipliers smooth away from the origin and homogeneous of degrees $-1 - 2s$ and $-2s$, and (5) holds;
- (iii) $E = \langle Au, u \rangle$ is conserved and satisfies $C^{-1}\|w\|_{\dot{H}^{-1-s}}^2 \leq E \leq c^{-1}\|w\|_{\dot{H}^{-1-s}}^2$ with c, C as in Lemma 8.2;
- (iv) Lemma 3.1, Propositions 4.3, 4.5 and 6.1, Theorems 5.3, 6.2 and 6.3 hold verbatim.

In particular, if $p > 3/(2s)$ and $\int_0^T \|w\|_{L^p} dt < \infty$ with $T < \infty$, the solution does not break down at or before T . The constants depend on α only through c, C and finitely many derivatives of α on S^2 .

Proof. (i) is Proposition 2.2 unchanged: it is an identity in m and never refers to A . For (ii), $\hat{m}(k) = ik|k|^{-2}k \times \hat{w}(k)$ is the same computation as before, since it uses only $\hat{m}(k) \perp k$; composing with $\alpha(k)^{-1}$, which by Lemma 8.2 is smooth away from the origin and homogeneous of degree $-2s$, gives a symbol for S smooth away from the origin and homogeneous of degree $-1 - 2s$, and multiplication by ik_j raises the degree by one. A symbol of this kind divided by $|k|^{-1-2s}$, respectively $|k|^{-2s}$, is smooth and homogeneous of degree zero, hence a Mihlin multiplier, which is (5).

For (iii), conservation is the Lie–Poisson cancellation of Lemma 2.5, which uses only that A is symmetric. For the comparison, write $\beta(k) := |k|^{-2s}\alpha(k)$, so that the eigenvalues of $\beta(k)|_{k^\perp}$ lie in $[c, C]$. Mode by mode, $|\hat{w}| = |k||\hat{m}|$ and $\hat{m} = \alpha\hat{u}$ give

$$|k|^{-2-2s}|\hat{w}|^2 = |k|^{-2s}|\alpha\hat{u}|^2 = |k|^{2s}|\beta\hat{u}|^2, \quad \langle \alpha\hat{u}, \hat{u} \rangle = |k|^{2s}\langle \beta\hat{u}, \hat{u} \rangle,$$

and with $v := \beta\hat{u}$ the ratio of the second to the first is $\langle \beta^{-1}v, v \rangle / |v|^2 \in [C^{-1}, c^{-1}]$. Summing over $k \neq 0$ gives the claim; at $\alpha(k) = |k|^{2s}P_k$ one has $c = C = 1$ and recovers the equality of Lemma 2.5.

For (iv): the proof of Lemma 3.1 uses of the symbol of T only that it is smooth on $\mathbb{R}^3 \setminus \{0\}$ and homogeneous of degree $-2s$, which is (ii); the argument and the admissible range $p > 3/(2s)$ are unaltered. Section 4 uses the symbol only through (5), the remaining ingredients — Kato–Ponce, Calderón, Moser — being statements about Λ and the drift. In Proposition 4.5 the bounds $\|v\|_{L^2} \leq \|\delta\|_{L^2}$ and $\|\nabla v\|_{L^2} \leq \|\delta\|_{L^2}$ become $\leq c^{-1}\|\delta\|_{L^2}$, since the symbols of S and T are

bounded by c^{-1} on $|k| \geq 1$ for $s > 0$; only the constant in the Grönwall changes. Section 5 uses that P_N is self-adjoint and commutes with the operators in play, which holds for every Fourier multiplier, and otherwise only (5). Section 6 quotes the preceding statements and nothing else. $\square$

Remark 8.4 (the counts of Section 7 are unchanged). The dilation argument uses only that S is homogeneous of degree $-1 - 2s$, and the conserved E scales exactly as $\|w\|_{\dot{H}^{-1-s}}^2$ does by (iii), the comparison constants being scale-invariant. So $s^* = \frac{d}{2} + 1$ and the closing condition $s \geq \frac{5}{2}$ hold for the whole class, and the criterion is stated in a subcritical quantity throughout it. Proposition 7.1 likewise survives: it uses the two-dimensional forms of Proposition 2.2 and Lemma 3.1 only.

Remark 8.5 (where scalarity is used). Lemma 7.6 is the exception, and the reason is visible in its proof. For a scalar symbol $\alpha = a(k)P_k$ the contracted symbol is

$$D(k; \xi', \xi) = - \frac{(k \cdot \xi) \det(\xi, k, \xi')}{a(k) |k|^2},$$

so the metric enters through a scalar factor which cannot disturb the determinant, and the lemma holds for every scalar anisotropic symbol with $|D| \leq c^{-1} |k|^{-2s} |\xi' - \xi|$. For a genuinely matrix-valued α it fails, and not marginally: for the perturbation of `aniso_check.py` the ratio of $|D|$ at $\xi' = \xi$ to its generic size is 0.20, 0.16, 0.18 and 0.13 at $s = \frac{1}{4}, \frac{1}{2}, 1, 2$, against machine zero in the scalar case. So the formulation of Section 7 should be sharpened: the depletion is carried by the ε tensor in $w = \text{curl } m$ together with the scalarity of the inertia operator, and a matrix-valued metric destroys it while leaving every other statement of this paper intact.

Remark 8.6 (numerical check). `aniso_check.py` verifies the above on a fixed 24^3 grid for a scalar symbol $|k|^{2s}(1 + 0.6n_1^2 + 0.3n_2^4)$ and a matrix symbol, at $s = \frac{1}{4}, \frac{1}{2}, 1, 2$: symmetry of A to 10^{-15} with the sphere eigenvalues positive; the reduction (i) to 10^{-12} or better; the homogeneities in (ii) to 10^{-13} ; conservation of E to 10^{-19} , with the measured comparison constants of (iii) equal to $[0.625, 1]$ in the scalar case and $[0.741, 1.132]$ in the matrix case; and the alignment cancellation of Remark 8.5.

Remark 8.7 (no conservation law substitutes for the criterion). This class carries a conserved quantity besides E — the helicity $H(m) = \int m \cdot \text{curl } m$ of the momentum, a Casimir of the Lie–Poisson structure and so conserved for every inertia symbol — and it is worth recording that it cannot take the criterion’s place, for a reason that is specific to the matrix-valued case. By [26, Cor. 4.3] the value of H , and not merely its conservation, depends on the symbol only through the single number $\det(\alpha(k)|_{k^\perp})$ at each k . For a scalar symbol that number is $a(k)^2$ and determines α , so the point is moot. For a symbol of Definition 8.1 it is not: $\alpha(k)|_{k^\perp}$ is a positive definite form on a two-plane and carries three real parameters, of which the determinant pins one, leaving two per direction — the ratio of the eigenvalues and the orientation of the principal axes inside $k^\perp$ — to which the helicity is blind. The equation is not blind to them. Holding every determinant fixed and moving along those directions on a fixed grid, H is reproduced to machine precision while the production $\langle w, (w \cdot \nabla)u \rangle$, whose time integrability Theorem 6.3 converts into persistence through Lemma 3.1, varies by a factor of several [26, Prop. 6.4]. So on the anisotropic symbols the criterion is supplying control that no invariant of the structure supplies.

9. SCOPE

Everything in Sections 2–6 is proved, granted the quoted commutator estimates of Kato–Ponce [17] and Calderón [10], the Moser product estimate [18], and the kernel asymptotics of [27].

Solutions are those of Definition 5.1, weakly continuous into H^r by Lemma 5.2, which is what the arguments here require. Strong continuity into H^r also holds, by the reversibility of (4) and the argument of Bona and Smith [8]; it is omitted because nothing here uses it, not because it is out of reach. Lemma 3.1 is stated on $\mathbb{T}^3$, where the kernel is compactly supported and no far field arises; on $\mathbb{R}^3$ the argument needs an additive low-frequency term and the statement changes accordingly. The case $s = 0$ is excluded from Theorem 6.3 and is [4]; the case $s < 0$ is not treated. Sections 2–7 are written for the isotropic multipliers (1); Section 8 shows that all of them but Lemma 7.6 hold for any inertia symbol smooth away from the origin and homogeneous of degree $2s$, matrix-valued included, and that Lemma 7.6 needs the symbol to be scalar. Nothing here bears on the global regularity of the Navier–Stokes or Euler equations, and by the count of Section 7 nothing here can.

ACKNOWLEDGEMENTS

The direction of this work is the author’s: the question of how far the completeness threshold for the metrics (1) can be displaced, and the decision to press on that side rather than on the breakdown side, are his. That a continuation criterion was the blocking prerequisite — that no Beale–Kato–Majda statement exists for this family, and that supplying one was the available next step — was identified by Claude, a generative AI system developed by Anthropic, in the course of that work. The author drove the iterative process that strengthened the results. The derivations, the numerical verification reported in Sections 2, 3 and 7, and a first draft of this text were produced in collaboration with Claude, working under the author’s direction. The author is responsible for the contents of this paper, including any errors.

The scripts `bkm_check.py`, `kernel_check.py`, `depletion_check.py`, `scaling_check.py`, `dim2_check.py`, `deplete_ratio.py` and `aniso_check.py` underlying the computational statements are available from the author.

REFERENCES

- [1] V. I. Arnold, *Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l’hydrodynamique des fluides parfaits*, Ann. Inst. Fourier **16** (1966), 319–361.
- [2] H. Bahouri, J.-Y. Chemin and R. Danchin, *Fourier Analysis and Nonlinear Partial Differential Equations*, Grundlehren der mathematischen Wissenschaften **343**, Springer, Heidelberg, 2011.
- [3] M. Bauer, B. Kolev and S. C. Preston, *Geodesic completeness of the $H^{3/2}$ metric on $\text{Diff}(S^1)$* , Monatsh. Math. **193** (2020), 233–245. [arXiv:1904.12523](#).
- [4] J. T. Beale, T. Kato and A. Majda, *Remarks on the breakdown of smooth solutions for the 3-D Euler equations*, Comm. Math. Phys. **94** (1984), 61–66.
- [5] H. Brezis and T. Gallouët, *Nonlinear Schrödinger evolution equations*, Nonlinear Anal. **4** (1980), 677–681.
- [6] H. Brezis and S. Wainger, *A note on limiting cases of Sobolev embeddings and convolution inequalities*, Comm. Partial Differential Equations **5** (1980), 773–789.
- [7] M. Bauer, S. C. Preston and J. Valletta, *Breakdown of smooth solutions to the subcritical EPDiff equation*, Calc. Var. Partial Differential Equations **64** (2025). [arXiv:2503.19780](#).
- [8] J. L. Bona and R. Smith, *The initial-value problem for the Korteweg–de Vries equation*, Philos. Trans. Roy. Soc. London Ser. A **278** (1975), 555–601.
- [9] M. Bruveris and F.-X. Vialard, *On completeness of groups of diffeomorphisms*, J. Eur. Math. Soc. **19** (2017), 1507–1544. [arXiv:1403.2089](#).
- [10] A. P. Calderón, *Commutators of singular integral operators*, Proc. Nat. Acad. Sci. U.S.A. **53** (1965), 1092–1099.
- [11] D. Chae, P. Constantin and J. Wu, *Inviscid models generalizing the two-dimensional Euler and the surface quasi-geostrophic equations*, Arch. Ration. Mech. Anal. (2011), doi:10.1007/s00205-011-0411-5. [arXiv:1010.1506](#).
- [12] P. Constantin and C. Fefferman, *Direction of vorticity and the problem of global regularity for the Navier–Stokes equations*, Indiana Univ. Math. J. **42** (1993), 775–789.

- [13] D. G. Ebin, G. Misiołek and S. C. Preston, *Singularities of the exponential map on the volume-preserving diffeomorphism group*, Geom. Funct. Anal. **16** (2006), 850–868.
- [14] D. G. Ebin and J. Marsden, *Groups of diffeomorphisms and the motion of an incompressible fluid*, Ann. of Math. **92** (1970), 102–163.
- [15] T. M. Elgindi, *Osgood’s lemma and some results for the slightly supercritical 2D Euler equations for incompressible flow*, Arch. Ration. Mech. Anal. (2014), doi:10.1007/s00205-013-0691-z. [arXiv:1308.1155](#).
- [16] J. Escher and B. Kolev, *Geodesic completeness for Sobolev H^s -metrics on the diffeomorphisms group of the circle*, J. Evol. Equ. **14** (2014), 949–968. [arXiv:1308.3570](#).
- [17] T. Kato and G. Ponce, *Commutator estimates and the Euler and Navier–Stokes equations*, Comm. Pure Appl. Math. **41** (1988), 891–907.
- [18] A. J. Majda and A. L. Bertozzi, *Vorticity and Incompressible Flow*, Cambridge Texts in Applied Mathematics **27**, Cambridge University Press, 2002.
- [19] G. Misiołek and S. C. Preston, *Fredholm properties of Riemannian exponential maps on diffeomorphism groups*, Invent. Math. **179** (2010), 191–227.
- [20] E. Olson and E. S. Titi, *Viscosity versus vorticity stretching: global well-posedness for a family of Navier–Stokes-alpha-like models*, Nonlinear Anal. **66** (2007), no. 11, 2427–2458.
- [21] S. C. Preston, *On the volumorphism group, the first conjugate point is always the hardest*, Comm. Math. Phys. **267** (2006), 493–513.
- [22] S. C. Preston, *The WKB method for conjugate points in the volumorphism group*, Indiana Univ. Math. J. **57** (2008), 3303–3327. [arXiv:0710.3870](#).
- [23] S. C. Preston, *A geometric rigidity theorem for hydrodynamical blowup*, Comm. Partial Differential Equations **35** (2010), 2007–2020.
- [24] S. C. Preston and A. Sarria, *Lagrangian aspects of the axisymmetric Euler equation*, [arXiv:1505.05569](#).
- [25] A. Salkimbayev, *Rigidity of the Euler–Arnold form under energy-preserving deformation*, companion paper.
- [26] A. Salkimbayev, *Momentum helicity separates cascade operators from Euler–Arnold operators*, companion paper.
- [27] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.
- [28] T. Tao, *Finite time blowup for Lagrangian modifications of the three-dimensional Euler equation*, Ann. PDE **2** (2016), no. 9. [arXiv:1606.08481](#).
- [29] K. Yamazaki, *On the global regularity of generalized Leray-alpha type models*, Nonlinear Anal. **75** (2012), 503–515.
- [30] A. Salkimbayev, *Eight conditions on the missing estimate for the three-dimensional Navier–Stokes equations*, companion paper.

ALMATY, KAZAKHSTAN

Email address: adilsalkimbayev@gmail.com