

CROSS-TRIAD RIGIDITY FOR CASCADE OPERATORS

ADILKHAN SALKIMBAYEV

ABSTRACT. Paper II shows that on any single triad a sum of cascade terms reproduces an Euler–Arnold vertex, so what remains of the sum question is whether one fixed coefficient set can do so on *all* triads at once. For a two-species cascade this is a gauge-trivialisation problem, which we answer negatively for every scalar inertia multiplier. A first invariant — the ratios of the blocks’ Cayley hyperdeterminants — excludes all but two exponents, $s = 0$ and $s = -2$, which are Euler and its vorticity dual. We show that this degeneracy is structural, by proving that on the relevant stratum the hyperdeterminant reduces to the product of the block’s four entries, and that it is nevertheless only a weakness of that invariant: the correct condition, that all the vertices lie in a single $GL(2)^3$ orbit, fails at $s = 0$ and $s = -2$ as well. The proof turns on an invariant *condition* rather than an invariant function — the rank of a flattening — which is blind to the degeneracy that silences the hyperdeterminant, and settles all scalar multipliers at once.

1. THE QUESTION

By Paper II, Theorem 5.7 every Euler–Arnold vertex on a triad is a sum of at most eight tensors with one rank-one block each, so no triad-local argument can exclude sums. What that construction does not produce is a cascade operator in Tao’s sense: its terms are built from the vertex itself and change from triad to triad, whereas a local cascade operator reuses one finite coefficient set on every triad and at every scale. The contribution of a cascade with m species to a triad $k + l + m = 0$ is

$$\sum_{i_1, i_2, i_3} \alpha_{i_1 i_2 i_3} \hat{\psi}_{i_1}(k) \otimes \hat{\psi}_{i_2}(l) \otimes \hat{\psi}_{i_3}(m),$$

the α constant, the divergence-free profiles $\hat{\psi}_i(k) \in k^\perp$ varying with the wavevector.

Question 1.1. *Do there exist m , profiles, constants α , and an inertia multiplier A with this equal to $T_A(k, l, m)$ on every triad of the support?*

For $m = 2$ the profiles form a frame G_k of each perp-plane and the question becomes a trivialisation,

$$T_A(k, l, m) = (G_k \otimes G_l \otimes G_m)\alpha, \quad \alpha \text{ independent of the triad}, \quad (1)$$

equivalently: all the vertices lie in one orbit of $GL(2)^3$ acting on $\mathbb{R}^{24}$.

2. A FIRST INVARIANT, AND WHERE IT FAILS

Proposition 2.1. *If (1) holds then the ratios $\text{Det}(T_0) : \text{Det}(T_1) : \text{Det}(T_2)$ of the blocks’ Cayley hyperdeterminants are the same on every triad.*

Proof. $\text{Det}((g_1 \otimes g_2 \otimes g_3)t) = (\det g_1 \det g_2 \det g_3)^2 \text{Det}(t)$, and block j carries its slots on the legs $j, j + 1, j + 2$, that is on all three legs. So every block is acted on by the same triple and each $\text{Det}(T_j)$ picks up the same factor. $\square$

Proposition 2.2 (numerical). *The ratios are not triad-independent. For $A = |k|^s I$ with $s \notin \{0, -2\}$, and for the anisotropic multiplier $A_k = \text{diag}(1, 1 + |k|)$, the hyperdeterminants are of order 10^{-3} to 10^{-2} relative to $\|T\|^4$ and $\text{Det}(T_0)/\text{Det}(T_2)$ varies: for $s = -3$ it takes the values 0.418, 0.827, 1.101, 0.743, 1.507 at radii $(1, 1.1, 1.2)$, $(1, 1.3, 1.6)$, $(1, 1.5, 1.8)$, $(1, 1.2, 2.0)$, $(1, 1.9, 2.4)$. Exactly two exponents escape: at $s = 0$ and $s = -2$, and for every constant $A = cI$, all three hyperdeterminants vanish identically, to 10^{-17} relative.*

3. THE DEGENERACY IS STRUCTURAL

For scalar A the operators P_i, Q_i are scalars, so a block is supported on the four monomials of $S^A \cup S^B$. Write its entries

$$p = t_{111}, \quad q = t_{122}, \quad r = t_{212}, \quad w = t_{221}.$$

Lemma 3.1. *On this stratum $\text{Det}(t) = 4pqrw$.*

Proof. In Cayley's expression every monomial contains one of $t_{112}, t_{121}, t_{211}, t_{222}$ except the single term $4t_{111}t_{122}t_{212}t_{221}$. $\square$

Theorem 3.2. *Let $A = f(|k|)I$. Then all three hyperdeterminants vanish on every triad if and only if f is constant or $f(x) = c/x^2$; that is, $s = 0$ or $s = -2$.*

Proof. With $c^A = u_l/(u_k u_m)$ and $c^B = u_k/(u_l u_m)$, $u_i = \mu_i^{1/2}$, and the coefficients of Paper I, Lemma 5.7,

$$q = \frac{1}{2}c^A\gamma, \quad r = \frac{1}{2}c^B\gamma', \quad w = \frac{J}{2z}(c^B - c^A), \quad p = \frac{J}{2z}\left(c^A\frac{y}{x} - c^B\frac{x}{y}\right).$$

Since $\gamma = -J/x$ and $\gamma' = J/y$ are non-zero on a non-degenerate triad, q and r never vanish, so by Lemma 3.1 the hyperdeterminant vanishes exactly when $p = 0$ or $w = 0$. Now $c^A/c^B = \mu_l/\mu_k$, so $w = 0$ on all triads iff $f(y) = f(x)$ for all admissible x, y , i.e. f is constant; and $p = 0$ on all triads iff $f(y)y^2 = f(x)x^2$, i.e. $f(x)x^2$ is constant. $\square$

So the two exceptions are not an accident of the sample: they are exactly the two inertia operators for which one of the four entries of every block vanishes identically. The two are exchanged by a duality fixed at $s = -1$, where the block is self-dual, its entries satisfying $p = w$ and $q = -r$.

4. BUT THE INVARIANT'S SILENCE IS AN ACCIDENT

Lemma 3.1 also shows why Proposition 2.1 is weak here: on this stratum the hyperdeterminant is merely a product of coordinates, and it reports nothing beyond whether all four are non-zero. The natural next invariants are no better. Pairing two blocks with the alternating form, $\langle X, Y \rangle_\varepsilon := \varepsilon^{aA} \varepsilon^{bB} \varepsilon^{cC} X_{abc} Y_{ABC}$ after aligning both by leg, is a relative invariant of weight one in each $\det G$, so ratios of such pairings would be absolute invariants. They vanish identically: the four monomials above are exactly those of even parity, ε couples even parity to odd, and so every such pairing is zero. This is the parity grading of Paper I, Lemma 5.9 reappearing.

A proxy does work, however. It is not an invariant *function* but an invariant *condition*, and it is blind to exactly the degeneracy that silences the others: the rank of a flattening.

Lemma 4.1. *If $X = (g_1 \otimes g_2 \otimes g_3)Y$ with each g_i invertible, then in each of the three slots the flattenings of X and Y have the same rank.*

Proof. Writing M_j for the 2×4 flattening in slot j , $M_j(X) = g_j^\top M_j(Y) (g_{j+1} \otimes g_{j+2})$, and both factors are invertible. $\square$

Lemma 4.2. *Let $A = f(|k|)I$ with $f > 0$, and let τ be a non-degenerate triad. The block with its slots on the legs (k, l, m) has third flattening of rank one when $|k| = |l|$ and of rank two otherwise — for every f .*

Proof. By Theorem 3.2 the block is supported on the four entries p, q, r, w , and $q = \frac{1}{2}c^A\gamma$, $r = \frac{1}{2}c^B\gamma'$ with $\gamma = -J/x$ and $\gamma' = J/y$ non-zero on a non-degenerate triad, so $q, r \neq 0$. The slot-3 flattening has rows

$$(t_{111}, t_{121}, t_{211}, t_{221}) = (p, 0, 0, w), \quad (t_{112}, t_{122}, t_{212}, t_{222}) = (0, q, r, 0),$$

whose supports are disjoint. The second row never vanishes, so the rank is one exactly when $p = w = 0$. By the computation in Theorem 3.2, $w = 0$ iff $f(x) = f(y)$ and $p = 0$ iff $f(x)x^2 = f(y)y^2$; together these give $x^2 = y^2$, hence $x = y$. Conversely $x = y$ gives both. $\square$

Theorem 4.3. *Let $A = f(|k|)I$ be any positive scalar multiplier. If the support contains a non-degenerate triad two of whose legs have equal length, and a non-degenerate triad whose three legs have pairwise distinct lengths, then Question 1.1 has a negative answer with $m = 2$.*

Proof. Suppose (1) holds. The three cyclic blocks at a triad are $(G \otimes G \otimes G)$ applied to the three cyclic relabellings of the single tensor α , and the frames are invertible, so by Lemma 4.1 the multiset of flattening ranks over the three blocks of a triad is the same at every triad of the support. By Lemma 4.2 the number of those blocks whose third flattening has rank one is the number of cyclically adjacent pairs of legs of equal length: it is one at the first triad and zero at the second. $\square$

This settles every scalar multiplier at once, the two exceptional exponents included — precisely where every hyperdeterminant and every ε pairing vanishes. What defeated the three earlier attempts was that each looked for an invariant function; a rank is not one.

The sharp condition is still the orbit one, and it is worth testing directly, both because it is sharper than any single necessary condition and because it does not need two triads of different shape.

Proposition 4.4 (numerical). *Let $A = f(|k|)I$ be a scalar multiplier. At a triad let V be the span of the twelve orbit directions $E \cdot T$, $E \in \mathfrak{gl}(2)^3$, together with the three directions $\partial T / \partial \mu_i$ — the latter included so that one computation covers every f at once, only the three values of f at the leg lengths entering. Then the shape derivatives $\partial T / \partial y$, $\partial T / \partial z$ do not lie in V . Their relative residuals after projection onto V , at radii $(1, 1.5, 1.8)$, are 0.065 and 0.055 for $s = 0$; 0.39 and 0.54 for $s = -2$; and between 0.15 and 0.35 for $s = -1$, $s = 3/2$ and $s = 3$.*

For a fixed f the three μ directions are not available and the test is correspondingly sharper. At $f \equiv 1$, where A stays literally constant as the shape moves, the residual off the twelve orbit directions alone runs from 0.099 to 0.855 over twelve shapes, and is stable to three significant figures across step sizes 10^{-4} , 10^{-5} , 10^{-6} of a five-point derivative.

Corollary 4.5. *For every scalar inertia multiplier, including Euler and the vorticity form, Question 1.1 has a negative answer with $m = 2$. Theorem 4.3 proves this whenever the support carries two triads of different shape; Proposition 4.4 is an independent check, and covers a single triad of fixed shape as well.*

Remark 4.6 (the Euler case). Euler satisfies three of the necessary conditions established in this paper. Its hyperdeterminants vanish identically (Theorem 3.2); the web condition holds exactly on every shared-leg pair (Proposition 6.2); and it carries canonical lines (Proposition 7.1). Taken together that looks like membership of the class it is supposed to be excluded from, and it is worth saying plainly why it is not. Those three conditions are necessary, not

sufficient, and each of the three sections says in its own terms that the condition is weak — the hyperdeterminant because on this stratum it is a product of coordinates, the web because its conditions are massively redundant, the lines because a gauge absorbs them. Satisfying a weak condition is not evidence of membership. The orbit test is the sharp one, and Euler fails it by a wide margin once A is held fixed, which for a constant f is the correct question.

The figure of 2.4×10^{-5} that an earlier version of this remark recorded at radii (1, 1.3, 1.6) was measured in the relaxed span including the three μ directions; it is a statement about the freedom in f , not about the Euler vertex, whose residual at that shape is 0.53. The recomputation used an orbit tangent built by differentiating the group action directly, which agrees with the construction used elsewhere here to 1.6×10^{-11} .

Remark 4.7. Residuals are reported rather than ranks throughout, because a rank count needs a tolerance and some of these residuals are small enough to make a fixed-tolerance rank unreliable.

Remark 4.8. The computation was checked three ways: the orbit directions lie in the kernel of the energy relations of Paper II, Theorem 5.1 to 5×10^{-17} , as they must since the gauge action preserves $\langle T(u, u), u \rangle$; a finite group motion lies in their span to the accuracy of the difference quotient; and the shape derivative lies in that kernel too.

5. THE SINGLE-TRIAD METHOD IS EXHAUSTED

Proposition 4.4 does not extend to matrix-valued multipliers, and the reason is not that the test was run too coarsely.

Proposition 5.1 (numerical). *Let A_k be a full positive definite operator on $k^\perp$, so that the admissible span is the twelve orbit directions together with the nine directions $\partial T / \partial A_i$. Then the shape derivatives lie in that span, to relative residual below 10^{-11} , for every inertia operator tested — scalar, diagonal anisotropic, fully anisotropic, and 200 random positive definite triples — and at both triads. The span has dimension 14 inside the 16-dimensional $\ker L$, so this is not a case of the admissible directions exhausting the ambient space.*

So with three functions per wavevector instead of one, the gauge freedom and the multiplier together absorb the shape variation of the vertex exactly, to first order. On a single triad there is nothing left for an argument of this kind to obstruct, and no refinement of the present one will recover an obstruction. That is a statement about shape derivatives and not about the images: for $m = 2$ a single triad does carry conditions, though not ones this method can see, and Remark 8.2 names two of them.

6. THE WEB

In both cases what defeats the argument is that a single triad sees each leg once. The information not used is that G_k and A_k are functions of the wavevector, shared by every triad through k . For $m = 2$ that sharing yields a condition in closed form, and one that holds for a matrix multiplier as well.

Lemma 6.1. *Suppose $X = (G_k \otimes G_l \otimes G_m)\alpha$ with $m = 2$. Then for $v \in k^\perp$ the slice $X(v) := X(v, \cdot, \cdot)$ satisfies $X(v) = G_l \alpha(G_k^\top v) G_m^\top$, so*

$$\det X(v) = \det G_l \det G_m \cdot q_\alpha(G_k^\top v), \quad q_\alpha(w) := \det \alpha(w, \cdot, \cdot).$$

The triad enters only through a scalar. Hence for any two triads τ, τ' sharing the leg k , the binary quadratic forms $Q_\tau(v) := \det T_A(\tau)|_v$ and $Q_{\tau'}$ on $k^\perp$ are proportional.

This is the first necessary condition available for a matrix-valued multiplier, Proposition 5.1 having closed the single-triad route. We tested it on a web: legs the 28 non-zero points of a planar lattice ball of radius 3, triads the 60 non-degenerate triples summing to zero, giving 395 pairs of triads with a common leg.

Proposition 6.2 (numerical). *The condition excludes generic multipliers. The proportion of shared-leg pairs passing is 0% for independent random positive definite A_k , with minimum separation 1.7×10^{-3} over 516 pairs; 15% for $A_k = \text{diag}(1, 1 + |k|)$; and 13% for $A = |k|^{3/2}I$. For $A = I$ and $A = |k|^{-2}I$ it holds exactly, on every pair.*

The naive count is encouraging and wrong. Proportionality of two binary quadratics is two conditions, so the web carries $\sum_k 2(n_k - 1) = 304$ of them against $3|S| = 84$ free parameters; conditions grow like $|S|^2$ and parameters like $|S|$. That does not survive contact with the computation.

Proposition 6.3 (numerical). *The conditions are massively redundant. Linearising the residual in the 84 parameters at $A = I$, the Jacobian has rank 26, so the local solution set is 58-dimensional; at $A = |k|^{-2}I$ the rank is 43 and the solution set is 41-dimensional. The web condition therefore does not isolate the two exceptional multipliers, and does not settle the matrix case.*

Proposition 6.4 (numerical). *The redundancy has the same source as every previous escape. At $A = I$ and $A = |k|^{-2}I$ each Q_τ has vanishing discriminant — to 0 and 5×10^{-15} respectively — so it is a perfect square ℓ_τ^2 ; and the line ℓ_τ is the same for every triad through a given leg, the directions agreeing to 3×10^{-16} . For $A = |k|^{3/2}I$ the discriminant is of order one and the directions spread maximally. Proportionality $\ell^2 \propto \ell'^2$ is therefore one condition rather than two, and it is satisfied automatically.*

Remark 6.5. That Q_τ is a perfect square is Theorem 3.2 again: the discriminant of the slice quadratic is the Cayley hyperdeterminant of the block, which vanishes identically exactly at $s = 0$ and $s = -2$. The three failures recorded in this paper — the hyperdeterminant ratios, the ε pairings, and now the web — are one failure, and it is that the blocks of the Euler and vorticity vertices are degenerate as $2 \times 2 \times 2$ tensors.

Remark 6.6. One positive fact emerges. For $A = I$ and $A = |k|^{-2}I$ every leg k carries a canonical line in $k^\perp$, the repeated root of Q_τ , and it does not depend on which triad through k is used to compute it. Any further attempt should either build on that line or explain why it cannot separate; it is the only triad-independent structure these two vertices have been found to carry.

7. THE CANONICAL LINES, AND WHY THEY DO NOT HELP

The last remark of Section 6 isolated the only triad-independent structure these two vertices carry. It can be identified, and it is richer than a single line.

Proposition 7.1 (numerical). *At $A = I$ every leg carries two distinct canonical lines: slicing a block in either of its input slots gives the plane normal ν , while slicing in the output slot gives $\hat{k}$. The discriminants vanish to 10^{-15} , and the two directions are exactly 90° apart, on every triad and at every leg. At $A = |k|^{-2}I$ all three slots give $\hat{k}$: a single line.*

If (1) holds, the lines at k are G_k applied to the corresponding canonical lines of α , which are fixed. So G_k is pinned to the stabiliser of the observed configuration: the diagonal torus, two parameters, at $A = I$; the Borel, three, at $A = |k|^{-2}I$; in place of the four of $GL(2)$. This is a real rigidification, and it is worth nothing.

Proposition 7.2 (numerical). *On a single triad the shape derivatives of T_A lie in the span of the nine derivatives $\partial T_A / \partial A_i$ alone, with no gauge directions at all. The relative residuals are below 3×10^{-10} for every inertia operator tested — $A = I$, $|k|^{-2}I$, $|k|^{3/2}I$, fully anisotropic — and at both triads, while that span has dimension only 8 or 9 inside the 16-dimensional $\ker L$. The statement is not vacuous: random vectors of $\ker L$ have residuals between 0.25 and 0.90 off the same span.*

Corollary 7.3. *Rigidifying the gauge cannot obstruct a matrix multiplier. Cutting the twelve gauge directions to the six allowed by Proposition 7.1 leaves the shape derivative absorbed to 1.4×10^{-12} , and cutting to the nine of the Borel case leaves it absorbed to 6×10^{-12} .*

Remark 7.4. The reason is now plain, and it is not about the gauge at all. Proposition 7.2 says that changing the shape of a triad is, to first order, the same as adjusting the inertia operator at its three legs. The multiplier freedom already covers the shape variation, so no restriction placed on G_k — by canonical lines or by anything else — can create an obstruction that a single triad is able to see. With A held fixed the picture is the opposite: the shape derivative then escapes the full twelve-dimensional gauge span with residuals between 0.39 and 0.71, which is Proposition 4.4 in another guise.

8. THE EXISTENCE QUESTION

A correction first. We said above that a further attempt must compare triads that do not share their multiplier data. That is backwards. Triads with disjoint legs carry independent A , so by Proposition 7.2 each absorbs its own shape and nothing is obstructed; rigidity can only come from triads that *do* share legs. That is what a web provides.

The question is also not undecidable. “There exist A , α and G such that $T_A(\tau) = (G \otimes G \otimes G)\alpha$ for every τ in a finite web” is a first-order sentence over the reals, and Tarski–Seidenberg decides such a sentence in principle. What is out of reach is the elimination, not the logic. So the question is one of measurement, and the quantity to measure is the dimension of two images: that of $\{A_k\} \mapsto (T_A(\tau))_\tau$, and that of $(\alpha, \{G_k\}) \mapsto ((G \otimes G \otimes G)\alpha)_\tau$. A representation exists only if they meet.

Both live in $\ker L$ on every triad, and the ambient is therefore $16|T|$ and not $24|T|$. For the vertex that is Paper II, Theorem 5.1. For the cascade it is a condition on α : summing the three blocks in one fixed slot order gives

$$(B_0)_{abc} + (B_1)_{bca} + (B_2)_{cab} = \sum_{ijp} (\alpha_{ijp} + \alpha_{jpi} + \alpha_{pij}) G_k(i, a) G_l(j, b) G_m(p, c),$$

which vanishes for every frame exactly when the cyclic symmetrisation of α does — Tao’s condition (4.3), and Paper II, Corollary 5.4. So α ranges not over $\mathbb{R}^{m^3}$ but over the kernel of that symmetrisation, of dimension m^3 less the number of cyclic orbits, that is $\frac{2}{3}(m^3 - m)$: four, sixteen and forty for $m = 2, 3, 4$.

Proposition 8.1 (numerical). *Measured by Jacobian rank at random points, on the planar lattice webs of radius 2 and 3:*

web	ambient $16 T $	$m = 2$	$m = 3$	$m = 4$
$R = 2$ (12 legs, 8 triads)	128	80	111	152
$R = 3$ (28 legs, 60 triads)	960	194	257	330

the entries being $\dim V + \dim C$. The vertex image attains its parameter count exactly, $\dim V = 3|S|$; the cascade image falls short of $\frac{2}{3}(m^3 - m) + 2m|S|$ only by the dimension of the gauge stabiliser. The codimension of the pair is 48, 17, -24 at $R = 2$ and 766, 703, 630 at $R = 3$.

Both dimensions grow linearly in $|S|$ for every fixed m , while the ambient grows like $|T| \sim |S|^2$. The deficit therefore diverges, and it does so for Tao's $m = 4$ as for $m = 2$ — but the small web is too small to see it. At $m = 4$ its parameters already over-saturate its ambient, and there the count says nothing at all; one has to go to $R = 3$, where the codimension is 630. This is the counting of Section 6 finally doing what was claimed for it there — and doing it because, unlike the proportionality conditions of Proposition 6.2, these images suffer no redundancy collapse.

Remark 8.2. What this does and does not establish. Two subvarieties whose dimensions sum to less than the ambient cannot meet in general position, but may still meet in special position, and a dimension count does not exclude that. So Proposition 8.1 is not a proof. It is, however, the strongest form the evidence can take short of elimination: the deficit is not marginal, it is not an artefact of redundancy, it grows without bound, and it is insensitive to the species count.

On one triad the two images meet for $m \geq 3$: there the cascade image fills $\ker L$, measured 16 of 16 at $m = 3$ and at $m = 4$, so a single triad imposes no condition. At $m = 2$ it does not. The image has dimension 10 inside the 16-dimensional $\ker L$, and two of the six conditions can be named. Writing $B_1 = (X \otimes Y \otimes Z)B_0$ with $X = G_l G_k^{-1}$, $Y = G_m G_l^{-1}$, $Z = G_k G_m^{-1}$ forces $ZYX = I$, hence $\det X \det Y \det Z = 1$, and since Det has weight two in each determinant,

$$\text{Det}(T_0) = \text{Det}(T_1) = \text{Det}(T_2) \quad \text{exactly, on every triad.}$$

This is sharper than Proposition 2.1, which compares triads and allows an arbitrary factor at each. It uses the invertibility of the frames, so it is special to $m = 2$; for $m \geq 3$ the frames are $2 \times m$ and no such relation survives. So for the species counts that matter for Tao it is indeed the web that separates the two classes, and the separation widens as the web grows; at $m = 2$ the separation already begins on a single triad.

9. CONCLUSION

Part of the sum question is settled and part is not, and the parts should not be run together.

Triad-locally it is settled, and settled against exclusion: by Paper II, Theorem 5.7 every Euler–Arnold vertex is a sum of at most eight admissible tensors, so no triad-local argument can exclude sums. That is a proof.

For two species and a scalar multiplier it is settled, and settled by proof: Theorem 4.3. That argument uses the invertibility of the frames, which is exactly the condition $m = 2$, and it needs two triads of different shape in the support.

Beyond that we have *not* proved that a local cascade operator cannot be an Euler–Arnold operator. What is proved is a sequence of statements about methods, each closing a route rather than the question. No linear functional separates the two classes, the only ones vanishing on the Euler–Arnold class being the energy identity that every cascade satisfies by construction (Paper II, Corollary 5.4). No invariant *function* of a single block separates them, the blocks at the two surviving inertia operators being degenerate as $2 \times 2 \times 2$ tensors (Theorem 3.2) — an invariant *condition* does, which is Theorem 4.3, but only at $m = 2$. No rigidification of the gauge separates them, since on one triad the inertia operator can imitate a change of shape (Proposition 7.2). Alongside these, Proposition 8.1 measures on a web a codimension that grows quadratically in the number of legs, for every species count.

That last is evidence and not proof, for the reason given in Remark 8.2: two subvarieties whose dimensions sum to less than the ambient need not be disjoint, and nothing here excludes an intersection in special position. So the position to quote is that every separating method we could construct fails, that the two classes are measured far apart, and that the distance between “far apart in dimension” and “disjoint” is exactly what remains.

Closing it is a question in real algebraic geometry rather than in fluid mechanics, and we stop here.

REFERENCES

- [1] A. Salkimbayev, *Rigidity of the Euler–Arnold form under energy-preserving deformation*, companion paper.
- [2] A. Salkimbayev, *The isotropic directional obstruction for cascade operators*, companion paper.
- [3] T. Tao, *Finite time blowup for an averaged three-dimensional Navier–Stokes equation*, J. Amer. Math. Soc. **29** (2016), 601–674.

ALMATY, KAZAKHSTAN

Email address: `adilsalkimbayev@gmail.com`