



National Comprehensive  
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

# Central Nervous System Cancers

Version 3.2020 — September 11, 2020

[NCCN Guidelines for Patients®](#)

**Continue**



National  
Comprehensive  
Cancer  
Network®

# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

[NCCN Guidelines Index](#)  
[Table of Contents](#)  
[Discussion](#)

**\*Louis Burt Nabors, MD/Chair Ψ**  
O'Neal Comprehensive Cancer Center at UAB

**\*Jana Portnow, MD/Vice-Chair † Ψ**  
City of Hope National Medical Center

**Manmeet Ahluwalia, MD ‡ † Ψ**  
Case Comprehensive Cancer Center/  
University Hospitals Seidman Cancer Center  
and Cleveland Clinic Taussig Cancer Institute

**Joachim Baehring, MD Ψ**  
Yale Cancer Center/Smilow Cancer Hospital

**Henry Brem, MD ¶ Ψ**  
The Sidney Kimmel Comprehensive  
Cancer Center at Johns Hopkins

**Steven Brem, MD ¶**  
Abramson Cancer Center  
at the University of Pennsylvania

**Nicholas Butowski, MD Ψ †**  
UCSF Helen Diller Family  
Comprehensive Cancer Center

**Jian L. Campian, MD, PhD Ψ †**  
Siteman Cancer Center at Barnes-  
Jewish Hospital and Washington  
University School of Medicine

**Stephen W. Clark, MD, PhD Ψ**  
Vanderbilt-Ingram Cancer Center

**Andrew J. Fabiano, MD ¶**  
Roswell Park Comprehensive Cancer Center

**Peter Forsyth, MD Ψ**  
Moffitt Cancer Center

**Jona Hattangadi-Gluth, MD §**  
UC San Diego Moores Cancer Center

**NCCN**  
**Mary Anne Bergman**  
**Susan Darlow, PhD**

**\*Matthias Holdhoff, MD, PhD †**  
The Sidney Kimmel Comprehensive  
Cancer Center at Johns Hopkins

**\*Craig Horbinski, MD, PhD ≠**  
Robert H. Lurie Comprehensive Cancer  
Center of Northwestern University

**Larry Junck, MD Ψ**  
University of Michigan Rogel Cancer Center

**\*Thomas Kaley, MD Ψ**  
Memorial Sloan Kettering Cancer Center

**\*Priya Kumthekar, MD Ψ**  
Robert H. Lurie Comprehensive Cancer  
Center of Northwestern University

**Jay S. Loeffler, MD § Ψ**  
Massachusetts General Hospital Cancer Center

**\*Maciej M. Mrugala, MD, PhD, MPH Ψ †**  
Mayo Clinic Cancer Center

**\*Seema Nagpal, MD † Ψ**  
Stanford Cancer Institute

**Manjari Pandey, MD †**  
St. Jude Children's  
Research Hospital/  
The University of Tennessee  
Health Science Center

**Ian Parney, MD, PhD Ψ ¶**  
Mayo Clinic Cancer Center

**Katherine Peters, MD, PhD Ψ**  
Duke Cancer Institute

**\*Vinay K. Puduvalli, MD Ψ**  
The Ohio State University Comprehensive  
Cancer Center - James Cancer Hospital  
and Solove Research Institute

**Ian Robins, MD, PhD †**  
University of Wisconsin  
Carbone Cancer Center

**Jason Rockhill, MD, PhD §**  
Fred Hutchinson Cancer Research Center/  
Seattle Cancer Care Alliance

**\*Chad Rusthoven, MD §**  
University of Colorado Cancer Center

**Nicole Shonka, MD † Ψ**  
Fred and Pamela Buffet Cancer Center

**Dennis C. Shrieve, MD, PhD §**  
Huntsman Cancer Institute  
at the University of Utah

**\*Lode J. Swinnen, MB, ChB ‡**  
The Sidney Kimmel Comprehensive  
Cancer Center at Johns Hopkins

**Stephanie Weiss, MD §**  
Fox Chase Cancer Center

**\*Patrick Yung Wen, MD Ψ**  
Dana-Farber Cancer Institute

**Nicole E. Willmarth, PhD ≠**  
American Brain Tumor Association

[NCCN Guidelines Panel Disclosures](#)

‡ Hematology/Hematology oncology	¥ Patient advocacy
¶ Internal medicine/Primary care	§ Radiotherapy/Radiation oncology
† Medical oncology	¶ Surgery/Surgical oncology
Ψ Neurology/Neuro-oncology	* Discussion Section Writing Committee
≠ Pathology	

**Continue**



### [NCCN Central Nervous System Cancers Panel Members](#) [Summary of the Guidelines Updates](#)

[Adult Low-Grade \(WHO Grade I or II\) Glioma/Pilocytic and Infiltrative Supratentorial Astrocytoma/Oligodendroglioma \(ASTR-1\)](#)  
[Anaplastic Gliomas/Glioblastoma \(GLIO-1\)](#)  
[Adult Intracranial and Spinal Ependymoma \(Excluding Subependymoma\) \(EPEN-1\)](#)  
[Adult Medulloblastoma \(AMED-1\)](#)  
[Primary CNS Lymphoma \(PCNS-1\)](#)  
[Primary Spinal Cord Tumors \(PSCT-1\)](#)  
[Meningiomas \(MENI-1\)](#)  
[Limited Brain Metastases \(LTD-1\)](#)  
[Extensive Brain Metastases \(MU-1\)](#)  
[Leptomeningeal Metastases \(LEPT-1\)](#)  
[Metastatic Spine Tumors \(SPINE-1\)](#)

#### Principles of:

- [Brain and Spine Tumor Imaging \(BRAIN-A\)](#)
- [Brain Tumor Surgery \(BRAIN-B\)](#)
- [Radiation Therapy for Brain and Spinal Cord \(BRAIN-C\)](#)
- [Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\)](#)
- [Brain and Spine Tumor Management \(BRAIN-E\)](#)
- [Brain Tumor Pathology \(BRAIN-F\)](#)

**Clinical Trials:** NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN Member Institutions, [click here: nccn.org/clinical\\_trials/member\\_institutions.aspx](#).

**NCCN Categories of Evidence and Consensus:** All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

**NCCN Categories of Preference:** All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2020.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Updates in Version 3.2020 of the NCCN Guidelines for Central Nervous System Cancers from Version 2.2020 include:

### MS-1

- The Discussion section was updated to reflect the changes in the algorithm.

Updates in Version 2.2020 of the NCCN Guidelines for Central Nervous System Cancers from Version 1.2020 include:

### BRAIN-D (8 of 15)

- *Tucatinib + trastuzumab + capecitabine was added as a category 2A recommendation for HER2-positive breast cancer patients previously treated with 1 or more anti-HER2-based regimens.*
  - ▶ The following reference was included: *Murthy RK, Loi S, Okines A, et al. Tucatinib, trastuzumab, and capecitabine for HER2-positive metastatic breast cancer. N Engl J Med. 2020;382:597-609.*

Updates in Version 1.2020 of the NCCN Guidelines for Central Nervous System Cancers from Version 3.2019 include:

### ASTR-1

#### Adjuvant Treatment

- *Consider BRAF and MEK inhibitors if BRAF V600E activating mutation*, is new to the page for pilocytic astrocytoma, PXA, ganglioglioma.

#### Follow-Up

- Modified, Brain MRI every 3–6 mo for 3–5 y then at least *annually every 6–12 mo or* as clinically indicated.

#### Footnotes

- "c": Modified to include, *and BRAF V600E mutation*.

### ASTR-2

- Algorithm has been streamlined.

- ▶ RT + adjuvant temozolomide for patients with high-risk disease is now a category 2A option.

### ASTR-3

#### Treatment

- Modified for prior fractionated EBRT, Consider reirradiation with *highly focused RT...especially if progression-free survival is greater than 2 y after prior RT or*

### GLIO-1

#### Footnotes

- "e": Biopsy *first prior to administration of steroids* if MRI compatible with CNS lymphoma.

### GLIO-2

#### Adjuvant Treatment

- ~~Fractionated external-beam RT~~ replaced with *Standard RT* throughout the page. (Also for [EPEN-2](#), [EPEN-3](#), [EPEN-4](#))

#### Follow-Up

- Brain MRI 2–8 wks (~~6 weeks~~). (Also for [GLIO-3](#) and [GLIO-4](#))
- corresponding to *Standard brain RT + concurrent and adjuvant lomustine and temozolomide (category 2B)*

[Continued](#)

UPDATES



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Updates in Version 1.2020 of the NCCN Guidelines for Central Nervous System Cancers from Version 3.2019 include:

### [GLIO-3](#)

#### Adjuvant Treatment

- **Standard brain RT + concurrent and adjuvant lomustine and temozolomide (category 2B)** for patients with good performance status and methylated tumors is new to the page.

#### Footnotes

- "r": Combination of agents modalities may lead to increased toxicity or radiographic changes. (Also for [GLIO-4](#)).
- "t": *Moderate to significant myelosuppression was observed, but the toxicity profile for this regimen is not yet fully defined*, is new to the page corresponding to **Standard brain RT + concurrent and adjuvant lomustine and temozolomide (category 2B)**

### [GLIO-4](#)

#### Footnotes

- "w": See *NCCN Guidelines for Older Adult Oncology*, corresponding to Age >70 y Glioblastoma.

### [EPEN-2](#)

#### Pathology

Ependymoma (Grades 1 and II). (Also for [EPEN-3](#)).

#### Footnotes

- "h": Lumbar puncture is indicated when there is clinical concern for meningeal dissemination. Lumbar puncture should be done after MRI of spine is performed to avoid a false-positive imaging result. Lumbar puncture for CSF should be delayed at least 2 weeks after surgery to avoid possible false-positive cytology. Lumbar puncture may be contraindicated (eg, posterior fossa mass). (Also for [EPEN-3](#), [EPEN-4](#)).
- "k": *Consider proton therapy if available to reduce toxicity*, is a new to the page and corresponds to Craniospinal RT. (Also for [EPEN-3](#), [EPEN-4](#), [AMED-2](#))

### [EPEN-3](#)

#### Postoperative Staging

- Grade I, top pathway: ~~Gross total~~ **En bloc** resection

### [EPEN-4](#)

- Resectable, evidence of metastasis, ~~Gross total or~~ removed

### [AMED-3](#)

#### Follow-Up

Brain MRI: every 3 mo for 2 y; then every 6–12 mo for 5–10 y; *then every 1–2 y or as clinically indicated*. For patients with previous spine disease, concurrent spine imaging as clinically indicated

### [PCNS-1](#)

#### Footnotes

- "f": Modified: Brain biopsy is recommended as the primary procedure to obtain diagnosis. CSF analysis should include flow cytometry, CSF cytology, *cell count, and* possibly gene rearrangements.

### [PCNS-2](#)

#### Evaluation

- *Lactate dehydrogenase (LDH) test*, is new to the page.

#### Induction Therapy

- Modified: High-dose methotrexate-based regimen *or other systemic therapy regimen if patient is unsuitable for or intolerant to high-dose methotrexate*
- Changed ~~vitreous~~ to **vitreoretinal** for eye exam.

### [PCNS-2A](#)

#### Footnotes

- "m": Dose adjusted for GFR *if dosing at 8 g/m<sup>2</sup>*.

### [PSCT-4](#)

#### Treatment for Recurrence

- Chemotherapy *relative to cell type if further surgery or RT not possible*

[Continued](#)  
**UPDATES**



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Updates in Version 1.2020 of the NCCN Guidelines for Central Nervous System Cancers from Version 3.2019 include:

### MENI-1

#### Footnotes

- "f" deleted: RT can be either external beam or SRS

### LTD-2

#### Treatment

- SRS in select patients for disseminated disease with poor systemic treatment options is new to the page

#### Footnotes

- "g": modified 2nd sentence to include: *Consultation with a radiation oncologist and close MRI surveillance is strongly recommended.*
- "l": Hippocampal avoidance preferred, [See BRAIN-C](#), corresponding to WBRT, (Also for [LTD-3](#)).

### LTD-3

#### Follow-Up

- Brain MRI modified: every 2–3 mo for 1–2 y then every 4–6 mo indefinitely (Also for [MU-2](#))

#### Treatment

- Surgery, modified: *followed by SRS or RT to the surgical bed*
- Chemotherapy was changed to *Systemic therapy* for all pathways on the page (Also for [MU-2](#))

#### Follow-Up

- "n": modified: After SRS, recurrence on MRI radiograph can be... (Also for [MU-2](#))

### LTD-4

- The following treatment options were added for patients with *stable systemic disease or reasonable systemic treatment options*:
  - ▶ Surgery
  - ▶ SRS
  - ▶ WBRT if no prior WBRT
  - ▶ Systemic therapy
- SRS in select patients was added as a treatment option for patients with systemic disease progression.

### MU-1

#### Primary Treatment

- WBRT, or SRS or Systemic therapy, with the following corresponding footnotes:
  - ▶ "g": *If an active agent exists (eg, cytotoxic, targeted, or immune modulating), trial of systemic therapy with good CNS penetration may be considered in select patients (eg, for patients with small asymptomatic brain metastases from melanoma or ALK rearrangement-positive NSCLC or EGFR-mutated NSCLC); it is reasonable to hold on treating with radiation to see if systemic therapy can control the brain metastases. Consultation with a radiation oncologist and close MRI surveillance is strongly recommended. There are no data from prospective clinical trials comparing the two strategies to assess what the impact of delayed radiation would be in terms of survival or in delay of neurologic deficit development.*
  - ▶ "h": [See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\)](#)

### LEPT-2

#### Treatment

- For Good Risk Status: SRS or RT (involved-field and/or whole brain) to bulky disease and neurologically symptomatic (such as cranial neuropathies) or painful sites. This replaced "WBRT and/or involved-field RT"

### BRAIN-A

- 1st bullet, 4th sub-bullet, Limitations modified: ~~patients with implantable devices are unsafe for cannot have an MRI...or renal insufficiency may occur~~

### BRAIN-C 1 of 8

#### Adult Low-Grade (WHO Grade I or II) Glioma/Pilocytic and Infiltrative Supratentorial Astrocytoma/Oligodendroglioma

- Tumor volumes are best defined using pre- and postoperative MRI imaging, usually T2 fluid-attenuated inversion recovery (FLAIR)/T2 and T1-, occasionally, enhanced T1 post-contrast sequences,...
- *New MRI for radiation treatment planning is recommended as there can be changes in mass effect and cytotoxic edema. Distinguishing non-enhancing tumor from vasogenic edema can be challenging and may warrant consultation with a neuroradiologist to inform treatment planning.*

[Continued](#)

UPDATES





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Updates in Version 1.2020 of the NCCN Guidelines for Central Nervous System Cancers from Version 3.2019 include:

### [BRAIN-C 1 of 8](#) (continued)

#### Anaplastic Gliomas/Glioblastoma High-Grade (Grades III/IV)

- Tumor volumes are best defined using pre- and postoperative MRI imaging using ~~enhanced post-contrast T1 with/without~~ and FLAIR/T2 sequences to define GTV. To account for sub-diagnostic tumor infiltration, the GTV is expanded 1–2 cm (CTV) for grade III ~~up to 2–2.5 cm (CTV) for grade...~~
- *Consider proton therapy for patients with good long-term prognosis (grade III IDH-mutant tumors [Buckner 2016] and grade III 1p19q co-deleted tumors [Shih 2015]) to better spare un-involved brain and preserve cognitive function, is new to the page.*

#### RT Dosing Information

- 4th bullet, second sentence modified: Typical fractionation schedules are 34 Gy/10 fx, 40.05 Gy/15 fx, ~~or 50 Gy/20 fx.~~

### [BRAIN-C 4 of 8](#)

- WHO Grade 1 Meningiomas  
2nd, sub,sub-bullet modified (25–30 Gy in 5 fractions)

### [BRAIN-C 6 of 8](#)

- 1st sub-bullet under RT dosing modified: ..spinal cord and/or nerve root
- 4th sub-bullet modified: When lower BED regimens are utilized upfront (ie, BED ≤60 Gy<sup>2</sup>, which includes up to 20 Gy in 5 fractions but does not include 30 Gy in 10 fractions), retreatment with similar BED regimens, such as 20 Gy in 5 fractions or 8 Gy in 1 fraction, can safely be considered as early as ~~4 weeks~~ to 6 weeks from initial treatment for pain relief.
- 5th sub-bullet modified: In other cases of retreatment, doses ranging from 15 Gy in 1 fraction *with SBRT* to 40 Gy in 20 fractions *with a conformal approach* have been utilized for tumor control, with careful consideration of tolerance of the spinal cord and/or nerve roots. In these instances, it is generally recommended that 6 months or more of time between treatments is required.

### [BRAIN-D \(1 of 15\)](#)

#### Adjuvant Treatment

Useful in Certain Circumstances; modified as follows:

- Pilocytic astrocytoma, PXA, ganglioglioma if *BRAF V600E* activating mutation
  - ▶ BRAF/MEK inhibitors:
    - ◇ dabrafenib/trametinib
    - ◇ vemurafenib/cobimetinib

### [BRAIN-D \(2 of 15\)](#)

#### Anaplastic Gliomas

- ~~Carboplatin category 2B~~ removed under Other Recommended Regimens for Recurrence Therapy (Also for [BRAIN-D 3 of 15](#))

#### Footnotes

"j": modified: Bevacizumab + chemotherapy can be considered if bevacizumab monotherapy fails *and it is desirable to continue the steroid sparing effects of bevacizumab* (Also for [BRAIN-D 3 of 15](#)).

### [BRAIN-D \(3 of 15\)](#)

#### Adjuvant Treatment for Glioblastoma

Preferred Regimens; modified as follows:

- RT with concurrent and adjuvant TMZ ± TTF

Useful in Certain Circumstances; the following is new to the page:

- *RT with concurrent and adjuvant lomustine and TMZ (for patients with MGMT promoter methylated tumors, KPS ≥60, and age ≤70 years) (category 2B), with the following corresponding footnote: Moderate to significant myelosuppression was observed, but the toxicity profile for this regimen is not yet fully defined.*

#### Recurrence Therapy for Glioblastoma

#### Preferred Regimens:

*Regorafenib* is new to the page

### [BRAIN-D \(5 of 15\)](#)

#### Induction Therapy for Primary CNS Lymphoma

Useful in Certain Circumstances

- *Patient is unsuitable for or intolerant to high-dose methotrexate*
  - ▶ See "Other Recommended Regimens" for Relapsed or Refractory Disease

#### Relapsed or Refractory Disease

Other Recommended Regimens; modified as follows:

- Retreat with high-dose methotrexate
  - ▶ with ~~or without~~ rituximab and ibrutinib
- Topotecan
- Dexamethasone, high-dose cytarabine, cisplatin

### [BRAIN-D \(7 of 15\)](#)

Other Recommended Regimens for Meningioma; modified as follows:

- Interferon alfa (category 2B)
- Somatostatin analogue, if octreotide scan positive

Useful in Certain Circumstances; modified as follows:

- Somatostatin analogue (category 2B) if octreotide scan positive



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Updates in Version 1.2020 of the NCCN Guidelines for Central Nervous System Cancers from Version 3.2019 include:

### BRAIN-D (8 of 15)

This page has been significantly revised. Newly diagnosed and recurrent disease have been combined.

#### Brain Metastases

- Breast Cancer/HER2 positive
  - Capecitabine + neratinib (category 2B)
  - Capecitabine (category 2B)
- Breast Cancer/HER2 non-specific
  - Cisplatin (category 2B)
  - Etoposide (category 2B)
  - High-dose methotrexate (category 2B)
- Melanoma, BRAF non-specific
  - Ipilimumab + nivolumab (preferred)
- Non-Small Cell Lung Cancer
  - EGFR-sensitizing mutation positive
    - ◊ Afatinib (category 2B)
    - ◊ Gefitinib (category 2B)
    - ◊ *Pulsatile* erlotinib
  - ALK rearrangement positive
    - ◊ Ceritinib (category 2B)
  - ALK rearrangement positive or ROS1 positive
    - ◊ Crizotinib (category 2B)
  - PD-L1 positive
    - ◊ Nivolumab (category 2B)
- Small Cell Lung Cancer
  - ◊ Topotecan (category 2B)

#### Footnotes

- "n": Consider glucarpidase (carboxypeptidase G2) for prolonged methotrexate clearance due to methotrexate-induced renal toxicity.
- "r": For patients with asymptomatic brain metastases, it is reasonable to treat with these systemic therapies in lieu of upfront WBRT and SRS.

### BRAIN-D (9 of 15)

#### Leptomeningeal and Spine Metastases

- Treatment
  - Intra-CSF chemotherapy
    - ◊ Interferon-alfa (category 2B)
  - Lymphoma/leukemias
    - ◊ Intra-CSF chemotherapy
      - Cytarabine (moved to the Treatment section for all leptomeningeal metastases)
      - Methotrexate (moved to the Treatment section for all leptomeningeal metastases)
      - Rituximab (lymphoma-only)
    - ◊ High-dose methotrexate (lymphoma-only)

### BRAIN-D (9 of 15) (continued)

- Breast Cancers
  - ◊ Intra-CSF chemotherapy
    - Trastuzumab (HER2 positive)
- The references have been updated for this section

### BRAIN-F 6 of 8

#### Principles of Brain Tumor Pathology Molecular Markers

- Medulloblastoma Molecular Subtyping
  - 3rd bullet, 2nd sentence modified: *Unlike in children, 50% of adult medulloblastomas with loss of 6q and positive nuclear catenin had no CTNNB1 mutations, pointing towards the possibility of alternative mechanisms of WNT pathway activation in adult medulloblastoma. Adult and pediatric medulloblastomas are genetically distinct and require different algorithms for molecular risk stratification. WNT-driven tumors will also usually contain monosomy 6. 6q loss is not confined to WNT in adults; it is also described in SHH and Group 4. Monosomy 6 is a specific marker for pediatric WNT, but not for adult WNT.*
  - 5th bullet, 2nd and 4th sentences modified: *behavior, whereas often in the setting of germline TP53 mutations, wildtype SHH-activated medulloblastomas have a variable course, and are uncommon in adults. Non-WNT/non-SHH medulloblastomas also show a variable course. WNT tumors and Group 4 tumors have worse prognosis in adults compared to children based on retrospective data. 6q loss and positive nuclear catenin have no clear prognostic role in adult medulloblastoma.*

#### References:

- The following references are new to this section:
  - Korshunov A, Remke M, Werft W, et al. Adult and pediatric medulloblastomas are genetically distinct and require different algorithms for molecular risk stratification. *J Clin Oncol* 2010;28:3054-3060.
  - Remke M, Hielscher T, Northcott PA, et al. Adult medulloblastoma comprises three major molecular variants. *J Clin Oncol* 2011;29(19):2717-2723.
  - Zhukova N, Ramaswamy V, Remke M, et al. Subgroup-specific prognostic implication of TP53 mutation in medulloblastoma. *J Clin Oncol*. 2013;31(23):2927-2935.
  - Kool M, Jones DT, Jager N, et al. Genome sequencing of SHH medulloblastoma predicts genotype-related response to smoothened inhibition. *Cancer Cell*. 2014;25(3):393-405.
  - Waszak SM, Northcott PA, Buchhalter I, et al. Spectrum and prevalence of genetic predisposition in medulloblastoma: a retrospective genetic study and prospective validation in a clinical trial cohort. *Lancet Oncol*. 2018;19(6):785-798.






National  
Comprehensive  
Cancer  
Network®

# NCCN Guidelines Version 3.2020

## Adult Low-Grade (WHO Grade I or II) Glioma/Pilocytic and Infiltrative Supratentorial Astrocytoma/Oligodendroglioma

[NCCN Guidelines Index](#)  
[Table of Contents](#)  
[Discussion](#)



<sup>a</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>b</sup>[See Principles of Brain Tumor Surgery \(BRAIN-B\).](#)

<sup>c</sup>Recommended molecular diagnostics include 1p19q chromosomal status, IDH1/2 mutation status, and *BRAF* V600E mutation. [See Principles of Brain Tumor Pathology \(BRAIN-F\).](#)

<sup>d</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>e</sup>Consider a multidisciplinary review in treatment planning, especially once pathology is available. [See Principles of Brain and Spine Tumor Management \(BRAIN-E\).](#)

<sup>f</sup>If radiographically the tumor appears to be a high-grade glioma, [see GLIO-1](#).

<sup>g</sup>The need to treat SEGAs or other findings in the appropriate tuberous sclerosis patient population should be determined by the patient's symptoms and/or change on serial radiologic studies. Referral to a neurofibromatosis or specialty center is recommended.

<sup>h</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




# NCCN Guidelines Version 3.2020

## Adult Low-Grade (WHO Grade I or II) Glioma/Pilocytic and Infiltrative Supratentorial Astrocytoma/Oligodendroglioma

PRESENTATION<sup>c,e,f</sup>

## ADJUVANT TREATMENT

## FOLLOW-UP



<sup>c</sup>Recommended molecular diagnostics include 1p19q chromosomal status, IDH1/2 mutation status, and BRAF V600E mutation. [See Principles of Brain Tumor Pathology \(BRAIN-F\)](#).

<sup>e</sup>Consider a multidisciplinary review in treatment planning, especially once pathology is available. [See Principles of Brain and Spine Tumor Management \(BRAIN-E\)](#).

<sup>f</sup>If radiographically the tumor appears to be a high-grade glioma, [see GLIO-1](#).

<sup>i</sup>Low-risk features: ≤40 y and gross total resection (GTR).

<sup>j</sup>High-risk features: >40 y or subtotal resection (STR) or open or stereotactic biopsy. Other high-risk factors that are sometimes taken into consideration are tumor size, neurologic deficits, and presence of sequencing verified IDH wild type.

<sup>k</sup>Regular follow-up is essential for patients receiving observation alone after resection.

<sup>l</sup>In the event that other risk factors are considered and treatment is warranted, treat as high risk. There may also be rare circumstances in which treating a patient with fractionated external beam RT alone (category 2B) or chemotherapy alone

(category 2B) may be considered. [See Principles of Brain and Spinal Cord Tumor Radiation \(BRAIN-C\)](#) or [Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\)](#).

<sup>m</sup>For low-grade gliomas, [See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\)](#).

<sup>n</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\)](#).

<sup>o</sup>The results of RTOG 9802 showed that there was a significant improvement in median overall survival in high-risk low-grade glioma patients treated with RT followed by PCV x 6 cycles compared with RT alone after a tissue diagnosis was made. However, this important study did not address whether all of these patients should be treated right away. Observation after diagnosis may be a reasonable option for a high-risk low-grade glioma patient who is neurologically asymptomatic or stable. Close monitoring with brain MRIs is important.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



National  
Comprehensive  
Cancer  
Network®


# NCCN Guidelines Version 3.2020

## Adult Low-Grade (WHO Grade I or II) Glioma/Pilocytic and Infiltrative Supratentorial Astrocytoma/Oligodendroglioma

[NCCN Guidelines Index](#)  
[Table of Contents](#)  
[Discussion](#)

### RECURRENCE<sup>P</sup>

### TREATMENT



<sup>a</sup>See [Principles of Brain and Spine Tumor Imaging \(BRAIN-A\)](#).

<sup>d</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>e</sup>Consider a multidisciplinary review in treatment planning, especially once pathology is available. See [Principles of Brain and Spine Tumor Management \(BRAIN-E\)](#).

<sup>f</sup>If radiographically the tumor appears to be a high-grade glioma, see [GLIO-1](#).

<sup>m</sup>For low-grade gliomas, See [Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\)](#).

<sup>n</sup>See [Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\)](#).

<sup>p</sup>If GTR is achieved in a patient with low-risk disease, consider further observation.

<sup>q</sup>Recurrence on neuroimaging can be confounded by treatment effects. To confirm

tumor recurrence and assess for possible transformation of tumor to higher grade, strongly consider tumor tissue sampling (biopsy at minimum) if there is a high index of suspicion of recurrence. Sixty percent or more of astrocytomas and 40%–50% of oligodendrogliomas will eventually undergo transformation to a higher grade. For treatment of patients with transformation to high-grade disease, see [GLIO-1](#).

<sup>r</sup>Brain MRI every 2–3 months while on treatment, then every 6 months indefinitely, to assess disease recurrence/progression during treatment with chemotherapy. (See [BRAIN-A](#)).

<sup>s</sup>RT alone is not encouraged, but may be appropriate for select cases (eg, poor performance status).

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020


## Anaplastic Gliomas<sup>a</sup>/Glioblastoma

### RADIOLOGIC PRESENTATION<sup>b</sup>

### CLINICAL IMPRESSION

### SURGERY<sup>c</sup>

### PATHOLOGY<sup>d</sup>



<sup>a</sup>This pathway includes the classification of mixed anaplastic oligoastrocytoma (AOA), anaplastic astrocytoma (AA), anaplastic oligodendroglioma (AO), and other rare anaplastic gliomas.

<sup>b</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>c</sup>[See Principles of Brain Tumor Surgery \(BRAIN-B\).](#)

<sup>d</sup>[See Principles of Brain Tumor Pathology \(BRAIN-F\).](#)

<sup>e</sup>Biopsy prior to administration of steroids if MRI compatible with CNS lymphoma.

<sup>f</sup>Consider a multidisciplinary review in treatment planning, especially once pathology is available. [See Principles of Brain and Spine Tumor Management \(BRAIN-E\).](#)

<sup>g</sup>If frozen section diagnosis supports high-grade glioma.

<sup>h</sup>Consider carmustine (BCNU) wafer implant during maximal safe resection (category 2B). Treatment with carmustine wafer may impact enrollment in adjuvant clinical trials.

<sup>i</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>j</sup>The 2016 WHO Classification of Tumors of the CNS has deleted oligoastrocytoma as a category, although “anaplastic oligoastrocytoma, NOS” may continue to be used for 1) patients with mixed histology and no available molecular data (ie, no tissue available for analysis) for determining whether to classify as oligodendroglioma versus astrocytoma; or 2) rare instances in which the tumor has regions with histologic features of oligoastrocytoma with 1p19q-codeletion, and distinct regions with histologic features of astrocytoma without 1p19q-codeletion.

<sup>k</sup>This pathway also includes gliosarcoma.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020


## Anaplastic Gliomas<sup>a</sup>/Glioblastoma

### ANAPLASTIC GLIOMAS (SEE [GLIO-3/GLIO-4](#) FOR GLIOBLASTOMA)

#### **PATHOLOGY<sup>d</sup>**

#### **ADJUVANT TREATMENT**

#### **FOLLOW-UP<sup>b</sup>**



<sup>a</sup>This pathway includes the classification of mixed AOA, AA, AO, and other rare anaplastic gliomas.

<sup>b</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>d</sup>[See Principles of Brain Tumor Pathology \(BRAIN-F\).](#)

<sup>j</sup>The 2016 WHO Classification of Tumors of the CNS has deleted oligoastrocytoma as a category, although “anaplastic oligoastrocytoma, NOS” may continue to be used for 1) patients with mixed histology and no available molecular data (ie, no tissue available for analysis) for determining whether to classify as oligodendroglioma versus astrocytoma; or 2) rare instances in which the tumor has regions with histologic features of oligoastrocytoma with 1p19q-codeletion, and distinct regions with

histologic features of astrocytoma without 1p19q-codeletion.

<sup>l</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>m</sup>The panel recommends that PCV be administered after RT (as per EORTC 26951) since the intensive PCV regimen given prior to RT (RTOG 9402) was not tolerated as well.

<sup>n</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>o</sup>Consider temozolomide if tumor is MGMT promoter methylated.

<sup>p</sup>Within the first 3 months after completion of RT and concomitant temozolomide, diagnosis of recurrence can be indistinguishable from pseudoprogression on neuroimaging.

**Note:** All recommendations are category 2A unless otherwise indicated.


**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





# NCCN Guidelines Version 3.2020

## Anaplastic Gliomas<sup>a</sup>/Glioblastoma



<sup>a</sup>This pathway includes the classification of mixed AOA, AA, AO, and other rare anaplastic gliomas.

<sup>b</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>d</sup>[See Principles of Brain Tumor Pathology \(BRAIN-F\).](#)

<sup>k</sup>This pathway also includes gliosarcoma.

<sup>l</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>n</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>o</sup>Consider temozolomide if tumor is MGMT promoter methylated.

<sup>p</sup>Within the first 3 months after completion of RT and concomitant temozolomide, diagnosis of recurrence can be indistinguishable from pseudoprogression on neuroimaging.

<sup>q</sup>MGMT= O6-methylguanine-DNA methyltransferase.

<sup>r</sup>Combination of modalities may lead to increased toxicity or radiographic changes.

<sup>s</sup>Benefit of treatment with temozolomide for glioblastomas beyond 6 months is unknown.

<sup>t</sup>Moderate to significant myelosuppression was observed, but the toxicity profile for this regimen is not yet fully defined.

<sup>u</sup>Clinical benefit from temozolomide is likely to be lower in patients whose tumors lack MGMT promoter methylation.

<sup>v</sup>Alternating electric field therapy is only an option for patients with supratentorial disease.


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Anaplastic Gliomas<sup>a</sup>/Glioblastoma



<sup>a</sup>This pathway includes the classification of mixed AOA, AA, AO, and other rare anaplastic gliomas.

<sup>b</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>d</sup>[See Principles of Brain Tumor Pathology \(BRAIN-F\).](#)

<sup>k</sup>This pathway also includes gliosarcoma.

<sup>l</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>n</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>o</sup>Consider temozolomide if tumor is MGMT promoter methylated.

<sup>p</sup>Within the first 3 months after completion of RT and concomitant temozolomide, diagnosis of recurrence can be indistinguishable from pseudoprogression on neuroimaging.

<sup>q</sup>MGMT= O6-methylguanine-DNA methyltransferase.

<sup>r</sup>Combination of modalities may lead to increased toxicity or radiographic changes.

<sup>s</sup>Benefit of treatment with temozolomide for glioblastomas beyond 6 months is unknown.

<sup>u</sup>Clinical benefit from temozolomide is likely to be lower in patients whose tumors lack MGMT promoter methylation.

<sup>v</sup>Alternating electric field therapy is only an option for patients with supratentorial disease.

<sup>w</sup>[See NCCN Guidelines for Older Adult Oncology.](#)


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Anaplastic Gliomas<sup>a</sup>/Glioblastoma



<sup>a</sup>This pathway includes the classification of mixed AOA, AA, AO, and other rare anaplastic gliomas.

<sup>b</sup>[See Principles of Brain and Spine Tumor Imaging \(BRIN-A\).](#)

<sup>i</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>j</sup>The 2016 WHO Classification of Tumors of the CNS has deleted oligoastrocytoma as a category, although “anaplastic oligoastrocytoma, NOS” may continue to be used for 1) patients with mixed histology and no available molecular data (ie, no tissue available for analysis) for determining whether to classify as oligodendroglioma versus astrocytoma; or 2) rare instances in which the tumor has regions with histologic features of oligoastrocytoma with 1p19q-codeletion, and distinct regions with histologic features of astrocytoma without 1p19q-codeletion.

<sup>l</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRIN-C\).](#)

<sup>n</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRIN-D\).](#)

<sup>p</sup>Within the first 3 months after completion of RT and concomitant temozolomide, diagnosis of recurrence can be indistinguishable from pseudoprogression on neuroimaging.

<sup>x</sup>Consider carmustine (BCNU) wafer implant during resection. Treatment with carmustine wafer may impact enrollment in clinical trials.

<sup>y</sup>The efficacy of standard-of-care treatment for recurrent glioblastoma is suboptimal, so for eligible patients consideration of clinical trials is highly encouraged. Prior treatment may impact enrollment in clinical trials.

<sup>z</sup>Consider biopsy, MR spectroscopy, MR perfusion, brain PET/CT, or brain PET/MRI, or re-image to follow changes that may be due to progression versus radionecrosis.

<sup>aa</sup>Anaplastic oligodendrogliomas have been reported to be especially sensitive to chemotherapy. Chemotherapy using temozolomide or nitrosourea-based regimens may be appropriate.

<sup>bb</sup>Especially if long interval since prior RT and/or if there was a good response to prior RT.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020


## Adult Intracranial and Spinal Ependymoma (Excluding Subependymoma)

### RADIOLOGIC PRESENTATION<sup>a</sup>

### CLINICAL IMPRESSION

### SURGERY<sup>d</sup>

### PATHOLOGY<sup>e</sup>


<sup>a</sup>See [Principles of Brain and Spine Tumor Imaging \(BRAIN-A\)](#).

<sup>b</sup>Consider a multidisciplinary review in treatment planning, especially once pathology is available. See [Principles of Brain and Spine Tumor Management \(BRAIN-E\)](#).

<sup>c</sup>If image-confirmed GTR not achieved, consider multidisciplinary review and resection.

<sup>d</sup>See [Principles of Brain Tumor Surgery \(BRAIN-B\)](#).

<sup>e</sup>Testing for RELA fusion may be recommended for ependymomas. See [Principles of Brain Tumor Pathology \(BRAIN-F\)](#).

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020


## Adult Intracranial and Spinal Ependymoma (Excluding Subependymoma)

### INTRACRANIAL EPENDYMOMA

#### PATHOLOGY

#### POSTOPERATIVE STAGING

#### ADJUVANT TREATMENT<sup>i</sup>



<sup>a</sup>See Principles of Brain and Spine Tumor Imaging (BRAIN-A).

<sup>f</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>g</sup>If not done preoperatively, spine MRI should be delayed by at least 2–3 weeks post surgery to avoid post-surgical artifacts.

<sup>h</sup>Lumbar puncture is indicated when there is clinical concern for meningeal dissemination. Lumbar puncture should be done after MRI of spine is performed to avoid a false-positive imaging result. Lumbar puncture for CSF should be

delayed at least 2 weeks after surgery to avoid possible false-positive cytology.

Lumbar puncture may be contraindicated (eg, posterior fossa mass).

<sup>i</sup>See Principles of Brain and Spinal Cord Tumor Radiation Therapy (BRAIN-C).

<sup>j</sup>Data supporting observation alone are based on retrospective studies.

<sup>k</sup>Consider proton therapy if available to reduce toxicity.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.






# NCCN Guidelines Version 3.2020

## Adult Intracranial and Spinal Ependymoma (Excluding Subependymoma)

### SPINAL EPENDYMOMA



<sup>a</sup>See [Principles of Brain and Spine Tumor Imaging \(BRAIN-A\)](#).

<sup>h</sup>Lumbar puncture is indicated when there is clinical concern for meningeal dissemination. Lumbar puncture should be done after MRI of spine is performed to avoid a false-positive imaging result. Lumbar puncture for CSF should be delayed at least 2 weeks after surgery to avoid possible false-positive cytology.

Lumbar puncture may be contraindicated (eg, posterior fossa mass).

<sup>i</sup>See [Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\)](#).

<sup>k</sup>Consider proton therapy if available to reduce toxicity.

<sup>l</sup>If not done preoperatively, spine MRI should be performed 48 h post surgery.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




# NCCN Guidelines Version 3.2020

## Adult Intracranial and Spinal Ependymoma (Excluding Subependymoma)

FOLLOW-UP<sup>a</sup>RECURRENCE STAGING WORKUP<sup>a</sup>TREATMENT FOR  
PROGRESSION OR  
RECURRENCE

- Imaging in the event of emergent signs or symptoms (brain and/or spine MRI)
- Imaging of tumor site (brain or spine MRI) every 3–4 mo for 1 y, then every 4–6 mo for year 2, then every 6–12 mo for 5–10 y, then as clinically indicated


Spine or  
brain  
recurrenceMRI imaging  
of spine,  
brain, and  
CSF analysis<sup>h</sup><sup>a</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)<sup>h</sup>Lumbar puncture is indicated when there is clinical concern for meningeal dissemination. Lumbar puncture should be done after MRI of spine is performed to avoid a false-positive imaging result. Lumbar puncture for CSF should be delayed at least 2 weeks after surgery to avoid possible false-positive cytology. Lumbar puncture may be contraindicated (eg, posterior fossa mass).<sup>i</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)<sup>k</sup>Consider proton therapy if available to reduce toxicity.<sup>m</sup>Consider stereotactic radiosurgery (SRS) if geometrically favorable.<sup>n</sup>Chemotherapy should be reserved for patients who are refractory to surgery or radiation.<sup>o</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### RADIOLOGIC PRESENTATION<sup>a</sup>

### CLINICAL IMPRESSION

### SURGERY<sup>d</sup>



<sup>a</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>b</sup>Consider a multidisciplinary review in treatment planning, before surgery and once pathology is available. [See Principles of Brain and Spine Tumor Management \(BRAIN-E\).](#)

<sup>c</sup>Placement of ventriculoperitoneal (VP) shunt for management of hydrocephalus is acceptable if needed.

<sup>d</sup>[See Principles of Brain Tumor Surgery \(BRAIN-B\).](#)


<sup>e</sup>Strongly recommend referring patient to a brain tumor center to be evaluated for possible further, more complete surgical resection.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### POSTOPERATIVE STAGING



<sup>a</sup>See [Principles of Brain and Spine Tumor Imaging \(BRAIN-A\)](#).

<sup>f</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>g</sup>Spine MRI should be delayed by at least 2–3 weeks post surgery to avoid post-surgical artifacts.

<sup>h</sup>Lumbar puncture should be done after spine MRI. Lumbar puncture for CSF should be delayed at least 2 weeks after surgery to avoid possible false-positive cytology.

<sup>i</sup>Bone scan; CT with contrast of chest, abdomen, and pelvis or whole body PET/CT; and bone marrow biopsy only if clinically indicated.

<sup>j</sup>Molecular profiling to identify clinically relevant subtypes is recommended to encourage opportunities for clinical trial involvement. See [Principles of Pathology \(BRAIN-F\)](#).

<sup>k</sup>See the modified Chang system for staging medulloblastoma. (Chang CH, Housepain EM, Herbert C. Radiology 1969;93:1351-1359 and Cohen ME, Duffner PK (Eds). Brain tumors in children, 2nd ed, McGraw-Hill, New York, 1994, p.187.)

<sup>l</sup>If only biopsy is possible, consider an attempt at resection at a center that specializes in the treatment of brain tumors.

<sup>m</sup>See [Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\)](#).

<sup>n</sup>Consider proton therapy if available to reduce toxicity.

<sup>o</sup>Omission of vincristine during radiotherapy phase of therapy or dose modification may be required for adults because they do not tolerate this regimen as well. Data supporting vincristine's use have been found in pediatric trials only. Patients should be closely monitored for neurologic toxicity with periodic exams. (Packer RJ, Gajjar A, Vezina G, et al. Phase III study of craniospinal radiation therapy followed by adjuvant chemotherapy for newly diagnosed average-risk medulloblastoma. J Clin Oncol 2006;24:4202-4208.)

<sup>p</sup>See [Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\)](#).

<sup>q</sup>Recommend a platinum-based chemotherapy regimen such as either of the treatment arms used in the Children's Oncology Group study referenced in footnote "k."

<sup>r</sup>Consider collecting stem cells before craniospinal radiation.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




### FOLLOW-UP<sup>a</sup>

### CLINICAL STAGING

### SURGERY

### TREATMENT FOR RECURRENCE

**Brain MRI:**  
every 3 mo for 2 y;  
then every 6–12 mo  
for 5–10 y; then every  
1–2 y or as clinically  
indicated  
For patients with  
previous spine disease,  
concurrent spine  
imaging as clinically  
indicated



<sup>a</sup>See [Principles of Brain and Spine Tumor Imaging \(BRAIN-A\)](#).

<sup>f</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>g</sup>Spine MRI should be delayed by at least 2–3 weeks post surgery to avoid post-surgical artifacts.

<sup>p</sup>See [Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\)](#).

<sup>s</sup>As clinically indicated, consider bone scan; contrast-enhanced CT scans of chest, abdomen, and pelvis; and/or bone marrow biopsy.

<sup>t</sup>Consider resection for palliation of symptoms where indicated.

<sup>u</sup>See [Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\)](#).

<sup>v</sup>Only if the patient is without evidence of disease after surgery or conventional dose re-induction chemotherapy.


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





### DIAGNOSIS BY TISSUE EVALUATION



<sup>a</sup>If patient is HIV positive, antiretroviral (ARV) therapy should be part of his/her treatment. ARVs can be administered safely with chemotherapy but consultation with an HIV specialist or pharmacist is important to optimize compatibility. [See NCCN Guidelines for Cancer in People Living with HIV.](#)

<sup>b</sup>For additional guidance on management of transplant recipients with PCNSL, [see NCCN Guidelines for B-Cell Lymphomas, Diffuse Large B-Cell Lymphoma.](#)

<sup>c</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>d</sup>Includes primary CNS lymphoma of the brain, spine, CSF, and leptomeninges. For lymphoma with primary tumor outside the CNS or involving only the eye, [See NCCN Guidelines for B-Cell Lymphomas, Diffuse Large B-Cell Lymphoma.](#)

<sup>e</sup>If stereotactic biopsy is not available refer to a specialized center.

<sup>f</sup>Brain biopsy is recommended as the primary procedure to obtain diagnosis. CSF analysis should include flow cytometry, CSF cytology, cell count, and possibly gene rearrangements.


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Primary CNS Lymphoma<sup>a,b</sup>



[See Follow-up \(PCNS-3\)](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### FOOTNOTES

<sup>a</sup>If patient is HIV positive, ARV therapy should be part of his/her treatment. ARVs can be administered safely with chemotherapy but consultation with an HIV specialist or pharmacist is important to optimize compatibility. [See NCCN Guidelines for Cancer in People Living with HIV.](#)

<sup>b</sup>For additional guidance on management of transplant recipients with PCNSL, [see NCCN Guidelines for B-Cell Lymphomas, Diffuse Large B-Cell Lymphoma.](#)

<sup>c</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>g</sup>May institute primary therapy and workup simultaneously.

<sup>h</sup>For full details regarding evaluation of extent of disease and response criteria for primary CNS lymphoma, refer to Abrey LE, Batchelor TT, Ferreri AJ, et al. Report of an international workshop to standardize baseline evaluation and response criteria for primary CNS lymphoma. J Clin Oncol 2005;23:5034-5043.

<sup>i</sup>CSF analysis should include flow cytometry, and CSF cytology, and may consider gene rearrangements.

<sup>j</sup>Caution is indicated in patients who are anticoagulated, thrombocytopenic, or who have a bulky intra-cranial mass.

<sup>k</sup>Recommend regular testicular exams. If PET/CT scan is negative, then there is no need for testicular ultrasound.

<sup>l</sup>A low KPS should not be a reason to withhold chemotherapy. KPS may improve dramatically after treatment.

<sup>m</sup>Dose adjusted for glomerular filtration rate (GFR) if dosing at 8 g/m<sup>2</sup>.

<sup>n</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>o</sup>If CSF positive or spinal MRI positive, consider alternative systemic chemotherapy regimens and/or intra-CSF chemotherapy (category 2B), especially for patients who cannot tolerate systemic methotrexate  $\geq 3$  g/m<sup>2</sup>.

<sup>p</sup>Due to a lack of strong evidence, it is not clear which consolidation regimen provides the most benefit.

<sup>q</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>r</sup>WBRT may increase neurotoxicity, especially in patients  $>60$  y.


<sup>s</sup>Includes primary CNS lymphoma of the brain, spine, CSF, and leptomeninges.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020 Primary CNS Lymphoma<sup>a,b</sup>


[See footnotes on PCNS-3A](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



National  
Comprehensive  
Cancer  
Network®

# NCCN Guidelines Version 3.2020

## Primary CNS Lymphoma<sup>a,b</sup>

[NCCN Guidelines Index](#)  
[Table of Contents](#)  
[Discussion](#)

### FOOTNOTES

<sup>a</sup>If patient is HIV positive, ARV therapy should be part of his/her treatment. ARVs can be administered safely with chemotherapy but consultation with an HIV specialist or pharmacist is important to optimize compatibility. [See NCCN Guidelines for Cancer in People Living with HIV.](#)

<sup>b</sup>For additional guidance on management of transplant recipients with PCNSL, [see NCCN Guidelines for B-Cell Lymphomas, Diffuse Large B-Cell Lymphoma.](#)

<sup>c</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>n</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>q</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>t</sup>The risk of neurotoxicity should be considered before administering high-dose therapy to a patient with prior WBRT.

<sup>u</sup>If the recurrent disease goes into complete remission with reinduction chemotherapy.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.






# NCCN Guidelines Version 3.2020

## Primary Spinal Cord Tumors

### RADIOLOGIC PRESENTATION<sup>a</sup>

### CLINICAL PRESENTATION

### SURGERY<sup>c</sup>



<sup>a</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>b</sup>Consider a multidisciplinary review in treatment planning, before surgery and once pathology is available. [See Principles of Brain and Spine Tumor Management \(BRAIN-E\).](#)

<sup>c</sup>[See Principles of Brain Tumor Surgery \(BRAIN-B\).](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Primary Spinal Cord Tumors

### RADIOLOGIC PRESENTATION

### PATHOLOGY

### CLINICAL PRESENTATION

### ADJUVANT TREATMENT

### FOLLOW-UP

Intradural  
Intramedullary  
tumor  
MRI well-defined/  
circumscribed

**Ependymoma**  
[See Adult Intracranial and Spinal  
Ependymoma \(EPEN-1\) and \(EPEN-3\)](#)

**Other subtypes:**  
 • Pilocytic  
astrocytoma  
 • Hemangioblastoma  
 ▶ Consider  
screening for  
von Hippel-  
Lindau  
syndrome<sup>d</sup>

Image-verified  
complete resection

Partial resection  
or biopsy

Asymptomatic

Symptomatic

Observation

Observation

RT<sup>e</sup>

**Follow-up**  
[See \(PSCT-4\)](#)

Intradural  
Intramedullary  
tumor  
MRI poorly  
defined/infiltrative

**Low-grade glioma**

• Anaplastic  
glioma  
 • Glioblastoma

[See Adult Low-Grade  
\(WHO Grade II\) Infiltrative  
Supratentorial Astrocytoma/  
Oligodendroglioma \(ASTR-1\)](#)

[See Anaplastic  
Gliomas/Glioblastoma  
\(GLIO-1\)](#)

<sup>d</sup>VHL Family Alliance: The VHL Handbook: What You Need to Know About VHL: A Reference Handbook for People with von Hippel-Lindau Disease, Their Families, and Support Personnel. Boston, MA, 2014 VHL Family Alliance.

<sup>e</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Primary Spinal Cord Tumors


### RADIOLOGIC PRESENTATION

### CLINICAL PRESENTATION

### SURGERY<sup>c</sup>

### PATHOLOGY

### FOLLOW-UP



<sup>a</sup>See Principles of Brain and Spine Tumor Imaging (BRAIN-A).

<sup>c</sup>See Principles of Brain Tumor Surgery (BRAIN-B).

<sup>f</sup>Spine MRI should be delayed by at least 2–3 weeks post surgery to avoid post-surgical artifacts.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




# NCCN Guidelines Version 3.2020

## Primary Spinal Cord Tumors

### FOLLOW-UP<sup>a</sup>

### RECURRENCE

### TREATMENT FOR RECURRENCE



<sup>a</sup>See Principles of Brain and Spine Tumor Imaging (BRAIN-A).

<sup>e</sup>See Principles of Brain and Spinal Cord Tumor Radiation Therapy (BRAIN-C).

<sup>g</sup>See Principles of Brain and Spinal Cord Tumor Systemic Therapy (BRAIN-D) for options according to disease histology.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Meningiomas

### PRESENTATION<sup>a</sup>

**Radiographic diagnosis by brain MRI:**

- Dural-based mass
- Homogeneously contrast-enhancing
- Dural tail
- CSF cleft

**Meningioma by radiographic criteria or Possible meningioma:**

- Consider resection
- Consider octreotide scan if diagnostic doubt exists

### TREATMENT<sup>b</sup>

Observe (preferred for small asymptomatic tumors; not generally recommended for symptomatic tumors)<sup>c</sup>

or

Surgery<sup>d,e</sup>  
(if accessible)<sup>f</sup>

or

RT<sup>f</sup>

### ADJUVANT TREATMENT

Consider RT<sup>f</sup> depending on factors in footnote "b"  
In general, postoperative management depends on grade,<sup>g</sup> extent of resection, and symptoms, as follows:

- Grade I: observation or consider RT (for symptomatic patients)
- Grade II with complete resection: consider RT
- Grade II with incomplete resection: RT
- Grade III: RT

**Follow-up**  
([See MENI-2](#))

<sup>a</sup>Multidisciplinary input for treatment planning if feasible.

<sup>b</sup>Treatment selection should be based on assessment of a variety of inter-related factors, including patient features (eg, age, performance score, comorbidities, treatment preferences), tumor features (eg, size, grade, growth rate, location [proximity to critical structures], potential for causing neurologic consequences if untreated, presence and severity of symptoms), and treatment-related factors (eg, potential for neurologic consequences from surgery/RT, likelihood of complete resection and/or complete irradiation with SRS, treatability of tumor if it progresses, available surgical or radiation oncology expertise and resources). The decision to administer RT after surgery also depends on the extent of resection

achieved. Multidisciplinary input for treatment planning is recommended.

<sup>c</sup>For asymptomatic meningiomas, observation is preferred for small tumors, with a suggested cutoff of ≤3 cm. Active treatment with surgery and/or RT is recommended in cases with one or more tumor- and/or treatment-related risk factors, such as proximity to the optic nerve.

<sup>d</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>e</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>f</sup>[See Principles of Brain Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>g</sup>WHO Grade I = Benign meningioma, WHO Grade II = Atypical meningioma, WHO Grade III = Malignant (anaplastic) meningioma.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### FOLLOW-UP<sup>h</sup>

### RECURRENCE/PROGRESSION

### TREATMENT

WHO Grade I and II<sup>e,i</sup>  
or unresected meningiomas:  
Brain MRI<sup>d,e</sup> at 3, 6, and 12  
mo,  
then every 6–12 mo for 5 y,  
then every 1–3 y as clinically  
indicated

Recurrent  
or  
progressive  
disease

Surgery if accessible

Brain MRI<sup>d,e</sup>

Consider clinical  
trial  
or  
RT<sup>f</sup> (if no prior RT)  
or  
Consider  
re-irradiation<sup>f</sup>

Not surgically accessible  
RT possible

RT<sup>f</sup>

Not surgically accessible  
RT not possible

Chemotherapy<sup>j</sup>

Treatment not clinically indicated

Observation

<sup>d</sup>Postoperative brain MRI within 48 hours after surgery.

<sup>e</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>f</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>h</sup>Consider less frequent follow-up after 5–10 y.

<sup>i</sup>More frequent imaging may be required for WHO Grade 3 meningiomas, and for meningiomas of any grade that are treated for recurrence or with chemotherapy.

<sup>j</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

**Note:** All recommendations are category 2A unless otherwise indicated.


**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





# NCCN Guidelines Version 3.2020

## Limited Brain Metastases



<sup>a</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>b</sup>Consider a multidisciplinary review in treatment planning, especially once pathology is available. [See Principles of Brain and Spine Tumor Management \(BRAIN-E\).](#)

<sup>c</sup>"Limited" brain metastases defines a group of patients for whom SRS is equally effective and offers significant cognitive protection compared with WBRT. The definition of "limited" brain metastases in terms of number of metastases or total intracranial disease volume is evolving and may depend on the specific clinical situation. (Yamamoto M, Serizawa T, Shuto T, et al. Stereotactic radiosurgery for patients with multiple brain metastases (JLGK0901): a multi-institutional prospective observational study. *Lancet Oncol* 2014;15:387-395.)

<sup>d</sup>[See Principles of Brain Tumor Surgery \(BRAIN-B\).](#)

<sup>e</sup>For secondary CNS lymphoma, treatment may include systemic treatment, whole-brain or focal RT, or combination.

<sup>f</sup>The decision to resect a tumor may depend on the need to establish histologic diagnosis, the size of the lesion, its location, and institutional expertise. For example, smaller (<2 cm), deep, asymptomatic lesions may be considered for treatment with SRS versus larger (>2 cm), symptomatic lesions that may be more appropriate for surgery. (Ewend MG, Morris DE, Carey LA, Ladha AM, Brem S. Guidelines for the initial management of metastatic brain tumors: role of surgery, radiosurgery, and radiation therapy. *J Natl Compr Cancer Netw* 2008;6:505-513.)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




# NCCN Guidelines Version 3.2020

## Limited Brain Metastases

### CLINICAL PRESENTATION

### TREATMENT<sup>g,h</sup>



<sup>e</sup>For secondary CNS lymphoma, treatment may include systemic treatment, whole-brain or focal RT, or combination.

<sup>g</sup>If an active agent exists (eg, cytotoxic, targeted, or immune modulating), trial of systemic therapy with good CNS penetration may be considered in select patients (eg, for patients with small asymptomatic brain metastases from melanoma or ALK rearrangement-positive NSCLC or EGFR-mutated NSCLC); it is reasonable to hold on treating with radiation to see if systemic therapy can control the brain metastases. Consultation with a radiation oncologist and close MRI surveillance is strongly recommended. There are no data from prospective clinical trials comparing the two strategies to assess what the impact of delayed radiation would be in terms of survival or in delay of neurologic deficit development.

<sup>h</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>i</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>j</sup>SRS is preferred when safe, especially for low tumor volume, to both the resection cavity and any other non-resected brain metastases. WBRT is generally not recommended but may be appropriate in some rare clinical circumstances.

<sup>k</sup>For brain metastases not managed with resection, SRS + WBRT is generally not recommended but may be appropriate in some rare clinical circumstances. Brown 2016 showed that for tumors <3 cm, SRS + WBRT improved local control compared with SRS alone, but did not significantly improve survival, and was associated with greater cognitive decline and poorer quality of life. (Brown PD, Jaeckle K, Ballman KV, et al. Effect of radiosurgery alone vs radiosurgery with whole brain radiation therapy on cognitive function in patients with 1 to 3 brain metastases: a randomized clinical trial. JAMA 2016;316:401-409.)

<sup>l</sup>Hippocampal avoidance preferred. [See BRAIN-C.](#)


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Limited Brain Metastases



<sup>a</sup>See [Principles of Brain and Spine Tumor Imaging \(BRAIN-A\)](#).

<sup>c</sup>"Limited" brain metastases defines a group of patients for whom SRS is equally effective and offers significant cognitive protection compared with WBRT. The definition of "limited" brain metastases in terms of number of metastases or total intracranial disease volume is evolving and may depend on the specific clinical situation. (Yamamoto M, Serizawa T, Shuto T, et al. Stereotactic radiosurgery for patients with multiple brain metastases (JLKG0901): a multi-institutional prospective observational study. *Lancet Oncol* 2014;15:387-395.)

<sup>h</sup>See [Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\)](#).

<sup>i</sup>See [Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\)](#).

<sup>l</sup>Hippocampal avoidance preferred. See [BRAIN-C](#).

<sup>m</sup>MRI every 2 months (instead of 3 mo) for those patients treated with SRS alone.

<sup>n</sup>After SRS, recurrence on MRI can be confounded by treatment effects; consider tumor tissue sampling if there is a high index of suspicion of recurrence.

<sup>p</sup>If patient had previous SRS with a good response >6 mo, then reconsider SRS if imaging supports active tumor and not necrosis.

<sup>q</sup>Imaging to evaluate emergent signs/symptoms is appropriate at any time.


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### RECURRENCE

### TREATMENT



<sup>d</sup>See Principles of Brain Tumor Surgery (BRAIN-B).

<sup>i</sup>See Principles of Brain and Spinal Cord Tumor Radiation Therapy (BRAIN-C).

<sup>h</sup>See Principles of Brain and Spinal Cord Tumor Systemic Therapy (BRAIN-D).


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Extensive Brain Metastases



<sup>a</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>b</sup>Consider a multidisciplinary review in treatment planning, especially once pathology is available. [See Principles of Brain and Spine Tumor Management \(BRAIN-E\).](#)

<sup>c</sup>Includes all cases that do not fit the definition of "limited brain metastases" on [LTD-1](#).

<sup>d</sup>[See Principles of Brain Tumor Surgery \(BRAIN-B\).](#)

<sup>e</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>f</sup>SRS can be considered for patients with good performance and low overall tumor volume and/or radioresistant tumors such as melanoma. (Yamamoto M, Serizawa T, Shuto T, et al. Stereotactic radiosurgery for patients with multiple brain metastases (JLGK0901): a multi-institutional prospective observational study. *Lancet Oncol* 2014;15:387-395.)

<sup>g</sup>If an active agent exists (eg, cytotoxic, targeted, or immune modulating), trial of systemic therapy with good CNS penetration may be considered in select patients (eg, for patients with small asymptomatic brain metastases from melanoma or ALK rearrangement-positive NSCLC or EGFR-mutated NSCLC); it is reasonable to hold on treating with radiation to see if systemic therapy can control the brain metastases. Consultation with a radiation oncologist and close MRI surveillance is strongly recommended. There are no data from prospective clinical trials comparing the two strategies to assess what the impact of delayed radiation would be in terms of survival or in delay of neurologic deficit development.

<sup>h</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




# NCCN Guidelines Version 3.2020

## Extensive Brain Metastases

### FOLLOW-UP<sup>a</sup>

### RECURRENCE

### TREATMENT



<sup>a</sup>See Principles of Brain and Spine Tumor Imaging (BRAIN-A).

<sup>e</sup>See Principles of Brain and Spinal Cord Tumor Radiation Therapy (BRAIN-C).

<sup>h</sup>See Principles of Brain and Spinal Cord Tumor Systemic Therapy (BRAIN-D).

<sup>i</sup>Imaging to evaluate emergent signs/symptoms is appropriate at any time.

<sup>j</sup>After SRS, recurrence on MRI can be confounded by treatment effects; consider tumor tissue sampling if there is a high index of suspicion of recurrence.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.






# NCCN Guidelines Version 3.2020

## Leptomeningeal Metastases

### WORKUP

### DIAGNOSIS

### RISK STATUS



<sup>a</sup>Consider a multidisciplinary review in treatment planning, especially once pathology is available. [See Principles of Brain and Spine Tumor Management \(BRAIN-E\)](#).

<sup>b</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\)](#).

<sup>c</sup>Caution is indicated in patients who are anticoagulated, thrombocytopenic, or who have a bulky intra-cranial mass.

<sup>d</sup>CSF analysis should include: a cell count, differential, glucose, and protein. For solid malignancies, order cytopathology. When available, assessment of circulating tumor cells (CTCs) increases sensitivity of tumor cell detection and assessment of response to treatment. For hematologic malignancies, use flow cytometry.

<sup>e</sup>Suggestive CSF includes high WBC count, low glucose, and high protein. If CSF is not positive for tumor cells, a second lumbar puncture is sometimes helpful. This is a volume-dependent test, and ideally ≥10 mL should be sent for cytologic analysis.

<sup>f</sup>Patients with exceptionally chemosensitive tumors (eg, small cell lung cancer, lymphoma) may be treated. Patients with a good risk status who do not desire further therapy may also be treated with palliative and/or best supportive care.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Leptomeningeal Metastases

### RISK STATUS

### TREATMENT

**Good risk:<sup>f</sup>**

- KPS ≥60
- No major neurologic deficits
- Minimal systemic disease
- Reasonable systemic treatment options, if needed

- Systemic chemotherapy<sup>g</sup>
- Intra-CSF chemotherapy<sup>g,h</sup>
  - ▶ If symptoms or imaging suggest CSF flow blockage, perform a CSF flow scan prior to starting intra-CSF chemotherapy. If flow abnormalities confirmed:
    - ◊ Fractionated external beam RT<sup>i</sup> to metastatic or painful sites of obstruction and repeat CSF flow scan to see if flow abnormalities have resolved.
    - or
    - ◊ High-dose methotrexate if breast cancer or lymphoma
- SRS or RT (involved-field and/or whole brain) to bulky disease and neurologically symptomatic (such as cranial neuropathies) or painful sites.<sup>i,j</sup>

[See Assessment of response \(LEPT-3\)](#)

**Poor risk:<sup>f</sup>**

- KPS <60
- Multiple, serious, major neurologic deficits
- Extensive systemic disease with few treatment options
- Bulky CNS disease
- Encephalopathy

Palliative/best supportive care and  
Consider involved-field RT<sup>i</sup> to neurologically symptomatic or painful sites for palliation (including spine and intracranial disease)

<sup>f</sup>Patients with exceptionally chemosensitive tumors (eg, small cell lung cancer, lymphoma) may be treated. Patients with a good risk status who do not desire further therapy may also be treated with palliative and/or best supportive care.

<sup>g</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>h</sup>Strongly consider Ommaya reservoir/intraventricular catheter.

<sup>i</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>j</sup>Due to substantial toxicity, craniospinal RT should only be considered in highly select patients (eg, leukemia, lymphoma).

**Note:** All recommendations are category 2A unless otherwise indicated.


**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Leptomeningeal Metastases

### TREATMENT



<sup>g</sup>See Principles of Brain and Spinal Cord Tumor Systemic Therapy (BRAIN-D).

<sup>i</sup>See Principles of Brain and Spinal Cord Tumor Radiation Therapy (BRAIN-C).

<sup>k</sup>If CSF cytology was initially negative, then assess response with MRI of spine/brain.

<sup>l</sup>If cytologic analysis is negative from CSF obtained from an Ommaya reservoir, then assess CSF obtained via a lumbar puncture to confirm CSF cytology is negative.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Metastatic Spine Tumors

### PRESENTATION

Patient diagnosed with cancer  
or patient with newly discovered abnormality  
suspicious for spine  
metastasis

Asymptomatic  
(Incidental finding)

### WORKUP

- Systemic imaging (ie, contrast-enhanced chest/abdominal/pelvic CT or whole body PET/CT, bone scan as indicated for metastatic workup)
- Biopsy<sup>a</sup> if it alters management

### TREATMENT

- Observation  
Spine MRI<sup>d</sup> in 6–8 weeks, then every 2–3 months until the nature of the lesion is established
- Surgery/focal RT<sup>e</sup> or chemotherapy<sup>f</sup> are options for patients with asymptomatic epidural disease

Symptomatic:  
• Severe, new, or  
progressive pain  
or neurologic  
symptoms or  
myelopathy

Spinal MRI<sup>b,c,d</sup>  
(urgent in the event of  
neurologic symptoms)

No tumor

Spinal cord compression<sup>g</sup>

No spinal cord compression<sup>d</sup>

[See  
SPINE-2](#)

<sup>a</sup>Biopsy if remote history of cancer.

<sup>b</sup>If the patient is unable to have an MRI, then a CT myelogram is recommended, which may also be useful for SRT planning.

<sup>c</sup>15%–20% of patients have additional lesions. Highly recommend complete spine imaging.

<sup>d</sup>[See Principles of Brain and Spine Tumor Imaging \(BRAIN-A\).](#)

<sup>e</sup>[See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>f</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>g</sup>Includes cauda equina syndrome.


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Metastatic Spine Tumors



<sup>f</sup>[See Principles of Brain and Spinal Cord Tumor Systemic Therapy \(BRAIN-D\).](#)

<sup>g</sup>Includes cauda equina syndrome.

<sup>h</sup>The recommended minimum dose of steroids is 4 mg of dexamethasone every 6 hours, although dose of steroids may vary (10–100 mg). A randomized trial supported the use of high-dose steroids (Sorensen PS, et al. Eur J Cancer 1994;30A:22-27).

<sup>i</sup>Spinal instability is grossly defined as the presence of significant kyphosis or subluxation (deformity), or of significantly retropulsed bone fragment.

<sup>j</sup>Consider alternative diagnosis of leptomeningeal disease ([See LEPT-1](#)).

<sup>k</sup>Tumor resection with or without spinal stabilization. Surgery should be focused on anatomic pathology.

Regarding surgery, note the following:

- Category 1 evidence supports the role of surgery in patients with a solitary epidural spinal cord compression by a tumor not known to be radiosensitive and who are willing to undergo surgery. (Patchell RA, et al. Lancet 2005;366(9486):643-648)
- For surgery, patients with hematologic tumors (ie, lymphoma, myeloma, leukemia) should be excluded, life expectancy should be ≥3 mo, and the patient should not be paraplegic for >24 h.
- Surgery is especially indicated if the patient has any of the following: spinal instability, no history of cancer, rapid neurologic deterioration during RT, previous RT to site, and single-site spinal cord compression.

<sup>m</sup>Postoperative spine MRI should be delayed by at least 2–3 weeks to avoid post-surgical artifacts. [See Principles of Brain Tumor Surgery \(BRAIN-B\).](#)

<sup>n</sup>Recommend SRS if oligometastases and radioresistant. [See Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\).](#)

<sup>o</sup>Vertebral augmentation: vertebroplasty, kyphoplasty.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




# NCCN Guidelines Version 3.2020

## Metastatic Spine Tumors

### FOLLOW-UP

### PRESENTATION (Symptom- or MRI-based)

### TREATMENT FOR RECURRENCE OR PROGRESSIVE DISEASE



<sup>d</sup>See [Principles of Brain and Spine Tumor Imaging \(BRAIN-A\)](#).

<sup>m</sup>Postoperative spine MRI should be delayed by at least 2–3 weeks to avoid post-surgical artifacts. See [Principles of Brain Tumor Surgery \(BRAIN-B\)](#).

<sup>n</sup>Recommend SRS if oligometastases and radioresistant. See [Principles of Brain and Spinal Cord Tumor Radiation Therapy \(BRAIN-C\)](#).

<sup>p</sup>Gary AK, et al. Prospective evaluation of spinal reirradiation by using stereotactic body radiation therapy: The University of Texas MD Anderson Cancer Center experience. Cancer 2011;117:3509-3516.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





### PRINCIPLES OF BRAIN AND SPINE TUMOR IMAGING<sup>1</sup>

This is a list of imaging modalities available and used in neuro-oncology primarily to make treatment decisions. The most common use for MR spectroscopy, MR perfusion, and PET scanning is to differentiate radiation necrosis from active tumor, as this might obviate the need for surgery or the discontinuation of an effective therapy. Imaging is always recommended to investigate emergent signs or symptoms.

- **MRI<sup>2</sup> of the brain and spine (with and without contrast):**
  - Gold standard
  - Provides a “static” picture of tumors
  - Benefits: Provides a reasonably good delineation of tumors; higher grade tumors and brain leptomeningeal metastasis usually enhance; lower grade tumors usually do not enhance
  - Limitations: Sensitive to movement, metallic objects cause artifact, implantable devices are unsafe for MRI, claustrophobia may be an issue, or renal insufficiency may occur
  - Postoperative brain MRI should be performed within 48 hours for gliomas and other brain tumors to determine extent of resection.
  - Postoperative spine MRI should be delayed by at least 2–3 weeks to avoid post-surgical artifacts.
  - Follow-up brain MRI should be performed at the frequency and intervals stated in the treatment algorithms. More frequent imaging may be done as clinically indicated by the treating physician, such as in the event of a clinical change such as development of seizures or neurologic deterioration.
- **CT of the brain and spine (with and without contrast):**
  - Should be used in patients who cannot have an MRI
  - Benefits: Claustrophobia or implantable devices are not an issue, can be done faster than an MRI
  - Limitations: Lacks resolution of MRI, especially in posterior fossa, or renal insufficiency
- **MR spectroscopy: Assess metabolites within tumors and normal tissue**
  - May be useful in differentiating tumor from radiation necrosis; may be helpful in grading tumors or assessing response
  - Area most abnormal would be the best place to target for a biopsy
  - Limitations: Tumors near vessels, air spaces, or bone. Extra time in MRI and others as noted under MRI
- **MR perfusion: Measures cerebral blood volume in tumors**
  - May be useful in differentiating grade of tumor or tumor versus radiation necrosis. Area of highest perfusion would be the best place to biopsy.
  - Limitations: Tumors near vessels, air spaces, bone, small-volume lesions, or tumors in the spinal cord. Extra time in MRI and others as noted under MRI.
- **Brain PET scanning: Assess metabolism within tumor and normal tissue by using radiolabeled tracers**
  - May be useful in differentiating tumor from radiation necrosis but has some limitations; may also correlate with tumor grade or provide the optimal area for biopsy
  - Limitations: Accuracy of interpretations, availability of equipment and isotopes

<sup>1</sup>The imaging modalities listed may not be available at every institution.

<sup>2</sup>Wen PY, Macdonald DR, Reardon DA, et al. Updated response assessment for high-grade gliomas: Response assessment in neuro-oncology working group. J Clin Oncol 2010;28:1963-1972.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF BRAIN TUMOR SURGERY

#### Guiding Principles

- Gross total resection (GTR) when appropriate
- Minimal surgical morbidity
- Accurate diagnosis

#### Factors

- Age
- Performance status (PS)
- Feasibility of decreasing the mass effect with surgery
- Resectability, including number of lesions, location of lesions, time since last surgery (recurrent patients)
- New versus recurrent tumor
- Suspected pathology – benign vs. malignant, possibility of other non-cancer diagnoses, projected natural history
- For patients with IDH1 mutations, there is evidence to suggest that a supramarginal resection is most appropriate, which would include not only enhancing areas but also T2/flair areas when appropriate in terms of a safe surgical approach, with the use of any and all surgical adjuncts possible.<sup>a</sup>

#### Options<sup>a</sup>

- GTR where feasible
- Stereotactic biopsy
- Open biopsy/debulking followed by planned observation or adjuvant therapy
- Chemotherapy implants, when indicated (See footnote h on [GLIO-1](#))
- Carmustine polymer wafer may be placed in the tumor resection cavity of patients.<sup>b,c</sup>

#### Tissue

- Sufficient tissue to pathologist for neuropathology evaluation and molecular correlates
- Frozen section analysis when possible to help with intraoperative decision-making
- Review by experienced neuropathologist
- Postoperative brain MRI should be performed within 48 hours for gliomas and other brain tumors to determine the extent of resection. Postoperative spine MRI should be delayed by at least 2–3 weeks to avoid post-surgical artifacts.
- The extent of resection should be judged on the postoperative study and used as a baseline to assess further therapeutic efficacy or tumor progression.

<sup>a</sup>A number of surgical adjuncts can be considered to facilitate safe brain tumor surgery, including use of an intraoperative microscope, frameless stereotactic image guidance, preoperative functional MRI and/or DTI fiber tracking, awake craniotomy, motor and/or speech mapping, intraoperative MRI, and intraoperative fluorescence-guided surgery with 5-ALA.

<sup>b</sup>Ewend MG, Brem S, Gilbert M, et al. Treatment of single brain metastasis with resection, intracavity carmustine polymer wafers, and radiation therapy is safe and provides excellent local control. Clin Cancer Res 2007;13:3637-3641.

<sup>c</sup>Brandes AA, Tosoni A, Amista P, et al. How effective is BCNU in recurrent glioblastoma in the modern era? A phase II trial. Neurology 2004;63:1281-1284.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF RADIATION THERAPY FOR BRAIN AND SPINAL CORD****Adult Low-Grade (WHO Grade I or II) Glioma/Pilocytic and Infiltrative Supratentorial Astrocytoma/Oligodendroglioma**

- Tumor volumes are best defined using pre- and postoperative MRI imaging, usually T2 fluid-attenuated inversion recovery (FLAIR) and T1 post-contrast sequences, to define gross tumor volume (GTV). Clinical target volume (CTV) (GTV plus 1–2 cm margin) should receive 45–54 Gy in 1.8–2.0 Gy fractions.<sup>1-3</sup> Consider RT dose escalation to 59.4–60 Gy for IDH wild-type low-grade gliomas, as these patients have a more aggressive course of disease.
- New MRI for radiation treatment planning is recommended as there can be changes in mass effect and cytotoxic edema. Distinguishing non-enhancing tumor from vasogenic edema can be challenging and may warrant consultation with a neuroradiologist to inform treatment planning.

**Anaplastic Gliomas/Glioblastoma High-Grade (Grades III/IV)*****Simulation and Treatment Planning***

- Tumor volumes are best defined using pre- and postoperative MRI imaging using post-contrast T1 and FLAIR/T2 sequences to define GTV. To account for sub-diagnostic tumor infiltration, the GTV is expanded 1–2 cm (CTV) for grade III and IV tumors. Although trials in glioblastoma have historically used CTV expansion in the range of 2 cm, smaller CTV expansions are supported in the literature and can be appropriate. A planning target volume (PTV) of margin of 3–5 mm is typically added to the CTV to account for daily setup errors and image registration. Daily image guidance is required if smaller PTV margins are used (3 mm or less). When edema as assessed by T2/FLAIR is included in the initial phase of treatment, fields are usually reduced for the last phase of the treatment (boost). The boost target volume will typically encompass only the gross residual tumor and the resection cavity. A range of acceptable clinical target volume margins exists. Both strategies appear to produce similar outcomes.<sup>4</sup>
- Consider proton therapy for patients with good long-term prognosis (grade III IDH-mutant tumors<sup>5</sup> and grade III 1p19q co-deleted tumors<sup>6</sup>) to better spare un-involved brain and preserve cognitive function.

***RT Dosing Information***

- The recommended dose is 60 Gy in 2.0 Gy fractions or 59.4 Gy in 1.8 Gy fractions.
- A slightly lower dose, such as 54–55.8 Gy in 1.8 Gy or 57 Gy in 1.9 Gy fractions, can be applied when the tumor volume is very large (gliomatosis), there is brainstem/spinal cord involvement, or for grade III astrocytoma.
- If a boost volume is used, the initial phase of the RT plan will receive 46 Gy in 2 Gy fractions or 45–50.4 Gy in 1.8 Gy fractions. The boost plan will typically then receive 14 Gy in 2 Gy fractions or 9–14.4 Gy in 1.8 Gy fractions.<sup>4</sup>
- In poorly performing patients or elderly patients, a hypofractionated accelerated course should be considered with the goal of completing the treatment in 2–4 weeks. Typical fractionation schedules are 34 Gy/10 fx or 40.05 Gy/15 fx.<sup>7,8</sup> Alternatively, a shorter fractionation schedule of 25 Gy/5 fx may be considered for elderly and/or frail patients with smaller tumors for whom a longer course of treatment would not be tolerable.<sup>9</sup>

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF RADIATION THERAPY FOR BRAIN AND SPINAL CORD

#### Adult Intracranial and Spinal Ependymoma

- **Limited Fields:**
  - ▶ Intracranial tumor volumes are best defined using pre- and postoperative MRI imaging, usually enhanced T1 and/or FLAIR/T2. GTV is defined as anatomic areas that are touched by preoperative tumor volume plus postoperative signal abnormality as seen on MRI.
  - ▶ RT Dosing Information:
    - ◊ CTV (GTV plus 1–2 cm margin) should receive 54–59.4 Gy in 1.8–2.0 Gy fractions. PTV of margin of 3–5 mm is typically added to the CTV to account for daily setup errors and image registration.
- **Craniospinal:**
  - ▶ To reduce toxicity from craniospinal irradiation (CSI) in adults, consider the use of intensity-modulated radiotherapy or protons if available.
  - ▶ RT Dosing Information:
    - ◊ Whole brain and spine (to bottom of thecal sac) receive 36 Gy in 1.8 Gy fractions, followed by limited field to spine lesions to 45 Gy. (Gross metastatic lesions below the conus could receive higher doses of 54–60 Gy.)<sup>10,11</sup>
    - ◊ Primary intracranial site should receive total dose of 54–59.4 Gy in 1.8–2.0 Gy fractions.
    - ◊ Consider boosting any gross intracranial metastatic sites to a higher dose while respecting normal tissue tolerances.
- **Spine Ependymoma:**
  - ▶ For spine ependymomas, see section on primary spinal cord tumors ([BRAIN-C 3 of 8](#)).<sup>12,13</sup>
  - ▶ CTV margins of 1–2 cm in the superior and inferior directions are recommended.
  - ▶ PTV of margin of 3–5 mm is typically added to the CTV to account for daily setup errors and image registration.

#### Adult Medulloblastoma

- **Standard Risk for Recurrence:**
  - ▶ Conventional dose: 30–36 Gy CSI<sup>14,†</sup> and boosting the primary brain site to 54–55.8 Gy with or without adjuvant chemotherapy.
  - ▶ Reduced dose: May consider reduced dose radiation with adjuvant chemotherapy: 23.4 Gy CSI<sup>14,15,††</sup> and boosting the primary brain site to 54–55.8 Gy.<sup>1</sup>
- **High Risk for Recurrence:**
  - ▶ 36 Gy CSI<sup>15,†</sup> with boosting primary brain site to 54–55.8 Gy with adjuvant chemotherapy.

†To reduce toxicity from craniospinal irradiation in adults, consider the use of intensity-modulated radiotherapy or protons if available.

††Regimen supported by data from pediatric trials only.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF RADIATION THERAPY FOR BRAIN AND SPINAL CORD

#### Primary CNS Lymphoma

- **WBRT may be withheld in the primary setting in patients treated with chemotherapy.**
  - ▶ **RT Dosing:**
    - ◊ **When used, low-dose WBRT should be limited to 23.4 Gy in 1.8 Gy fractions following a complete response (CR) to chemotherapy.<sup>16</sup>**
    - ◊ **For less than CR, consider WBRT to 30–36 Gy followed by a limited field to gross disease to 45 Gy or focal radiation to residual disease only.<sup>17-20</sup>**
  - ▶ **For patients who are not candidates for chemotherapy:**
    - ◊ **WBRT doses of 24–36 Gy followed by a boost to gross disease for a total dose of 45 Gy.**

#### Primary Spinal Cord Tumors

- ▶ **RT Dosing:**
  - ◊ **Doses of 45–54 Gy are recommended using fractions of 1.8 Gy.**
  - ◊ **In tumors below the conus medullaris higher doses up to 60 Gy may be delivered.**
  - ◊ **CTV margins of 1–2 cm in the superior and inferior directions are recommended.**
  - ◊ **PTV margins of 3–5 mm are typically added to the CTV to account for daily setup errors and image registration.**

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF RADIATION THERAPY FOR BRAIN AND SPINAL CORD

#### Meningiomas

- General Treatment Information

- ▶ If appropriate, may be treated using SRS or fractionated SRS.
- ▶ Highly conformal fractionated RT techniques (eg, 3D-CRT, IMRT, VMAT, proton therapy) are recommended to spare critical structures and uninvolved tissue.
- ▶ Stereotactic or image-guided therapy is recommended when using tight margins or when close to critical structures.

- **WHO Grade I Meningiomas:**

- ▶ **RT Dosing:**
  - ◊ 45–54 Gy.
  - ◊ WHO grade I meningiomas may also be treated with SRS doses of 12–16 Gy in a single fraction when appropriate, or consider hypofractionated SRT (25–30 Gy in 5 fractions) if near critical structures. Optimal dosing has not been determined.

- **WHO Grade II Meningiomas:**

- ▶ General Treatment Information

- ◊ Treatment should be directed to gross tumor (if present), surgical bed, and a margin (1–2 cm) to account for microscopic disease.
- ◊ Limit margin expansion into the brain parenchyma if there is no evidence of brain invasion.

- ▶ **RT Dosing:**

- ◊ 54–60 Gy in 1.8–2.0 Gy fractions.

- **WHO Grade III Meningiomas:**

- ▶ General Treatment Information

- ◊ Treat as malignant tumors with treatment directed to gross tumor (if present), surgical bed, and a margin (2–3 cm).

- ▶ **RT Dosing:**

- ◊ 59.4–60 Gy in 1.8–2.0 Gy fractions.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





### PRINCIPLES OF RADIATION THERAPY FOR BRAIN AND SPINAL CORD

#### Brain Metastases

- **WBRT: Doses vary between 20 and 40 Gy delivered in 5–20 fractions.**
  - ▶ The standard regimens include 30 Gy in 10 fractions or 37.5 Gy in 15 fractions.
  - ▶ Nevertheless, 20 Gy in 5 fractions is a good option for patients with poor predicted prognosis.<sup>21</sup>
  - ▶ For patients with a better prognosis, consider memantine during and after WBRT for a total of 6 months.<sup>22</sup>
  - ▶ For patients with a better prognosis (4 months or greater), consider hippocampal-sparing WBRT.<sup>23-24</sup>
- ▶ **SRS: Maximum marginal doses from 15–24 Gy based on tumor volume is recommended.<sup>25-28</sup>**
  - ◊ Consider fractionated SRS for brain tumor >3 cm.
    - Most common fractionated SRT doses include: 27 Gy in 3 fx and 30 Gy in 5 fx.
  - ◊ Postoperative SRS or SRT: Local recurrence rates after brain metastasis resections remain high (in the range of 50% at 1–2 years) even in the setting of a radiographic GTR. Postoperative SRS or SRT to the surgical cavity is supported by randomized data to improve local control over observation and to offer similar overall survival and superior cognitive preservation to postoperative WBRT.<sup>29,30</sup> A consensus statement regarding radiation target delineation has been published.<sup>31</sup> Fractionated SRT may be preferred for larger cavities.<sup>32</sup> Common dose-fractionation schedules include 16–20 Gy in 1 fraction, 27 Gy in 3 fractions, and 30 Gy in 5 fractions.
- ◊ SRS is generally preferred over WBRT for limited brain metastases.

#### Leptomeningeal Metastases

- ▶ Volume and dose depend on primary source and sites requiring palliation.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF RADIATION THERAPY FOR BRAIN AND SPINAL CORD****Metastatic Spine Tumors****• General Treatment Information**

- ▶ Doses to vertebral body metastases will depend on patient's PS, spine stability, location in relationship to spinal cord, primary histology, presence of epidural disease, and overall treatment intent (pain relief, long-term local control, or cure).
- ▶ Stereotactic radiation approaches (SRS/stereotactic body radiotherapy [SBRT]) for spinal cases may be preferred for patients with oligometastatic disease where tumor ablation is a goal of treatment and in tumors considered radioresistant (eg, renal cell, melanoma, sarcoma, hepatocellular, and some colorectal and NSCLC cases).
- ▶ Stereotactic radiation approaches may also be preferred in the setting of tumor recurrence after prior radiation as a strategy to limit radiation dose to the spinal cord or other critical structures. Careful adherence to consensus guidelines for radiosurgery planning and delivery is recommended.<sup>33-35</sup>

**• RT Dosing:**

- ▶ Generally, conventional external beam radiation doses of 8 Gy/1 fx, 20 Gy/5 fx, or 30 Gy/10 fx can be used. It is critical to consider tolerance at the spinal cord and/or nerve root. In selected cases, or recurrences after previous radiation, SBRT is appropriate.
- ▶ Common recommended doses for spine SRS/SBRT may include:
  - ◊ 16–24 Gy x 1 fx;
  - ◊ 24 Gy in 2 fx;
  - ◊ 24–27 Gy in 3 fx;
  - ◊ 30–35 Gy in 5 fx
- ▶ In patients with uncomplicated spine metastases that are treated primarily for pain relief, 8 Gy in 1 fraction has been shown to provide equivalent pain control to longer fractionation schedules. Single fraction treatment is more convenient for patients and an important consideration for patients with poor prognoses. This treatment may be associated with higher rates of retreatment, and a consideration for patients with a prognosis that exceeds 6 months or greater.
- ▶ When lower BED regimens are utilized upfront (ie, BED ≤60 Gy, which includes up to 20 Gy in 5 fractions but does not include 30 Gy in 10 fractions), retreatment with similar BED regimens, such as 20 Gy in 5 fractions or 8 Gy in 1 fraction, can safely be considered as early as 6 weeks from initial treatment for pain relief.
- ▶ In other cases of retreatment, doses ranging from 15 Gy in 1 fraction with SBRT to 40 Gy in 20 fractions with a conformal approach have been utilized for tumor control, with careful consideration of tolerance of the spinal cord and/or nerve roots. In these instances, it is generally recommended that 6 months or more of time between treatments is required.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR RADIATION THERAPY****REFERENCES**

- <sup>1</sup>Karim AB, Maat B, Hatlevoll R, et al. A randomized trial on dose-response in radiation therapy of low-grade cerebral glioma: European Organization for Research and Treatment of Cancer (EORTC) Study 22844. *Int J Radiat Oncol Biol Phys* 1996;36:549-556.
- <sup>2</sup>Shaw E, Arusell R, Scheithauer B, et al. Prospective randomized trial of low- versus high-dose radiation therapy in adults with supratentorial low-grade glioma: initial report of a North Central Cancer Treatment Group/Radiation Therapy Oncology Group/Eastern Cooperative Oncology Group study. *J Clin Oncol* 2002;20:2267-2276.
- <sup>3</sup>van den Bent MJ, Afra D, de Witte O, et al. Long-term efficacy of early versus delayed radiotherapy for low-grade astrocytoma and oligodendroglioma in adults: the EORTC 22845 randomised trial. *Lancet* 2005;366(9490):985-990.
- <sup>4</sup>Cabrera AR, Kirkpatrick JP, Fiveash JB, et al. Radiation therapy for glioblastoma: Executive summary of an American Society for Radiation Oncology Evidence-Based Clinical Practice Guideline. *Pract Radiat Oncol* 2016;6:217-225.
- <sup>5</sup>Buckner JC, Shaw EG, Pugh SL, et al. Radiation plus procarbazine, CCNU, and vincristine in low-grade glioma. *N Engl J Med* 2016;374:1344-1355.
- <sup>6</sup>Shih HA, Sherman JC, Nachtigall LB, et al. Proton therapy for low-grade gliomas: results from a prospective trial. *Cancer* 2015;121:1712-1719.
- <sup>7</sup>Malmström A, Grønberg BH, Marosi C, et al. Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised, phase 3 trial. *Lancet Oncol* 2012;13:916-920.
- <sup>8</sup>Roa W, Brasher PM, Bauman G, et al. Abbreviated course of radiation therapy in older patients with glioblastoma multiforme: a prospective randomized clinical trial. *J Clin Oncol* 2004;22:1583-1588.
- <sup>9</sup>Roa W, Kepka L, Kumar N, et al. International Atomic Energy Agency randomized phase III study of radiation therapy in elderly and/or frail patients with newly diagnosed glioblastoma multiforme. *J Clin Oncol* 2015;33:4145-4150.
- <sup>10</sup>Emami B, Lyman J, Brown A, et al. Tolerance of normal tissue to therapeutic irradiation. *Int J Radiat Oncol Biol Phys* 1991;21:109-122.
- <sup>11</sup>Pieters RS, Niemierko A, Fullerton BC, Munzenrider JE. Cauda equina tolerance to high-dose fractionated irradiation. *Int J Radiat Oncol Biol Phys* 2006;64:251-257.
- <sup>12</sup>Abdel-Wahab M, et al. Spinal cord gliomas: A multi-institutional retrospective analysis. *Int J Radiat Oncol Biol Phys*, 2006; 64:1060-1071.
- <sup>13</sup>Lee SH, Chung CK, Kim CH, et al. Long-term outcomes of surgical resection with or without adjuvant radiation therapy for treatment of spinal ependymoma: a retrospective multicenter study by the Korea Spinal Oncology Research Group. *Neuro Oncol* 2013;15:921-929.
- <sup>14</sup>Brown AP, Barney CL, Grosshans DR. Proton beam craniospinal irradiation reduces acute toxicity for adults with medulloblastoma. *Int J Radiat Oncol Biol Phys* 2013;86:277-284.
- <sup>15</sup>Packer RJ, Gajjar A, Vezina G, et al. Phase III study of craniospinal radiation therapy followed by adjuvant chemotherapy for newly diagnosed average-risk medulloblastoma. *J Clin Oncol* 2006;24:4202-4208.
- <sup>16</sup>Morris PG, Correa DD, Yahalom J, et al. Rituximab, methotrexate, procarbazine, and vincristine followed by consolidation reduced-dose whole-brain radiotherapy and cytarabine in newly diagnosed primary CNS lymphoma: final results and long-term outcome. *J Clin Oncol* 2013;31:3971-3979.
- <sup>17</sup>Ghesquière H, Ferlay C, Sebban C, et al. Long-term follow-up of an age-adapted C5R protocol followed by radiotherapy in 99 newly diagnosed primary CNS lymphomas: a prospective multicentric phase II study of the Groupe d'Etude des Lymphomes de l'Adulte (GELA). *Ann Oncol* 2010;4:842-850.
- <sup>18</sup>Shah GD, Yahalom J, Correa DD, et al. Combined immunochemotherapy with reduced whole-brain radiotherapy for newly diagnosed primary CNS lymphoma. *J Clin Oncol* 2007;30:4730-4735.
- <sup>19</sup>Thiel E, Korfel A, Martus P, et al. High-dose methotrexate with or without whole brain radiotherapy for primary CNS lymphoma (G-PCNSL-SG-1): a phase 3, randomised, non-inferiority trial. *Lancet Oncol* 2010;11:1036-1047.
- <sup>20</sup>Kasenda B, Loeffler J, Illerhaus G, et al. The role of whole brain radiation in primary CNS lymphoma. *Blood* 2016;1:32-36.
- <sup>21</sup>Andrews DW, Scott CB, Sperduto PW, et al. Whole brain radiation therapy with or without stereotactic radiosurgery boost for patients with one to three brain metastases: phase III results of the RTOG 9508 randomized trial. *Lancet* 2004;363:1665-1672.
- <sup>22</sup>Brown PD, Pugh S, Laack NN, Wefel JS, et al. Memantine for the prevention of cognitive dysfunction in patients receiving whole-brain radiotherapy: a randomized, double-blind, placebo-controlled trial. *Neuro Oncol* 2013;15:1429-37.
- <sup>23</sup>Gondi V, Pugh S, Brown PD, et al. NCOG-01. Preservation of neurocognitive function (NCF) with hippocampal avoidance during whole-brain radiotherapy (WBRT) for brain metastases: preliminary results of phase III trial NRG Oncology CC001. *Neuro Oncol* 2018;20:vi:172-vi:172.
- <sup>24</sup>Gondi V, Pugh SL, Tome WA, et al. Preservation of memory with conformal avoidance of the hippocampal neural stem-cell compartment during whole-brain radiotherapy for brain metastases (RTOG 0933): a phase II multi-institutional trial. *J Clin Oncol* 2014;32:3810-3816.

**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR RADIATION THERAPY

### REFERENCES

- <sup>25</sup>Aoyama H, Shirato H, Tago M, et al. Stereotactic radiosurgery plus whole-brain radiation therapy vs stereotactic radiosurgery alone for treatment of brain metastases: A randomized controlled trial. *JAMA* 2006;295:2483-2491.
- <sup>26</sup>Chang WS, Kim HY, Chang JW, et al. Analysis of radiosurgical results in patients with brain metastases according to the number of brain lesions: is stereotactic radiosurgery effective for multiple brain metastases? *J Neurosurg* 2010;113 Suppl:73-78.
- <sup>27</sup>Shaw E, Scott C, Souhami L, et al. Single dose radiosurgical treatment of recurrent previously irradiated primary brain tumors and brain metastases: final report of RTOG protocol 90-05. *Int J Radiat Oncol Biol Phys* 2000;47:291-298.
- <sup>28</sup>Yamamoto M, Serizawa T, Shuto T, et al. Stereotactic radiosurgery for patients with multiple brain metastases (JLGK0901): a multi-institutional prospective observational study. *Lancet Oncol* 2014;15:387-395.
- <sup>29</sup>Brown PD, Ballman KV, Cerhan JH, et al. Postoperative stereotactic radiosurgery compared with whole brain radiotherapy for resected metastatic brain disease (NCCTG N107C/CEC.3): a multicentre, randomised, controlled, phase 3 trial. *Lancet Oncol* 2017;18:1049-1060.
- <sup>30</sup>Mahajan A, Ahmed S, McAleer MF, et al. Post-operative stereotactic radiosurgery versus observation for completely resected brain metastases: a single-centre, randomised, controlled, phase 3 trial. *Lancet Oncol* 2017;18:1040-1048.
- <sup>31</sup>Soliman H, Ruschin M, Angelov L, et al. Consensus Contouring Guidelines for postoperative completely resected cavity stereotactic radiosurgery for brain metastases. *Int J Radiat Oncol Biol Phys* 2018;100:436-442.
- <sup>32</sup>Soltys SG, Seiger K, Modlin LA, et al. A phase I/II dose-escalation trial of 3-fraction stereotactic radiosurgery (SRS) for large resection cavities of brain metastases. *Int J Radiat Oncol Biol Phys* 2015;93:S38.
- <sup>33</sup>Cox BW, Spratt DE, Lovelock M, et al. International Spine Radiosurgery Consortium consensus guidelines for target volume definition in spinal stereotactic radiosurgery. *Int J Radiat Oncol Biol Phys* 2012;83:e597-e605.
- <sup>34</sup>Redmond KJ, Lo SS, Soltys SG, et al. Consensus guidelines for postoperative stereotactic body radiation therapy for spinal metastases: results of an international survey. *J Neurosurg Spine* 2017;26:299-306.
- <sup>35</sup>Redmond KJ, Robertson S, Lo SS, et al. Consensus contouring guidelines for postoperative stereotactic body radiation therapy for metastatic solid tumor malignancies to the spine. *Int J Radiat Oncol Biol Phys* 2017;97:64-74.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

#### ADULT LOW-GRADE GLIOMA/PILOCYTIC AND INFILTRATIVE SUPRATENTORIAL ASTROCYTOMA/OLIGODENDROGLIOMA

	Preferred Regimens	Other Recommended Regimens	Useful in Certain Circumstances
Adjuvant Treatment	<ul style="list-style-type: none"> <li>• RT + adjuvant PCV (category 1)<sup>1,2</sup></li> <li>• RT + concurrent and adjuvant temozolomide<sup>3-5</sup></li> <li>• RT + adjuvant temozolomide<sup>3-5</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Temozolomide<sup>b,3,4</sup></li> <li>• PCV<sup>b</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Pilocytic astrocytoma, PXA, ganglioglioma if <i>BRAF</i> V600E activation mutation               <ul style="list-style-type: none"> <li>▸ BRAF/MEK inhibitors:                   <ul style="list-style-type: none"> <li>◊ Dabrafenib/trametinib<sup>6,7</sup></li> <li>◊ Vemurafenib/cobimetinib<sup>8,9</sup></li> </ul> </li> </ul> </li> <li>• Subependymal giant cell astrocytoma               <ul style="list-style-type: none"> <li>▸ mTOR inhibitor (eg, everolimus)<sup>10,11</sup></li> </ul> </li> </ul>
Recurrent or Progressive Disease <sup>a</sup>	<ul style="list-style-type: none"> <li>• None</li> </ul>	<ul style="list-style-type: none"> <li>• RT + adjuvant PCV</li> <li>• RT + adjuvant temozolomide</li> <li>• RT+ concurrent and adjuvant temozolomide</li> <li>• Temozolomide<sup>c,4,12,13</sup></li> <li>• Lomustine or carmustine</li> <li>• PCV<sup>14</sup></li> <li>• Platinum-based regimens<sup>d,15-17</sup></li> </ul>	<ul style="list-style-type: none"> <li>• None</li> </ul>

<sup>a</sup>There are multiple reasonable options, but there is no uniform standard of care at this time for recurrent disease.

<sup>b</sup>In rare circumstances, treating a patient with chemotherapy without RT may be considered.

<sup>c</sup>For patients not previously treated.

<sup>d</sup>Platinum-based regimens include cisplatin or carboplatin.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

#### ANAPLASTIC GLIOMAS

	Preferred Regimens	Other Recommended Regimens	Useful in Certain Circumstances
<b>Adjuvant Treatment</b> Anaplastic oligodendroglioma (1p19q co-deleted) (KPS ≥60)	<ul style="list-style-type: none"> <li>• RT with adjuvant PCV (category 1)<sup>e,18</sup></li> <li>• RT with neoadjuvant PCV (category 1)<sup>e,19</sup></li> </ul>	<ul style="list-style-type: none"> <li>• RT with concurrent and adjuvant TMZ<sup>20</sup></li> <li>• RT with adjuvant TMZ<sup>21,22</sup></li> </ul>	<ul style="list-style-type: none"> <li>• None</li> </ul>
<b>Adjuvant Treatment</b> Anaplastic astrocytoma/anaplastic oligoastrocytoma, NOS <sup>f</sup> (KPS ≥60)	<ul style="list-style-type: none"> <li>• RT with concurrent and adjuvant TMZ<sup>23,24</sup></li> <li>• RT followed by adjuvant TMZ (12 cycles)<sup>24</sup></li> </ul>	<ul style="list-style-type: none"> <li>• RT with adjuvant PCV<sup>e,25,26</sup></li> <li>• RT with neoadjuvant PCV<sup>e</sup></li> </ul>	<ul style="list-style-type: none"> <li>• None</li> </ul>
<b>Adjuvant Treatment</b> Anaplastic gliomas (KPS <60)	<ul style="list-style-type: none"> <li>• None</li> </ul>	<ul style="list-style-type: none"> <li>• TMZ<sup>9</sup> (category 2B)<sup>27</sup></li> </ul>	<ul style="list-style-type: none"> <li>• None</li> </ul>
<b>Recurrence Therapy<sup>h</sup></b>	<ul style="list-style-type: none"> <li>• TMZ<sup>12,13,28,29</sup></li> <li>• Lomustine or carmustine<sup>30</sup></li> <li>• PCV<sup>31</sup></li> <li>• Bevacizumab<sup>i, 32-34</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Chemotherapy<sup>j</sup> + bevacizumab               <ul style="list-style-type: none"> <li>▶ Carmustine or lomustine + bevacizumab<sup>35</sup></li> <li>▶ TMZ + bevacizumab<sup>36</sup></li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• If failure or intolerance to the preferred or other recommended regimens               <ul style="list-style-type: none"> <li>▶ Etoposide<sup>37,38</sup> (category 2B)</li> <li>▶ Platinum-based regimens<sup>d,39-41</sup> (category 3)</li> </ul> </li> </ul>

<sup>d</sup>Platinum-based regimens include cisplatin or carboplatin.

<sup>e</sup>The panel recommends that PCV be administered after RT (as per EORTC 26951) since the intensive PCV regimen given prior to RT (RTOG 9402) was not tolerated as well.

<sup>f</sup>The WHO 2016 classification of CNS tumors has deleted oligoastrocytoma as a diagnostic category; however, oligoastrocytoma NOS and anaplastic oligoastrocytoma NOS may continue to be used for tumors that cannot be classified as either astrocytoma or oligodendroglioma due to the absence of appropriate molecular testing.

<sup>g</sup>Consider TMZ if tumor is MGMT promoter methylated.

<sup>h</sup>Strongly suggest consideration of clinical trials prior to treating recurrent disease with standard chemotherapy, as additional therapies may eliminate the majority of clinical trial options.

<sup>i</sup>Patients who have evidence of radiographic progression may benefit from continuation of bevacizumab to prevent rapid neurologic deterioration.

<sup>j</sup>Bevacizumab + chemotherapy can be considered if bevacizumab monotherapy fails and it is desirable to continue the steroid-sparing effects of bevacizumab.

**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

GLIOBLASTOMA			
	Preferred Regimens	Other Recommended Regimens	Useful in Certain Circumstances
Adjuvant Treatment	<ul style="list-style-type: none"> <li>• RT with concurrent and adjuvant TMZ<sup>42,43</sup> ± TTF<sup>44</sup></li> </ul>		<ul style="list-style-type: none"> <li>• RT with concurrent and adjuvant TMZ (for patients age 70 or younger and KPS &lt;60)<sup>45</sup></li> <li>• TMZ (for patients with MGMT promoter-methylated tumors and KPS &lt;60 or age &gt;70 years and KPS ≥60)<sup>42,46</sup></li> <li>• RT with concurrent and adjuvant lomustine and TMZ (for patients with MGMT promoter-methylated tumors, KPS ≥60, and age ≤70 years) (category 2B)<sup>k,47</sup></li> </ul>
Recurrence Therapy <sup>h,m</sup>	<ul style="list-style-type: none"> <li>• Bevacizumab<sup>i,l, 48-51</sup></li> <li>• Temozolomide<sup>13,29,52,53</sup></li> <li>• Lomustine or carmustine<sup>54-57</sup></li> <li>• PCV<sup>58,59</sup></li> <li>• Regorafenib<sup>60</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Chemotherapy<sup>j</sup> + bevacizumab               <ul style="list-style-type: none"> <li>▸ Carmustine or lomustine + bevacizumab<sup>61,62</sup></li> <li>▸ TMZ + bevacizumab<sup>63,64</sup></li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• If failure or intolerance to the preferred or other recommended regimens               <ul style="list-style-type: none"> <li>▸ Etoposide (category 2B)<sup>37</sup></li> <li>▸ Platinum-based regimens<sup>d</sup> (category 3)<sup>65,66</sup></li> </ul> </li> </ul>

<sup>d</sup>Platinum-based regimens include cisplatin or carboplatin.

<sup>h</sup>Strongly suggest consideration of clinical trials prior to treating recurrent disease with standard chemotherapy, as additional therapies may eliminate the majority of clinical trial options.

<sup>i</sup>Patients who have evidence of radiographic progression may benefit from continuation of bevacizumab to prevent rapid neurologic deterioration.

<sup>j</sup>Bevacizumab + chemotherapy can be considered if bevacizumab monotherapy fails and it is desirable to continue the steroid-sparing effects of bevacizumab.

<sup>k</sup>Moderate to significant myelosuppression was observed, but the toxicity profile for this regimen is not yet fully defined.

<sup>l</sup>An FDA-approved biosimilar is an appropriate substitute for bevacizumab.

<sup>m</sup>There are no identified targeted agents with demonstrated efficacy in glioblastoma. However, the panel encourages molecular testing of tumor because if a driver mutation is detected, it may be reasonable to treat with a targeted therapy on a compassionate use basis and/or the patient may have more treatment options in the context of a clinical trial. Molecular testing also has a valuable role in improving diagnostic accuracy and prognostic stratification that may inform treatment selection.

**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

#### ADULT INTRACRANIAL AND SPINAL EPENDYMOMA (EXCLUDING SUBEPENDYMOMA)

	<u>Preferred Regimens</u>	<u>Other Recommended Regimens</u>	<u>Useful in Certain Circumstances</u>
<b>Recurrence Therapy</b>	<ul style="list-style-type: none"> <li>• None</li> </ul>	<ul style="list-style-type: none"> <li>• Platinum-based regimens:<sup>d</sup> Single agent or combination<sup>67,68</sup></li> <li>• Etoposide<sup>69,70</sup></li> <li>• Lomustine or carmustine<sup>67</sup></li> <li>• Bevacizumab<sup>i,71</sup></li> <li>• TMZ<sup>72</sup></li> <li>• Lapatinib + TMZ (category 2B)<sup>73</sup></li> </ul>	<ul style="list-style-type: none"> <li>• None</li> </ul>

#### ADULT MEDULLOBLASTOMA

<b>Regimens following weekly vincristine<sup>n</sup> during craniospinal radiation therapy</b>	<ul style="list-style-type: none"> <li>• Cisplatin, cyclophosphamide, and vincristine<sup>n,74</sup></li> <li>• Cisplatin, lomustine, and vincristine<sup>n,74</sup></li> </ul>	<ul style="list-style-type: none"> <li>• None</li> </ul>	<ul style="list-style-type: none"> <li>• None</li> </ul>
<b>Recurrence Therapy</b>	<ul style="list-style-type: none"> <li>• None</li> </ul>	<ul style="list-style-type: none"> <li>• No prior chemotherapy               <ul style="list-style-type: none"> <li>▸ High-dose cyclophosphamide ± etoposide</li> <li>▸ Carboplatin, etoposide, and cyclophosphamide</li> <li>▸ Cisplatin, etoposide, and cyclophosphamide</li> </ul> </li> <li>• Prior chemotherapy               <ul style="list-style-type: none"> <li>▸ High-dose cyclophosphamide ± etoposide</li> <li>▸ Oral etoposide<sup>75,76</sup></li> <li>▸ TMZ<sup>12,77</sup></li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• Consider high-dose chemotherapy with autologous stem cell reinfusion<sup>78</sup> in patients who achieve a CR with conventional doses of chemotherapy or have no residual disease after re-resection</li> <li>• Vismodegib (for mutations in the sonic hedgehog pathway and if prior chemotherapy)<sup>79</sup></li> </ul>

<sup>d</sup>Platinum-based regimens include cisplatin or carboplatin.

<sup>i</sup>Patients who have evidence of radiographic progression may benefit from continuation of bevacizumab to prevent rapid neurologic deterioration.

<sup>n</sup>Omission of vincristine during radiotherapy phase of therapy or dose modification may be required for adults because they do not tolerate this regimen as well. Data supporting vincristine's use have been found in pediatric trials only. Patients should be monitored closely for neurologic toxicity with periodic exams.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

#### PRIMARY CNS LYMPHOMA

	Preferred Regimens	Other Recommended Regimens	Useful in Certain Circumstances
<b>Induction Therapy:</b>	<ul style="list-style-type: none"> <li>• <b>Systemic therapy</b> <ul style="list-style-type: none"> <li>▸ High-dose methotrexate 8 g/m<sup>2</sup> combined with the following:<sup>o,80</sup> <ul style="list-style-type: none"> <li>◊ Rituximab<sup>p,81-84</sup></li> <li>◊ Rituximab and TMZ<sup>p,85</sup></li> </ul> </li> <li>▸ High-dose methotrexate 3.5 g/m<sup>2</sup> combined with the following, and consider WBRT:<sup>o,q</sup> <ul style="list-style-type: none"> <li>◊ Vincristine, procarbazine, and rituximab (R-MPV)<sup>86-89</sup></li> <li>◊ TMZ + rituximab followed by post-RT TMZ<sup>90</sup></li> </ul> </li> </ul> </li> </ul>	NONE	<ul style="list-style-type: none"> <li>• <b>Intra-CSF therapy</b> <ul style="list-style-type: none"> <li>▸ If CSF positive or spinal MRI positive <ul style="list-style-type: none"> <li>◊ Methotrexate</li> <li>◊ Cytarabine</li> <li>◊ Rituximab<sup>91</sup></li> </ul> </li> </ul> </li> <li>• <b>Patient is unsuitable for or intolerant to high-dose methotrexate</b> <ul style="list-style-type: none"> <li>▸ See "Other Recommended Regimens" for Relapsed or Refractory Disease</li> </ul> </li> </ul>
<b>Consolidation Therapy:</b>	<ul style="list-style-type: none"> <li>• <b>High-dose chemotherapy with stem cell rescue</b> <ul style="list-style-type: none"> <li>▸ Carmustine + thiotepa<sup>92,93</sup></li> <li>▸ Thiotepa, busulfan, and cyclophosphamide (TBC)<sup>94</sup></li> </ul> </li> <li>• High-dose cytarabine + etoposide (EA)<sup>95</sup></li> <li>• High-dose cytarabine<sup>86-88</sup></li> </ul>	NONE	NONE
<b>Relapsed or Refractory Disease:</b>	NONE	<ul style="list-style-type: none"> <li>• Retreat with high-dose methotrexate<sup>o,80</sup> <ul style="list-style-type: none"> <li>▸ With or without rituximab<sup>p</sup></li> <li>▸ With rituximab<sup>p</sup> and ibrutinib<sup>r,96</sup></li> </ul> </li> <li>• Ibrutinib<sup>r,96,97</sup></li> <li>• TMZ<sup>98</sup></li> <li>• Rituximab ± TMZ<sup>99-101</sup></li> <li>• Lenalidomide with or without rituximab<sup>102</sup></li> <li>• High-dose cytarabine<sup>103</sup></li> <li>• Pemetrexed<sup>104</sup></li> <li>• Pomalidomide<sup>105</sup></li> </ul>	<ul style="list-style-type: none"> <li>▸ Consider high-dose chemotherapy with autologous stem cell reinfusion in eligible patients<sup>92,106,107</sup></li> </ul>

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

#### PRIMARY CNS LYMPHOMA

#### FOOTNOTES

<sup>o</sup>Consider glucarpidase (carboxypeptidase G2) for prolonged methotrexate clearance due to methotrexate-induced renal toxicity. Ramsey LB, Balis FM, O'Brien MM, et al. Consensus guideline for use of glucarpidase in patients with high-dose methotrexate induced acute kidney injury and delayed methotrexate clearance. *Oncologist* 2018;23:52-61.

<sup>p</sup>Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen. If positive, check viral load and consult with gastroenterologist. [The NCCN Guidelines for B-Cell Lymphomas \(NHODG-B, 2 of 4\)](#) also have information about HBV testing for patients considering rituximab.

<sup>q</sup>Other combinations with methotrexate may be used.

<sup>r</sup>Ibrutinib is associated with risk of aspergillus infection.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

#### MENINGIOMAS

<u>Preferred Regimens</u>	<u>Other Recommended Regimens</u>	<u>Useful in Certain Circumstances</u>
<ul style="list-style-type: none"> <li>• None</li> </ul>	<ul style="list-style-type: none"> <li>• Sunitinib (category 2B)<sup>108</sup></li> <li>• Bevacizumab<sup>i,109,110</sup></li> <li>• Bevacizumab<sup>i</sup> + everolimus (category 2B)<sup>111</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Somatostatin analogue (category 2B)<sup>112</sup></li> </ul>

<sup>i</sup>Patients who have evidence of radiographic progression may benefit from continuation of bevacizumab to prevent rapid neurologic deterioration.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

#### BRAIN METASTASES

- **Tumor agnostic<sup>s</sup>**
  - ▶ ***NTRK* gene-fusion tumors**
    - ◊ Larotrectinib<sup>113,114</sup>
    - ◊ Entrectinib<sup>115,116</sup>
  - ▶ Temozolomide 5/28 schedule
- **Breast Cancer<sup>u</sup>**
  - ▶ **HER2-positive**
    - ◊ Capecitabine + lapatinib<sup>117,118</sup>
    - ◊ Capecitabine + neratinib<sup>119,120</sup>
    - ◊ Paclitaxel + neratinib (category 2B)<sup>121</sup>
    - ◊ Capecitabine (category 2B)<sup>122-126</sup>
    - ◊ Tucatinib + trastuzumab + capecitabine (if previously treated with 1 or more anti-HER2-based regimens)<sup>127</sup>
  - ▶ **HER2 non-specific**
    - ◊ Cisplatin (category 2B)<sup>128,129</sup>
    - ◊ Etoposide (category 2B)<sup>128,129</sup>
    - ◊ Cisplatin + etoposide (category 2B)<sup>129,130</sup>
    - ◊ High-dose methotrexate (category 2B)<sup>m,131</sup>
- **Melanoma<sup>u</sup>**
  - ▶ ***BRAF* V600E positive**
    - ◊ Dabrafenib<sup>132-134</sup>/trametinib<sup>135</sup>
    - ◊ Vemurafenib<sup>136,137</sup>/cobimetinib<sup>t</sup> (category 2B)
  - ▶ ***BRAF* non-specific**
    - ◊ Ipilimumab + nivolumab (preferred)<sup>138-140</sup>
    - ◊ Ipilimumab<sup>141</sup>
    - ◊ Nivolumab<sup>139</sup>
    - ◊ Pembrolizumab<sup>142</sup>
- **Non-Small Cell Lung Cancer<sup>u</sup>**
  - ▶ **EGFR-sensitizing mutation positive**
    - ◊ Osimertinib (EGFR T790M positive)<sup>143-145</sup>
    - ◊ Pulsatile erlotinib<sup>146-148</sup>
    - ◊ Afatinib (category 2B)<sup>149</sup>
    - ◊ Gefitinib (category 2B)<sup>150,151</sup>
  - ▶ **ALK rearrangement positive**
    - ◊ Brigatinib<sup>152,153</sup>
    - ◊ Alectinib<sup>154,155</sup>
    - ◊ Ceritinib<sup>156</sup>
  - ▶ **ALK rearrangement positive or ROS1 positive**
    - ◊ Crizotinib (category 2B)<sup>157</sup>
  - ▶ **PD-L1 positive**
    - ◊ Pembrolizumab<sup>142,158</sup>
    - ◊ Nivolumab<sup>159-161</sup>
- **Small Cell Lung Cancer<sup>u</sup>**
  - ◊ Topotecan (category 2B)
- **Lymphoma<sup>u</sup>**
  - ◊ High-dose methotrexate<sup>162</sup>

<sup>s</sup>See the appropriate NCCN treatment guidelines for systemic therapy recommendations for newly diagnosed brain metastases for any cancers not listed here.

<sup>t</sup>Although there are no published prospective studies on the combination of vemurafenib and cobimetinib for melanoma patients with brain metastases, there is high-quality evidence that for melanoma with distant metastasis, combination therapy with vemurafenib and cobimetinib is associated with improved outcomes and safety compared with single-agent vemurafenib.

<sup>u</sup>Use active agents against primary tumor.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY

#### LEPTOMENINGEAL AND SPINE METASTASES

##### • Treatment

▶ Systemic therapy specific to primary cancer type; emphasizing drugs with good CNS penetration

▶ Intra-CSF chemotherapy<sup>163</sup>

◊ Thiotepe<sup>164</sup>

◊ Topotecan<sup>165</sup>

◊ Etoposide<sup>166</sup>

◊ Cytarabine<sup>167-170</sup>

◊ Methotrexate<sup>169,171-173</sup>

▶ Lymphoma

◊ Intra-CSF chemotherapy

– Rituximab<sup>168</sup>

◊ High-dose methotrexate<sup>o,162</sup>

▶ Breast cancer

◊ Intra-CSF chemotherapy

– Methotrexate<sup>169,171,172</sup>

– Trastuzumab (*HER2*-positive)<sup>174</sup>

◊ High-dose methotrexate<sup>o,131,175,176</sup>

▶ Non-small cell lung cancer

◊ Osimertinib (EGFR mutation positive)<sup>177,178</sup>

◊ Weekly pulse erlotinib for (EGFR exon 19 deletion or exon 21 L858R mutation) (category 2B)<sup>146</sup>

##### Metastatic Spine Tumors

• Use regimen for disease-specific site

<sup>o</sup>Consider glucarpidase (carboxypeptidase G2) for prolonged methotrexate clearance due to methotrexate-induced renal toxicity. Ramsey LB, Balis FM, O'Brien MM, et al. Consensus guideline for use of glucarpidase in patients with high-dose methotrexate induced acute kidney injury and delayed methotrexate clearance. *Oncologist* 2018;23:52-61.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY**  
**REFERENCES**

- <sup>1</sup>Buckner JC, Shaw EG, Pugh SL, et al. Radiation plus procarbazine, CCNU, and vincristine in low-grade glioma. *N Engl J Med* 2016;374:1344-1355.
- <sup>2</sup>Shaw EG, Wang M, Coons SW, et al. Randomized trial of radiation therapy plus procarbazine, lomustine, and vincristine chemotherapy for supratentorial adult low-grade glioma: initial results of RTOG 9802. *J Clin Oncol* 2012;30:3065-3070.
- <sup>3</sup>Pouratian N, Gasco J, Sherman JH, et al. Toxicity and efficacy of protracted low dose temozolomide for the treatment of low grade gliomas. *J Neurooncol* 2007;82:281-288.
- <sup>4</sup>Kesari S, Schiff D, Drappatz J, et al. Phase II study of protracted daily temozolomide for low-grade gliomas in adults. *Clin Cancer Res* 2009;15:330-337.
- <sup>5</sup>Fisher BJ, Hu C, Macdonald DR, et al. Phase 2 study of temozolomide-based chemoradiation therapy for high-risk low-grade gliomas: preliminary results of Radiation Therapy Oncology Group 0424. *Int J Radiat Oncol Biol Phys* 2015;91:497-504.
- <sup>6</sup>Brown NF, Carter T, Kitchen N, Mulholland P. Dabrafenib and trametinib in BRAFV600E mutated glioma. *CNS Oncol* 2017;6:291-296.
- <sup>7</sup>Marks AM, Bindra RS, DiLuna ML, et al. Response to the BRAF/MEK inhibitors dabrafenib/trametinib in an adolescent with a BRAF V600E mutated anaplastic ganglioglioma intolerant to vemurafenib. *Pediatr Blood Cancer* 2018;65:e26969.
- <sup>8</sup>Hyman DM, Puzanov I, Subbiah V, et al. Vemurafenib in multiple nonmelanoma cancers with BRAF V600 mutations. *N Engl J Med* 2015;373:726-736.
- <sup>9</sup>Kaley T, Touat M, Subbiah V, et al. BRAF inhibition in BRAF(V600)-mutant gliomas: results from the VE-BASKET Study. *J Clin Oncol* 2018;JCO2018789990.
- <sup>10</sup>Franz DN, Belousova E, Sparagana S, et al. Everolimus for subependymal giant cell astrocytoma in patients with tuberous sclerosis complex: 2-year open-label extension of the randomised EXIST-1 study. *Lancet Oncol* 2014;15:1513-1520.
- <sup>11</sup>Franz DN, Belousova E, Sparagana S, et al. Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial. *Lancet* 2013;381:125-132.
- <sup>12</sup>Nicholson HS, Kretschmar CS, Krailo M, et al. Phase 2 study of temozolomide in children and adolescents with recurrent central nervous system tumors: a report from the Children's Oncology Group. *Cancer* 2007;110:1542-1550.
- <sup>13</sup>Perry JR, Rizek P, Cashman R, et al. Temozolomide rechallenge in recurrent malignant glioma by using a continuous temozolomide schedule: the "rescue" approach. *Cancer* 2008;113:2152-2157.
- <sup>14</sup>Triebels VH, Taphoorn MJ, Brandes AA, et al. Salvage PCV chemotherapy for temozolomide-resistant oligodendrogliomas. *Neurology* 2004;63:904-906.
- <sup>15</sup>Massimino M, Spreafico F, Riva D, et al. A lower-dose, lower-toxicity cisplatin-etoposide regimen for childhood progressive low-grade glioma. *J Neurooncol* 2010;100:65-71.
- <sup>16</sup>Moghrabi A, Friedman HS, Ashley DM, et al. Phase II study of carboplatin (CBDCA) in progressive low-grade gliomas. *Neurosurg Focus* 1998;4:e3.
- <sup>17</sup>Brandes AA, Basso U, Vastola F, et al. Carboplatin and teniposide as third-line chemotherapy in patients with recurrent oligodendroglioma or oligoastrocytoma: a phase II study. *Ann Oncol* 2003;14:1727-1731.
- <sup>18</sup>van den Bent MJ, Brandes AA, Taphoorn MJ, et al. Adjuvant procarbazine, lomustine, and vincristine chemotherapy in newly diagnosed anaplastic oligodendroglioma: long-term follow-up of EORTC brain tumor group study 26951. *J Clin Oncol* 2013;31:344-350.
- <sup>19</sup>Cairncross G, Wang M, Shaw E, et al. Phase III trial of chemoradiotherapy for anaplastic oligodendroglioma: long-term results of RTOG 9402. *J Clin Oncol* 2013;31:337-343.
- <sup>20</sup>Jaecle K, Vogelbaum M, Ballman K, et al. CODEL (Alliance-N0577; EORTC-26081/22086; NRG-1071; NCIC-CEC-2): phase III randomized study of RT vs. RT+TMZ vs. TMZ for newly diagnosed 1p/19q-codeleted anaplastic oligodendroglial tumors. *Neurology* 2016;86.
- <sup>21</sup>Panageas KS, Iwamoto FM, Cloughesy TF, et al. Initial treatment patterns over time for anaplastic oligodendroglial tumors. *Neuro Oncol* 2012;14:761-767.
- <sup>22</sup>Shin JY, Diaz AZ. Utilization and impact of adjuvant therapy in anaplastic oligodendroglioma: an analysis on 1692 patients. *J Neurooncol* 2016;129:567-575.
- <sup>23</sup>Brandes AA, Nicolardi L, Tosoni A, et al. Survival following adjuvant PCV or temozolomide for anaplastic astrocytoma. *Neuro Oncol* 2006;8:253-260.
- <sup>24</sup>van den Bent MJ, Baumert B, Erridge SC, et al. Interim results from the CATNON trial (EORTC study 26053-22054) of treatment with concurrent and adjuvant temozolomide for 1p/19q non-co-deleted anaplastic glioma: a phase 3, randomised, open-label intergroup study. *Lancet* 2017;390:1645-1653.
- <sup>25</sup>Levin VA, Hess KR, Choucair A, et al. Phase III randomized study of postradiotherapy chemotherapy with combination alpha-difluoromethylornithine-PCV versus PCV for anaplastic gliomas. *Clin Cancer Res* 2003;9:981-990.
- <sup>26</sup>Levin VA, Silver P, Hannigan J, et al. Superiority of post-radiotherapy adjuvant chemotherapy with CCNU, procarbazine, and vincristine (PCV) over BCNU for anaplastic gliomas: NCOG 6G61 final report. *Int J Radiat Oncol Biol Phys* 1990;18:321-324.
- <sup>27</sup>Ducray F, del Rio MS, Carpentier C, et al. Up-front temozolomide in elderly patients with anaplastic oligodendroglioma and oligoastrocytoma. *J Neurooncol* 2011;101:457-462.
- <sup>28</sup>Yung WK, Prados MD, Yaya-Tur R, et al. Multicenter phase II trial of temozolomide in patients with anaplastic astrocytoma or anaplastic oligoastrocytoma at first relapse. *Temodal Brain Tumor Group. J Clin Oncol* 1999;17:2762-2771.
- <sup>29</sup>Perry JR, Belanger K, Mason WP, et al. Phase II trial of continuous dose-intense temozolomide in recurrent malignant glioma: RESCUE study. *J Clin Oncol* 2010;28:2051-2057.
- <sup>30</sup>Prados M, Rodriguez L, Chamberlain M, et al. Treatment of recurrent gliomas with 1,3-bis (2-chloroethyl)-1-nitrosourea and alpha-difluoromethylornithine. *Neurosurgery* 1989;24:806-809.
- <sup>31</sup>Soffietti R, Ruda R, Bradac GB, Schiffer D. PCV chemotherapy for recurrent oligodendrogliomas and oligoastrocytomas. *Neurosurgery* 1998;43:1066-1073.
- <sup>32</sup>Norden AD, Young GS, Setayesh K, et al. Bevacizumab for recurrent malignant gliomas: efficacy, toxicity, and patterns of recurrence. *Neurology* 2008;70:779-787.
- <sup>33</sup>Chamberlain MC, Johnston S. Bevacizumab for recurrent alkylator-refractory anaplastic oligodendroglioma. *Cancer* 2009;115:1734-1743.
- <sup>34</sup>Chamberlain MC, Johnston S. Salvage chemotherapy with bevacizumab for recurrent alkylator-refractory anaplastic astrocytoma. *J Neurooncol* 2009;91:359-367.

**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



**PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY**  
**REFERENCES**

- <sup>35</sup>Vaccaro V, Fabi A, Vidiri A, et al. Activity and safety of bevacizumab plus fotemustine for recurrent malignant gliomas. *Biomed Res Int* 2014;2014:351252.
- <sup>36</sup>Liu Y, Feng F, Ji P, et al. Improvement of health related quality of life in patients with recurrent glioma treated with bevacizumab plus daily temozolomide as the salvage therapy. *Clin Neurol Neurosurg* 2018;169:64-70.
- <sup>37</sup>Fulton D, Urtasun R, Forsyth P. Phase II study of prolonged oral therapy with etoposide (VP16) for patients with recurrent malignant glioma. *J Neurooncol* 1996;27:149-155.
- <sup>38</sup>Leonard A, Wolff JE. Etoposide improves survival in high-grade glioma: a meta-analysis. *Anticancer Res* 2013;33:3307-3315.
- <sup>39</sup>Murray LJ, Bridgewater CH, Levy D. Carboplatin chemotherapy in patients with recurrent high-grade glioma. *Clin Oncol (R Coll Radiol)* 2011;23:55-61.
- <sup>40</sup>Roci E, Cakani B, Brace G, et al. Platinum-based chemotherapy in recurrent high-grade glioma patients: retrospective study. *Med Arch* 2014;68:140-143.
- <sup>41</sup>Soffietti R, Nobile M, Ruda R, et al. Second-line treatment with carboplatin for recurrent or progressive oligodendroglial tumors after PCV (procarbazine, lomustine, and vincristine) chemotherapy: a phase II study. *Cancer* 2004;100:807-813.
- <sup>42</sup>Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 2005;352:987-996.
- <sup>43</sup>Stupp R, Hegi ME, Mason WP, et al. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol* 2009;10:459-466.
- <sup>44</sup>Stupp R, Taillibert S, Kanner A, et al. Effect of tumor-treating fields plus maintenance temozolomide vs maintenance temozolomide alone on survival in patients with glioblastoma: a randomized clinical trial. *JAMA* 2017;318:2306-2316.
- <sup>45</sup>Muni R, Minniti G, Lanzetta G, et al. Short-term radiotherapy followed by adjuvant chemotherapy in poor-prognosis patients with glioblastoma. *Tumori* 2010;96:60-64.
- <sup>46</sup>Malmstrom A, Gronberg BH, Marosi C, et al. Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised, phase 3 trial. *Lancet Oncol* 2012;13:916-926.
- <sup>47</sup>Herrlinger U, Tzaridis T, Mack F, et al. Lomustine-temozolomide combination therapy versus standard temozolomide therapy in patients with newly diagnosed glioblastoma with methylated MGMT promoter (CeTeG/NOA-09): a randomised, open-label, phase 3 trial. *Lancet* 2019;393:678-688.
- <sup>48</sup>Cloughesy T, Prados MD, Mikkelsen T. A phase 2 randomized non-comparative clinical trial of the effect of bevacizumab alone or in combination with irinotecan on 6-month progression free survival in recurrent refractory glioblastoma [abstract]. *J Clin Oncol* 2008;26(Suppl 15):2010b.
- <sup>49</sup>Friedman HS, Prados MD, Wen PY, et al. Bevacizumab alone and in combination with irinotecan in recurrent glioblastoma. *J Clin Oncol* 2009;27:4733-4740.
- <sup>50</sup>Kreisl TN, Kim L, Moore K, et al. Phase II trial of single-agent bevacizumab followed by bevacizumab plus irinotecan at tumor progression in recurrent glioblastoma. *J Clin Oncol* 2009;27:740-745.
- <sup>51</sup>Vredenburgh JJ, Desjardins A, Herndon JE, 2nd, et al. Phase II trial of bevacizumab and irinotecan in recurrent malignant glioma. *Clin Cancer Res* 2007;13:1253-1259.
- <sup>52</sup>Yung WK, Albright RE, Olson J, et al. A phase II study of temozolomide vs. procarbazine in patients with glioblastoma multiforme at first relapse. *Br J Cancer* 2000;83:588-593.
- <sup>53</sup>Weller M, Tabatabai G, Kastner B, et al. MGMT promoter methylation is a strong prognostic biomarker for benefit from dose-intensified temozolomide rechallenge in progressive glioblastoma: the DIRECTOR trial. *Clin Cancer Res* 2015;21:2057-2064.
- <sup>54</sup>Wick W, Puduvalli VK, Chamberlain MC, et al. Phase III study of enzastaurin compared with lomustine in the treatment of recurrent intracranial glioblastoma. *J Clin Oncol* 2010;28:1168-1174.
- <sup>55</sup>Brandes AA, Tosoni A, Amista P, et al. How effective is BCNU in recurrent glioblastoma in the modern era? A phase II trial. *Neurology* 2004;63:1281-1284.
- <sup>56</sup>Reithmeier T, Graf E, Piroth T, et al. BCNU for recurrent glioblastoma multiforme: efficacy, toxicity and prognostic factors. *BMC Cancer* 2010;10:30.
- <sup>57</sup>Taal W, Oosterkamp HM, Walenkamp AM, et al. Single-agent bevacizumab or lomustine versus a combination of bevacizumab plus lomustine in patients with recurrent glioblastoma (BELOB trial): a randomised controlled phase 2 trial. *Lancet Oncol* 2014;15:943-953.
- <sup>58</sup>Carvalho BF, Fernandes AC, Almeida DS, et al. Second-line chemotherapy in recurrent glioblastoma: a 2-cohort study. *Oncol Res Treat* 2015;38:348-354.
- <sup>59</sup>Schmidt F, Fischer J, Herrlinger U, et al. PCV chemotherapy for recurrent glioblastoma. *Neurology* 2006;66:587-589.
- <sup>60</sup>Lombardi G, DeSalvo GL, Brandes AA, et al. Regorafenib compared with lomustine in patients with relapsed glioblastoma (REGOMA): a multicentre, open-label, randomised, controlled, phase 2 trial. *Lancet Oncol* 2019;20(1):110-119.
- <sup>61</sup>Wick W, Gorlia T, Bendszus M, et al. Lomustine and bevacizumab in progressive glioblastoma. *N Engl J Med* 2017;377:1954-1963.
- <sup>62</sup>Soffietti R, Trevisan E, Bertero L, et al. Bevacizumab and fotemustine for recurrent glioblastoma: a phase II study of AINO (Italian Association of Neuro-Oncology). *J Neurooncol* 2014;116:533-541.
- <sup>63</sup>Badruddoja MA, Pazzi M, Sanan A, et al. Phase II study of bi-weekly temozolomide plus bevacizumab for adult patients with recurrent glioblastoma. *Cancer Chemother Pharmacol* 2017;80:715-721.
- <sup>64</sup>Desjardins A, Reardon DA, Coan A, et al. Bevacizumab and daily temozolomide for recurrent glioblastoma. *Cancer* 2012;118:1302-1312.
- <sup>65</sup>Field KM, Simes J, Nowak AK, et al. Randomized phase 2 study of carboplatin and bevacizumab in recurrent glioblastoma. *Neuro Oncol* 2015;17:1504-1513.
- <sup>66</sup>Kaloshi G, Diamandi P, Cakani B, et al. The added value of bevacizumab concomitantly administered with carboplatin versus carboplatin alone in patients with recurrent glioblastomas. *Tumori* 2015;101:41-45.
- <sup>67</sup>Gornet MK, Buckner JC, Marks RS, et al. Chemotherapy for advanced CNS ependymoma. *J Neurooncol* 1999;45:61-67.
- <sup>68</sup>Brandes AA, Cavallo G, Reni M, et al. A multicenter retrospective study of chemotherapy for recurrent intracranial ependymal tumors in adults by the Gruppo Italiano Cooperativo di Neuro-Oncologia. *Cancer* 2005;104:143-148.
- <sup>69</sup>Chamberlain MC. Recurrent intracranial ependymoma in children: salvage therapy with oral etoposide. *Pediatr Neurol* 2001;24:117-121.

**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY**  
**REFERENCES**

- <sup>70</sup>Sandri A, Massimino M, Mastrodicasa L, et al. Treatment with oral etoposide for childhood recurrent ependymomas. *J Pediatr Hematol Oncol* 2005;27:486-490.
- <sup>71</sup>Green RM, Cloughesy TF, Stupp R, et al. Bevacizumab for recurrent ependymoma. *Neurology* 2009;73:1677-1680.
- <sup>72</sup>Ruda R, Bosa C, Magistrello M, et al. Temozolomide as salvage treatment for recurrent intracranial ependymomas of the adult: a retrospective study. *Neuro Oncol* 2016;18:261-268.
- <sup>73</sup>Gilbert M, Yuan Y, Wani K, et al. AT-23 a phase II study of lapatinib and dose-dense temozolomide (TMZ) for adults with recurrent ependymoma: a CERN clinical trial. *Neuro-Oncology* 2014;16:v13-v13.
- <sup>74</sup>Packer RJ, Gajjar A, Vezina G, et al. Phase III study of craniospinal radiation therapy followed by adjuvant chemotherapy for newly diagnosed average-risk medulloblastoma. *J Clin Oncol* 2006;24:4202-4208.
- <sup>75</sup>Ashley DM, Meier L, Kerby T, et al. Response of recurrent medulloblastoma to low-dose oral etoposide. *J Clin Oncol* 1996;14:1922-1927.
- <sup>76</sup>Chamberlain MC, Kormanik PA. Chronic oral VP-16 for recurrent medulloblastoma. *Pediatr Neurol* 1997;17:230-234.
- <sup>77</sup>Wang CH, Hsu TR, Wong TT, Chang KP. Efficacy of temozolomide for recurrent embryonal brain tumors in children. *Childs Nerv Syst* 2009;25:535-541.
- <sup>78</sup>Dunkel IJ, Gardner SL, Garvin JH, Jr., et al. High-dose carboplatin, thiopeta, and etoposide with autologous stem cell rescue for patients with previously irradiated recurrent medulloblastoma. *Neuro Oncol* 2010;12:297-303.
- <sup>79</sup>Robinson GW, Orr BA, Wu G, et al. Vismodegib exerts targeted efficacy against recurrent sonic hedgehog-subgroup medulloblastoma: results from phase II Pediatric Brain Tumor Consortium Studies PBTC-025B and PBTC-032. *J Clin Oncol* 2015;33:2646-2654.
- <sup>80</sup>Batchelor T, Carson K, O'Neill A, et al. Treatment of primary CNS lymphoma with methotrexate and deferred radiotherapy: a report of NABTT 96-07. *J Clin Oncol* 2003;21:1044-1049.
- <sup>81</sup>Chamberlain MC, Johnston SK. High-dose methotrexate and rituximab with deferred radiotherapy for newly diagnosed primary B-cell CNS lymphoma. *Neuro Oncol* 2010;12:736-744.
- <sup>82</sup>Gregory G, Arumugaswamy A, Leung T, et al. Rituximab is associated with improved survival for aggressive B cell CNS lymphoma. *Neuro Oncol* 2013;15:1068-1073.
- <sup>83</sup>Holdhoff M, Ambady P, Abdelaziz A, et al. High-dose methotrexate with or without rituximab in newly diagnosed primary CNS lymphoma. *Neurology* 2014;83:235-239.
- <sup>84</sup>Ly KI, Crew LL, Graham CA, Mrugala MM. Primary central nervous system lymphoma treated with high-dose methotrexate and rituximab: A single-institution experience. *Oncol Lett* 2016;11:3471-3476.
- <sup>85</sup>Wieduwilt MJ, Valles F, Issa S, et al. Immunochemotherapy with intensive consolidation for primary CNS lymphoma: a pilot study and prognostic assessment by diffusion-weighted MRI. *Clin Cancer Res* 2012;18:1146-1155.
- <sup>86</sup>DeAngelis LM, Seiferheld W, Schold SC, et al. Combination chemotherapy and radiotherapy for primary central nervous system lymphoma: Radiation Therapy Oncology Group Study 93-10. *J Clin Oncol* 2002;20:4643-4648.
- <sup>87</sup>Shah GD, Yahalom J, Correa DD, et al. Combined immunochemotherapy with reduced whole-brain radiotherapy for newly diagnosed primary CNS lymphoma. *J Clin Oncol* 2007;25:4730-4735.
- <sup>88</sup>Gavrilovic IT, Hormigo A, Yahalom J, et al. Long-term follow-up of high-dose methotrexate-based therapy with and without whole brain irradiation for newly diagnosed primary CNS lymphoma. *J Clin Oncol* 2006;24:4570-4574.
- <sup>89</sup>Omuro A, Correa DD, DeAngelis LM, et al. R-MPV followed by high-dose chemotherapy with TBC and autologous stem-cell transplant for newly diagnosed primary CNS lymphoma. *Blood* 2015;125:1403-1410.
- <sup>90</sup>Glass J, Won M, Schultz CJ, et al. Phase I and II study of induction chemotherapy with methotrexate, rituximab, and temozolomide, followed by whole-brain radiotherapy and postirradiation temozolomide for primary CNS lymphoma: NRG Oncology RTOG 0227. *J Clin Oncol* 2016;34:1620-1625.
- <sup>91</sup>Schulz H, Pels H, Schmidt-Wolf I, et al. Intraventricular treatment of relapsed central nervous system lymphoma with the anti-CD20 antibody rituximab. *Haematologica* 2004;89:753-754.
- <sup>92</sup>Illerhaus G, Muller F, Feuerhake F, et al. High-dose chemotherapy and autologous stem-cell transplantation without consolidating radiotherapy as first-line treatment for primary lymphoma of the central nervous system. *Haematologica* 2008;93:147-148.
- <sup>93</sup>Kasenda B, Schorb E, Fritsch K, et al. Prognosis after high-dose chemotherapy followed by autologous stem-cell transplantation as first-line treatment in primary CNS lymphoma--a long-term follow-up study. *Ann Oncol* 2012;23:2670-2675.
- <sup>94</sup>DeFilipp Z, Li S, El-Jawahri A, et al. High-dose chemotherapy with thiopeta, busulfan, and cyclophosphamide and autologous stem cell transplantation for patients with primary central nervous system lymphoma in first complete remission. *Cancer* 2017;123:3073-3079.
- <sup>95</sup>Rubenstein JL, Hsi ED, Johnson JL, et al. Intensive chemotherapy and immunotherapy in patients with newly diagnosed primary CNS lymphoma: CALGB 50202 (Alliance 50202). *J Clin Oncol* 2013;31:3061-3068.
- <sup>96</sup>Soussain C, Choquet S, Blonski M, et al. Ibrutinib monotherapy for relapse or refractory primary CNS lymphoma and primary vitreoretinal lymphoma: final analysis of the phase II 'proof-of-concept' iLOC study by the Lymphoma study association (LYSA) and the French oculo-cerebral lymphoma (LOC) network. *Eur J Cancer* 2019;117:121-130.
- <sup>97</sup>Grommes C, Pastore A, Palaskas N, et al. Ibrutinib unmasks critical role of bruton tyrosine kinase in primary CNS lymphoma. *Cancer Discov* 2017;7:1018-1029.
- <sup>98</sup>Makino K, Nakamura H, Hide T, Kuratsu J. Salvage treatment with temozolomide in refractory or relapsed primary central nervous system lymphoma and assessment of the MGMT status. *J Neurooncol* 2012;106:155-160.
- <sup>99</sup>Enting RH, Demopoulos A, DeAngelis LM, Abrey LE. Salvage therapy for primary CNS lymphoma with a combination of rituximab and temozolomide. *Neurology* 2004;63:901-903.
- <sup>100</sup>Batchelor TT, Grossman SA, Mikkelsen T, et al. Rituximab monotherapy for patients with recurrent primary CNS lymphoma. *Neurology* 2011;76:929-930.
- <sup>101</sup>Nayak L, Abrey LE, Drappatz J, et al. Multicenter phase II study of rituximab and temozolomide in recurrent primary central nervous system lymphoma. *Leuk Lymphoma* 2013;54:58-61.
- <sup>102</sup>Rubenstein JL, Geng H, Fraser EJ, et al. Phase 1 investigation of lenalidomide/rituximab plus outcomes of lenalidomide maintenance in relapsed CNS lymphoma. *Blood Adv* 2018;2:1595-1607.

**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY**  
**REFERENCES**

- <sup>103</sup>DeAngelis LM, Kreis W, Chan K, et al. Pharmacokinetics of ara-C and ara-U in plasma and CSF after high-dose administration of cytosine arabinoside. *Cancer Chemother Pharmacol* 1992;29:173-177.
- <sup>104</sup>Raizer JJ, Rademaker A, Evens AM, et al. Pemetrexed in the treatment of relapsed/refractory primary central nervous system lymphoma. *Cancer* 2012;118:3743-3748.
- <sup>105</sup>Tun HW, Johnston PB, DeAngelis LM, et al. Phase 1 study of pomalidomide and dexamethasone for relapsed/refractory primary CNS or vitreoretinal lymphoma. *Blood* 2018;132:2240-2248.
- <sup>106</sup>Soussain C, Hoang-Xuan K, Taillandier L, et al. Intensive chemotherapy followed by hematopoietic stem-cell rescue for refractory and recurrent primary CNS and intraocular lymphoma: Societe Francaise de Greffe de Moelle Osseuse-Therapie Cellulaire. *J Clin Oncol* 2008;26:2512-2518.
- <sup>107</sup>Kasenda B, Ihorst G, Schroers R, et al. High-dose chemotherapy with autologous haematopoietic stem cell support for relapsed or refractory primary CNS lymphoma: a prospective multicentre trial by the German Cooperative PCNSL study group. *Leukemia* 2017;31:2623-2629.
- <sup>108</sup>Kaley TJ, Wen P, Schiff D, et al. Phase II trial of sunitinib for recurrent and progressive atypical and anaplastic meningioma. *Neuro Oncol* 2015;17:116-121.
- <sup>109</sup>Lou E, Sumrall AL, Turner S, et al. Bevacizumab therapy for adults with recurrent/progressive meningioma: a retrospective series. *J Neurooncol* 2012;109:63-70.
- <sup>110</sup>Nayak L, Iwamoto FM, Rudnick JD, et al. Atypical and anaplastic meningiomas treated with bevacizumab. *J Neurooncol* 2012;109:187-193.
- <sup>111</sup>Shih KC, Chowdhary S, Rosenblatt P, et al. A phase II trial of bevacizumab and everolimus as treatment for patients with refractory, progressive intracranial meningioma. *J Neurooncol* 2016;129:281-288.
- <sup>112</sup>Chamberlain MC, Glantz MJ, Fadul CE. Recurrent meningioma: salvage therapy with long-acting somatostatin analogue. *Neurology* 2007;69:969-973.
- <sup>113</sup>Drilon A, Laetsch TW, Kummar W, et al. Efficacy of larotrectinib in TRK fusion-positive cancers in adults and children. *N Engl J Med* 2018;378:731-739.
- <sup>114</sup>Drilon AE, DuBois SG, Farago AF, et al. Activity of larotrectinib in TRK fusion cancer patients with brain metastases or primary central nervous system tumors. *J Clin Oncol* 2019;37(no. 15\_suppl): Abstract 2006.
- <sup>115</sup>Demetri GD, Paz-Ares LG, Farago AF, et al. Efficacy and safety of entrectinib in patients with NTRK fusion-positive (NTRK-fp) tumours: pooled analysis of STARTRK-2, STARTRK-1, and ALKA-372-001. *ESMO Congress* 2018.
- <sup>116</sup>Siena S, Doebele RC, Shaw AT, et al. Efficacy of entrectinib in patients with solid tumors and central nervous system metastases: integrated analysis from three clinical trials. *J Clin Oncol* 2019;37:3017.
- <sup>117</sup>Bachelot T, Romieu G, Campone M, et al. Lapatinib plus capecitabine in patients with previously untreated brain metastases from HER2-positive metastatic breast cancer (LANDSCAPE): a single-group phase 2 study. *Lancet Oncol* 2013;14:64-71.
- <sup>118</sup>Petrelli F, Ghidini M, Lonati V, et al. The efficacy of lapatinib and capecitabine in HER-2 positive breast cancer with brain metastases: A systematic review and pooled analysis. *Eur J Cancer* 2017;84:141-148.
- <sup>119</sup>Freedman RA, Gelman RS, Anders CK, et al. TBCRC 022: a phase II trial of neratinib and capecitabine for patients with human epidermal growth factor receptor 2-positive breast cancer and brain metastases. *J Clin Oncol* 2019;37:1081-1089.
- <sup>120</sup>Saura C, Oliveira M, Feng YH, et al. Neratinib + capecitabine versus lapatinib + capecitabine in patients with HER2+ metastatic breast cancer previously treated with ≥ 2 HER2-directed regimens: findings from the multinational, randomized, phase III NALA trial. *J Clin Oncol* 2019;37(no. 15\_suppl):Abstract 1002.
- <sup>121</sup>Awada A, Colomer R, Inoue K, et al. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEFERT-T randomized clinical trial. *JAMA Oncol* 2016;2:1557-1564.
- <sup>122</sup>Rivera E, Meyers C, Groves M, et al. Phase I study of capecitabine in combination with temozolomide in the treatment of patients with brain metastases from breast carcinoma. *Cancer* 2006;107:1348-1354.
- <sup>123</sup>Fabi A, Vidiri A, Ferretti G, et al. Dramatic regression of multiple brain metastases from breast cancer with capecitabine: another arrow at the bow? *Cancer Invest* 2006;24:466-468.
- <sup>124</sup>Siegelmann-Danieli N, Stein M, Bar-Ziv J. Complete response of brain metastases originating in breast cancer to capecitabine therapy. *Isr Med Assoc J* 2003;5:833-834.
- <sup>125</sup>Wang ML, Yung WK, Royce ME, et al. Capecitabine for 5-fluorouracil-resistant brain metastases from breast cancer. *Am J Clin Oncol* 2001;24:421-424.
- <sup>126</sup>Hikino H, Yamada T, Johbara K, et al. Potential role of chemo-radiation with oral capecitabine in a breast cancer patient with central nervous system relapse. *Breast* 2006;15:97-99.
- <sup>127</sup>Murthy RK, Loi S, Okines A, et al. Tucatinib, trastuzumab, and capecitabine for HER2-positive metastatic breast cancer. *N Engl J Med* 2020;382:597-609.
- <sup>128</sup>Cocconi G, Lottici R, Bisagni G, et al. Combination therapy with platinum and etoposide of brain metastases from breast carcinoma. *Cancer Invest* 1990;8:327-334.
- <sup>129</sup>Franciosi V, Cocconi G, Michiara M, et al. Front-line chemotherapy with cisplatin and etoposide for patients with brain metastases from breast carcinoma, nonsmall cell lung carcinoma, or malignant melanoma: a prospective study. *Cancer* 1999;85:1599-1605.
- <sup>130</sup>Vinolas N, Graus F, Mellado B, et al. Phase II trial of cisplatin and etoposide in brain metastases of solid tumors. *J Neurooncol* 1997;35:145-148.
- <sup>131</sup>Lassman AB, Abrey LE, Shah GD, et al. Systemic high-dose intravenous methotrexate for central nervous system metastases. *J Neurooncol* 2006;78:255-260.
- <sup>132</sup>Long GV, Trefzer U, Davies MA, et al. Dabrafenib in patients with Val600Glu or Val600Lys BRAF-mutant melanoma metastatic to the brain (BREAK-MB): a multicentre, open-label, phase 2 trial. *Lancet Oncol* 2012;13:1087-1095.
- <sup>133</sup>Falchook GS, Long GV, Kurzrock R, et al. Dabrafenib in patients with melanoma, untreated brain metastases, and other solid tumours: a phase 1 dose-escalation trial. *Lancet* 2012;379:1893-1901.
- <sup>134</sup>Azer MW, Menzies AM, Haydu LE, et al. Patterns of response and progression in patients with BRAF-mutant melanoma metastatic to the brain who were treated with dabrafenib. *Cancer* 2014;120:530-536.
- <sup>135</sup>Davies MA, Saiag P, Robert C, et al. Dabrafenib plus trametinib in patients with BRAFV600-mutant melanoma brain metastases (COMBI-MB): a multicentre, multicohort, open-label, phase 2 trial. *Lancet Oncol* 2017;18:863-873.

**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY**  
**REFERENCES**

- <sup>136</sup>McArthur GA, Maio M, Arance A, et al. Vemurafenib in metastatic melanoma patients with brain metastases: an open-label, single-arm, phase 2, multicentre study. *Ann Oncol* 2017;28:634-641.
- <sup>137</sup>Dummer R, Goldinger SM, Turttschi CP, et al. Vemurafenib in patients with BRAF(V600) mutation-positive melanoma with symptomatic brain metastases: final results of an open-label pilot study. *Eur J Cancer* 2014;50:611-621.
- <sup>138</sup>Tawbi HA, Forsyth PA, Algazi AP, et al. Combined nivolumab and ipilimumab in melanoma metastatic to the brain. *N Engl J Med* 2018;379:722-730.
- <sup>139</sup>Long GV, Atkinson V, Lo S, et al. Combination nivolumab and ipilimumab or nivolumab alone in melanoma brain metastases: a multicentre randomised phase 2 study. *Lancet Oncol* 2018;19:672-681.
- <sup>140</sup>Long GV, Atkinson V, Lo S, et al. long-term outcomes from the randomized phase II study of nivolumab or nivolumab + ipilimumab in patients with melanoma brain metastases: Anti-PD-1 Brain Collaboration (the ABC trial). *ESMO 2019 Congress*.
- <sup>141</sup>Margolin K, Ernstoff MS, Hamid O, et al. Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial. *Lancet Oncol* 2012;13:459-465.
- <sup>142</sup>Goldberg SB, Gettinger SN, Mahajan A, et al. Pembrolizumab for patients with melanoma or non-small-cell lung cancer and untreated brain metastases: early analysis of a non-randomised, open-label, phase 2 trial. *Lancet Oncol* 2016;17:976-983.
- <sup>143</sup>Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. *N Engl J Med* 2018;378:113-125.
- <sup>144</sup>Goss G, Tsai CM, Shepherd FA, et al. CNS response to osimertinib in patients with T790M-positive advanced NSCLC: pooled data from two phase II trials. *Ann Oncol* 2018;29:687-693.
- <sup>145</sup>Mok TS, Wu YL, Ahn MJ, et al. Osimertinib or platinum-pemetrexed in EGFR T790M-positive lung cancer. *N Engl J Med* 2017;376:629-640.
- <sup>146</sup>Grommes C, Oxnard GR, Kris MG, et al. "Pulsatile" high-dose weekly erlotinib for CNS metastases from EGFR mutant non-small cell lung cancer. *Neuro Oncol* 2011;13:1364-1369.
- <sup>147</sup>Katayama T, Shimizu J, Suda K, et al. Efficacy of erlotinib for brain and leptomeningeal metastases in patients with lung adenocarcinoma who showed initial good response to gefitinib. *J Thorac Oncol* 2009;4:1415-1419.
- <sup>148</sup>Arbour KC, Kris MG, Riely GJ, et al. Twice weekly pulse and daily continuous-dose erlotinib as initial treatment for patients with epidermal growth factor receptor-mutant lung cancers and brain metastases. *Cancer* 2018;124(1):105-109.
- <sup>149</sup>Hoffknecht P, Tufman A, Wehler T, et al. Efficacy of the irreversible ErbB family blocker afatinib in epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI)-pretreated non-small-cell lung cancer patients with brain metastases or leptomeningeal disease. *J Thorac Oncol* 2015;10:156-163.
- <sup>150</sup>Ceresoli GL, Cappuzzo F, Gregorc V, et al. Gefitinib in patients with brain metastases from non-small-cell lung cancer: a prospective trial. *Ann Oncol* 2004;15:1042-1047.
- <sup>151</sup>Wu C, Li YL, Wang ZM, et al. Gefitinib as palliative therapy for lung adenocarcinoma metastatic to the brain. *Lung Cancer* 2007;57:359-364.
- <sup>152</sup>Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib versus crizotinib in ALK-positive non-small-cell lung cancer. *N Engl J Med* 2018;379:2027-2039.
- <sup>153</sup>Kim DW, Tiseo M, Ahn MJ, et al. Brigatinib in patients with crizotinib-refractory anaplastic lymphoma kinase-positive non-small-cell lung cancer: a randomized, multicenter phase II trial. *J Clin Oncol* 2017;35:2490-2498.
- <sup>154</sup>Peters S, Camidge DR, Shaw AT, et al. Alectinib versus crizotinib in untreated ALK-positive non-small-cell lung cancers. *N Engl J Med* 2017;377:829-838.
- <sup>155</sup>Gandhi L, Ou SI, Shaw AT, et al. Efficacy of alectinib in central nervous system metastases in crizotinib-resistant ALK-positive non-small-cell lung cancer: comparison of RECIST 1.1 and RANO-HGG criteria. *Eur J Cancer* 2017;82:27-33.
- <sup>156</sup>Kim DW, Mehra R, Tan DSW, et al. Activity and safety of ceritinib in patients with ALK-rearranged non-small-cell lung cancer (ASCEND-1): updated results from the multicentre, open-label, phase 1 trial. *Lancet Oncol* 2016;17:452-463.
- <sup>157</sup>Costa DB, Shaw AT, Ou SH, et al. Clinical experience with crizotinib in patients with advanced ALK-rearranged non-small-cell lung cancer and brain metastases. *J Clin Oncol* 2015;33:1881-1888.
- <sup>158</sup>Kluger HM, Chiang V, Mahajan A, et al. Long-term survival of patients with melanoma with active brain metastases treated with pembrolizumab on a phase II trial. *J Clin Oncol* 2019;37:52-60.
- <sup>159</sup>Gauvain C, Vauleon E, Chouaid C, et al. Intracerebral efficacy and tolerance of nivolumab in non-small-cell lung cancer patients with brain metastases. *Lung Cancer* 2018;116:62-66.
- <sup>160</sup>Goldman JW, Crino L, Vokes EE, et al. P2.36: Nivolumab (nivo) in patients (pts) with advanced (adv) NSCLC and central nervous system (CNS) metastases (mets). *J Thorac Oncol* 2016;11:S238-S239.
- <sup>161</sup>Rizvi NA, Mazieres J, Planchard D, et al. Activity and safety of nivolumab, an anti-PD-1 immune checkpoint inhibitor, for patients with advanced, refractory squamous non-small-cell lung cancer (CheckMate 063): a phase 2, single-arm trial. *Lancet Oncol* 2015;16:257-265.
- <sup>162</sup>Bokstein F, Lossos A, Lossos IS, Siegal T. Central nervous system relapse of systemic non-Hodgkin's lymphoma: results of treatment based on high-dose methotrexate combination chemotherapy. *Leuk Lymphoma* 2002;43:587-593.
- <sup>163</sup>Chamberlain MC. Leptomeningeal metastasis. *Curr Opin Neurol* 2009;22:665-674.
- <sup>164</sup>Gutin PH, Weiss HD, Wiernik PH, Walker MD. Intrathecal N, N', N"-triethylenethiophosphoramidate [thio-TEPA (NSC 6396)] in the treatment of malignant meningeal disease: phase I-II study. *Cancer* 1976;38:1471-1475.
- <sup>165</sup>Groves MD, Glantz MJ, Chamberlain MC, et al. A multicenter phase II trial of intrathecal topotecan in patients with meningeal malignancies. *Neuro Oncol* 2008;10:208-215.
- <sup>166</sup>Chamberlain MC, Tsao-Wei DD, Groshen S. Phase II trial of intracerebrospinal fluid etoposide in the treatment of neoplastic meningitis. *Cancer* 2006;106:2021-2027.
- <sup>167</sup>Jaekle KA, Phuphanich S, Bent MJ, et al. Intrathecal treatment of neoplastic meningitis due to breast cancer with a slow-release formulation of cytarabine. *Br J Cancer* 2001;84:157-163.
- <sup>168</sup>Chamberlain MC, Johnston SK, Van Horn A, Glantz MJ. Recurrent lymphomatous meningitis treated with intra-CSF rituximab and liposomal ara-C. *J Neurooncol* 2009;91:271-277.
- <sup>169</sup>Glantz MJ, Jaekle KA, Chamberlain MC, et al. A randomized controlled trial comparing intrathecal sustained-release cytarabine (DepoCyt) to intrathecal methotrexate in patients with neoplastic meningitis from solid tumors. *Clin Cancer Res* 1999;5:3394-3402.

**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**





### PRINCIPLES OF BRAIN AND SPINAL CORD TUMOR SYSTEMIC THERAPY REFERENCES

- <sup>170</sup>Glantz MJ, LaFollette S, Jaeckle KA, et al. Randomized trial of a slow-release versus a standard formulation of cytarabine for the intrathecal treatment of lymphomatous meningitis. *J Clin Oncol* 1999;17:3110-3116.
- <sup>171</sup>Grossman SA, Finkelstein DM, Ruckdeschel JC, et al. Randomized prospective comparison of intraventricular methotrexate and thiopeta in patients with previously untreated neoplastic meningitis. Eastern Cooperative Oncology Group. *J Clin Oncol* 1993;11:561-569.
- <sup>172</sup>Hitchins RN, Bell DR, Woods RL, Levi JA. A prospective randomized trial of single-agent versus combination chemotherapy in meningeal carcinomatosis. *J Clin Oncol* 1987;5:1655-1662.
- <sup>173</sup>Ziegler JL, Bluming AZ. Intrathecal chemotherapy in Burkitt's lymphoma. *Br Med J* 1971;3:508-512.
- <sup>174</sup>Zagouri F, Sergentanis TN, Bartsch R, et al. Intrathecal administration of trastuzumab for the treatment of meningeal carcinomatosis in HER2-positive metastatic breast cancer: a systematic review and pooled analysis. *Breast Cancer Res Treat* 2013;139:13-22.
- <sup>175</sup>Glantz MJ, Cole BF, Recht L, et al. High-dose intravenous methotrexate for patients with nonleukemic leptomeningeal cancer: is intrathecal chemotherapy necessary? *J Clin Oncol* 1998;16(4):1561-1567.
- <sup>176</sup>Mrugala MM, Kim B, Sharma A, et al. Phase II study of systemic high-dose methotrexate and intrathecal liposomal cytarabine for treatment of leptomeningeal carcinomatosis from breast cancer. *Clin Breast Cancer* 2019;19(5):311-316.
- <sup>177</sup>Yang JCH, Cho BC, Kim DW, et al. Osimertinib for patients (pts) with leptomeningeal metastases (LM) from EGFR-mutant non-small cell lung cancer (NSCLC): updated results from the BLOOM study. *J Clin Oncol* 2017;35(no. 15\_suppl): Abstract 2020.
- <sup>178</sup>Nanjo S, Hata A, Okuda C, et al. Standard-dose osimertinib for refractory leptomeningeal metastases in T790M-positive EGFR-mutant non-small cell lung cancer. *Br J Cancer* 2017;118(1):32-37.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF BRAIN AND SPINE TUMOR MANAGEMENT****General**

Patients diagnosed with a tumor involving the brain, spinal cord, and related support structures should be referred to practitioners who are experienced in the diagnosis and management of these lesions.<sup>1</sup> The patient may (and should) be presented with options for care, which may include procedures or treatments best done by other specialists. The care options should then be discussed with the patient and their chosen supports in a manner that is understandable and culturally and educationally sensitive. It is strongly encouraged to discuss goals of care with the patient.

**Multidisciplinary Care**

- During the course of their treatment, most patients will be seen by multiple subspecialists. Close and regular communication among all providers across disciplines is essential. Brain tumor board or multidisciplinary clinic care models are strongly recommended. These models facilitate interactions among multiple subspecialists, ideally including allied health services (ie, physical, occupational and speech therapies, nursing, psychology, social work) for optimizing treatment plan recommendations.
- As treatment proceeds, it is important that the patient and family understand the role of each team member. One practitioner should be identified early on as the main point of contact for follow-up care questions. This individual can facilitate referral to the appropriate specialist.
- Offering patients the option of participation in a clinical trial is strongly encouraged. Practitioners should discuss any local, regional, and national options for which the patient may be eligible and the advantages and disadvantages of participation. Centers treating neuro-oncology patients are encouraged to participate in large collaborative trials in order to have local options to offer patients.
- Patients should be educated on the importance of informed consent and side effects when receiving chemotherapy.
- Throughout treatment the patient's quality of life should remain the highest priority and guide clinical decision-making. While responses on imaging are benchmarks of successful therapy, other indicators of success such as overall well-being, function in day-to-day activities, social and family interactions, nutrition, pain control, long-term consequences of treatment, and psychological issues must be considered.
- Patients should be informed of the possibility of pseudoprogression, its approximate incidence, and potential investigations that may be needed in the event that pseudoprogression is suspected. Close follow-up imaging, MR perfusion, MR spectroscopy, PET/CT imaging, and repeat surgery may be necessary if clinically indicated. Educate patients on the uncertainty of imaging as a whole, and the potential need for corollary testing to interpret scans.
- For patients with spine tumors, it is important to assemble a multidisciplinary team to integrate diagnosis, treatment, symptom management, and rehabilitation. Patients with spine tumors have complex physical, psychological, and social care needs.
- Optimal management requires a multidisciplinary team including the following expertise: neuro-oncology/medical and radiation oncology; surgery (ie, neurosurgery, orthopedic surgery, surgical oncology); radiology; interventional pain specialties; physical and rehabilitation medicine; physiatrists; experts in bowel and bladder care, back care, and ambulation support; physical therapy; occupational therapy; psychological and/or social services; and nutritional support.
- Practitioners should become familiar with palliative and hospice care resources that are available in their community in order to help educate patients and families that involvement of these services does not indicate a state of hopelessness, no further treatment, or abandonment. Palliative and pain management care should be integrated into management of neuro-oncology patients early in the course of their treatment.

[See NCCN Guidelines for Palliative Care](#) and Ferrell BR, Temel JS, Temin S, et al. Integration of Palliative Care Into Standard Oncology Care: American Society of Clinical Oncology Clinical Practice Guideline Update. J Clin Oncol 2017;35:96-112.

<sup>1</sup>Depending on local referral patterns and available expertise, this physician may be a neurosurgeon, neurologist, medical oncologist, or radiation oncologist.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**PRINCIPLES OF BRAIN AND SPINE TUMOR MANAGEMENT****Medical Management****1. Mass Effect, Brain Edema, Radiation Necrosis**

- Steroid therapy should be carefully monitored. If a patient is asymptomatic, steroids may be unnecessary. Careful questioning for subtle symptoms should be undertaken if edema is extensive on imaging. In general, the lowest dose of steroids should be used for the shortest time possible.<sup>2</sup> Downward titration of the dose should be attempted whenever possible. Patients with extensive mass effect should receive steroids for at least 24 hours before radiation therapy. Patients with a high risk of GI side effects (ie, perioperative patients, prior history of ulcers/GI bleed, receiving NSAIDs or anticoagulation) should receive H<sub>2</sub> blockers or proton pump inhibitors. Care should be taken to watch for development of steroid side effects.<sup>3</sup> Consider prophylactic treatment of pneumocystis jiroveci pneumonia for patients undergoing long-term steroid therapy. Consider short-course bevacizumab for management of symptoms driven by RT necrosis,<sup>4,5</sup> poorly controlled vasogenic edema, or mass effect in patients with brain metastases and primary brain tumors, particularly those with deep-seated unresectable tumors, as it may allow overall quality-of-life improvements by reducing steroid dose and improving functional status.<sup>6</sup>

**2. Seizures**

- Seizures are frequent in patients with primary or metastatic brain tumors. Despite this, studies have shown that the use of older, “traditional” anti-seizure medications, including phenytoin, phenobarbital, and valproic acid as prophylaxis against seizures in patients who have never had a seizure or who are undergoing neurosurgical procedures, is ineffective and is not recommended. Newer agents (ie, levetiracetam, topiramate, lamotrigine, pregabalin) have not yet been systematically studied.
- Seizure prophylaxis is not recommended as routine in asymptomatic patients but is reasonable to consider perioperatively.
- Many anti-seizure medications have significant effects on the cytochrome P450 system, and may have effects on the metabolism of numerous chemotherapeutic agents such as irinotecan, gefitinib, erlotinib, and temsirolimus among others. Where possible, such enzyme-inducing anti-seizure medications (EIAEDs) should be avoided (ie, phenytoin, phenobarbital, carbamazepine), and non-EIAEDs should be used instead (ie, levetiracetam, topiramate, valproic acid, lacosamide). Patients should be closely monitored for any adverse effects of the anti-seizure medications or chemotherapeutic agents.

**3. Endocrine Disorders**

- Endocrinopathies are common with brain tumor patients. This may be affected by concomitant steroid use as well as by radiotherapy, surgery, and certain medical therapies. Patients who present with a declining sense of well-being or quality of life should be evaluated not only for abnormalities related to their hypothalamic pituitary and adrenal axis, but also with regard to thyroid and gonad function. For patients who received prior RT, long-term monitoring of the hypothalamic pituitary and adrenal axis may be considered (eg, ACTH stimulation test, thyroid monitoring).

**4. Fatigue (Also see the [NCCN Guidelines for Cancer-Related Fatigue](#))**

- Fatigue is commonly experienced by brain tumor patients. This symptom can be severe, persistent, emotionally overwhelming, and not related to the degree or duration of physical activity. Screening should be initiated to identify any underlying medical sources of this symptom, after which patients can be taught energy conservation and organizational skills to help manage this effect. Supervised, moderate exercise may be of assistance for those in otherwise good general medical condition. More data are needed on the use of CNS stimulants and these agents are not routinely recommended.

<sup>2</sup>An exception to this rule is in the case of suspected CNS lymphoma. Steroids should be avoided where possible ([see PCNS-1](#)) prior to biopsy to allow best chance of diagnosis.

<sup>3</sup>Refractory hyperglycemia, skin changes, visual changes, fluid retention, and myopathy. If any of these changes occur, it is imperative to evaluate potential palliative treatments for them and also to evaluate the current dose of steroids to see if it can be reduced in an attempt to mitigate these side effects. Clinical monitoring for adrenal insufficiency is recommended when weaning steroids for patients who have been on long-term steroid therapy.

<sup>4</sup>Nayak L, Iwamoto FM, Rudnick JD, et al. Atypical and anaplastic meningiomas treated with bevacizumab. J Neurooncol 2012;109(1):187-193.

<sup>5</sup>Lou E, Sumrall AL, Turner S, et al. Bevacizumab therapy for adults with recurrent/progressive meningioma: a retrospective series. J Neurooncol 2012;109(1):63-70.

<sup>6</sup>Kaley T, Nolan C, Carver A, Omuro A. Bevacizumab for acute neurologic deterioration in patients with glioblastoma. CNS Oncol 2013;2:413-418.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**[Continued](#)**

**BRAIN-E**  
**2 OF 3**





### PRINCIPLES OF BRAIN AND SPINE TUMOR MANAGEMENT

#### Medical Management—continued

#### 5. Psychiatric Disorders (Also see the [NCCN Guidelines for Distress Management including NCCN Distress Thermometer \[DIS-A\]](#))

- Depression and/or anxiety is common in neuro-oncology patients. These symptoms are greater than simple sadness or anxiety associated with the diagnosis of a tumor. The vegetative symptoms associated with depression or severe anxiety may become very disabling for the patient and distressing for the family. These symptoms will respond to psychotropic medications as they do in non-tumor patients. If less severe, strong support from behavioral health allies and other qualified counselors is also extremely beneficial. Physicians, and other members of their health care teams, should be sensitive to these symptoms and inquire about them in follow-up visits in order to determine if the patient may be a candidate for psychological or psychiatric treatment. Communication between members of the patient's health care team regarding the patient's response to treatment is important. See Andersen BL, DeRubeis RJ, Berman BS, et al. Screening, assessment, and care of anxiety and depressive symptoms in adults with cancer: an American Society of Clinical Oncology guideline adaptation. J Clin Oncol 2014;32:1605-1619.
- Anti-seizure medications, anxiolytics, some chemotherapy agents, antiemetics, and other agents used directly in cancer therapy may affect mental status, alertness, and mood. Alterations in thought processes should trigger an investigation for any treatable causes, including endocrine disorders, infection, side effects of medication, or tumor progression.

#### 6. Venous Thromboembolism (VTE)

- See the [NCCN Guidelines for Cancer-Associated Venous Thromboembolic Disease](#).

#### Allied Services

- Physical therapy, occupational therapy, and speech therapy may be helpful for many patients with CNS tumors, either benign or malignant. Surgical intervention is not a prerequisite for referral, and these therapies should not be withheld from patients because of the uncertain course of certain malignant tumors. Many patients with aggressive, malignant primary brain tumors or CNS metastases can benefit from inpatient rehabilitation.
- Practitioners are encouraged to serve as a resource and to refer patients to social services, support groups, and cancer patient advocacy organizations. Institutional or community resources that can assist patients and families in dealing with financial, insurance, and legal issues are important.
- Practitioners should be familiar with their state laws concerning seizures and driving so that they can advise patients and families appropriately.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF BRAIN TUMOR PATHOLOGY

- Incorporation of relevant diagnostic markers, including histopathologic and molecular information, as per the WHO 2016 Classification of Tumors of the Central Nervous System should be considered standard practice for tumor classification.
- Molecular/genetic characterization complements standard histologic analysis, providing additional diagnostic and prognostic information that can greatly improve diagnostic accuracy, influence treatment selection, and possibly improve management decision-making.

#### Standard Histopathologic Examination and Classification

- Histologic subgrouping of CNS neoplasms provides valuable prognostic information, as is described in the WHO Classification of Tumors of the Central Nervous System.<sup>1</sup>
- Inter-observer discrepancies in histologic diagnosis and grading are a recognized issue, due to the inherently subjective nature of certain aspects of histopathologic interpretation (eg, astrocytic vs. oligodendroglial morphology). Also, surgical sampling does not always capture all the relevant diagnostic features in morphologically heterogeneous tumors.
- Even so, the traditional histologic classification of CNS neoplasms into primary neuroectodermal neoplasms (eg, glial, neuronal, embryonal), other primary CNS neoplasms (eg, lymphoma, germ cell, meningeal), metastatic neoplasms, and non-neoplastic conditions mimicking tumors remains fundamental to any pathologic assessment.

#### Molecular Characterization

- With the use of genetic and molecular testing, histologically similar CNS neoplasms can be differentiated more accurately in terms of prognosis and, in some instances, response to different therapies.<sup>2-6</sup>
- Molecular characterization of primary CNS tumors has substantially impacted clinical trial eligibility and risk stratification in the past 10 years, thereby evolving the standard of care towards an integrated tumor diagnosis in neuro-oncology.
- Molecular/genetic characterization does not replace standard histologic assessment, but serves as a complementary approach to provide additional diagnostic and prognostic information that often enhances treatment selection.
- There are no identified targeted agents with demonstrated efficacy in glioblastoma. However, the panel encourages molecular testing of tumor because if a driver mutation is detected, it may be reasonable to treat with a targeted therapy on a compassionate use basis and/or the patient may have more treatment options in the context of a clinical trial. Molecular testing also has a valuable role in improving diagnostic accuracy and prognostic stratification that may inform treatment selection.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)  
[References](#)

**BRAIN-F**  
**1 OF 8**



### PRINCIPLES OF BRAIN TUMOR PATHOLOGY MOLECULAR MARKERS

The following molecular markers are often used by neuropathologists to facilitate characterization of gliomas and/or by neuro-oncologists to guide treatment decisions:

#### **Isocitrate Dehydrogenase 1 and 2 (IDH1 and IDH2) Mutation**

- **Recommendation:** IDH mutation testing is required for the workup of glioma.
- **Description:** IDH1 and IDH2 are metabolic enzymes. Specific mutations in genes encoding these enzymes lead to the aberrant production of D-2-hydroxyglutarate, an oncometabolite that causes epigenetic modifications in affected cells.<sup>7</sup>
- **Detection:** The most common IDH1 mutation (R132H) is reliably screened by mutation-specific immunohistochemistry, which is recommended for all glioma patients. If the R132H immunostain result is negative, in the appropriate clinical context, sequencing of IDH1 and IDH2 is highly recommended to detect less common IDH1 and IDH2 mutations. Prior to age 55 years, sequencing of IDH1 and IDH2 is required if the R132H immunostain result is negative. Standard sequencing methods include Sanger sequencing, pyrosequencing, and next-generation sequencing, and should be performed on formalin-fixed, paraffin-embedded tissue.<sup>7</sup>
- **Diagnostic value:**
  - IDH mutations define WHO grade II and III astrocytomas and oligodendrogliomas, and the secondary grade IV glioblastomas into which astrocytomas often evolve. Their presence distinguishes lower-grade gliomas from primary glioblastomas, which are IDH-wild type.<sup>8,9</sup> Detection of these mutations in a specimen that is otherwise equivocal for tumor may also be regarded as evidence that a diffusely infiltrative glioma is present.<sup>7</sup>
  - True grade I non-infiltrative gliomas, such as pilocytic astrocytomas and gangliogliomas, do not contain IDH mutations. In such cases, detection of an IDH mutation indicates that the tumor is at least a grade II diffusely infiltrative glioma.<sup>7</sup>
- **Prognostic value:**
  - IDH mutations are commonly associated with MGMT promoter methylation.<sup>4</sup>
  - IDH1 or 2 mutations are associated with a relatively favorable prognosis and are important in stratification for clinical trials.<sup>10</sup>
  - In grade II or III infiltrative gliomas, wild-type IDH1 or 2 is associated with increased risk of aggressive disease.<sup>4</sup>
  - IDH1 or 2 mutations are associated with a survival benefit for patients treated with radiation or alkylating chemotherapy.<sup>11,12</sup>

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)  
[References](#)

**BRAIN-F**  
**2 OF 8**



### PRINCIPLES OF BRAIN TUMOR PATHOLOGY MOLECULAR MARKERS

The following molecular markers are often used by neuropathologists to facilitate characterization of gliomas and/or by neuro-oncologists to guide treatment decisions:

#### Codeletion of 1p and 19q

- **Recommendation:** 1p19q testing is an essential part of molecular diagnostics for oligodendroglioma.
- **Description:** This codeletion represents an unbalanced translocation (1;19)(q10;p10), leading to whole-arm deletion of 1p and 19q.<sup>13</sup>
- **Detection:** The codeletion of 1p and 19q is detectable by fluorescence in situ hybridization (FISH) or polymerase chain reaction (PCR). Additional methods, including array-based genomic copy number testing and next-generation sequencing, may also be employed.
- **Diagnostic value:** It is strongly associated with oligodendroglial histology and helps confirm the oligodendroglial character of tumors with equivocal or mixed histologic features.<sup>14</sup>
  - ▶ IDH-mutated gliomas that do NOT show loss of ATRX (for example, by IHC) should be strongly considered for 1p19q testing, even if not clearly oligodendroglial by histology. Conversely, IDH1 wild-type gliomas do not contain true whole-arm 1p/19q codeletion.<sup>15</sup> Therefore, 1p/19q testing is unnecessary if a glioma is not IDH-mutant, and a glioma should not be regarded as 1p/19q-codeleted without an accompanying IDH mutation, regardless of test results.
  - ▶ A tumor should only be diagnosed as an oligodendroglioma if it contains both an IDH mutation and 1p/19q codeletion. Furthermore, the term “oligoastrocytoma” should no longer be used, as such morphologically ambiguous tumors can reliably be resolved into astrocytomas and oligodendrogliomas with molecular testing.<sup>16</sup>
- **Prognostic value:** The codeletion confers a favorable prognosis and is predictive of response to alkylating chemotherapy and combination therapy with radiation and alkylating chemotherapy.<sup>17,18</sup>

#### MGMT Promoter Methylation

- **Recommendation:** MGMT promoter methylation is an essential part of molecular diagnostics for all high-grade gliomas (grade III and IV).
- **Description:** MGMT (O<sup>6</sup>-methylguanine-DNA methyltransferase) is a DNA repair enzyme that reverses the DNA damage caused by alkylating agents, resulting in tumor resistance to temozolomide and nitrosourea-based chemotherapy. Methylation of the MGMT promoter silences MGMT, making the tumor more sensitive to treatment with alkylating agents.<sup>19</sup>
- **Detection:** Methylation of the MGMT promoter is detectable by methylation-specific PCR,<sup>20</sup> pyrosequencing,<sup>21</sup> or array-based technologies.<sup>22</sup>
- **Prognostic value:**
  - ▶ MGMT promoter methylation is strongly associated with IDH mutations and genome-wide epigenetic changes (G-CIMP phenotype).<sup>4</sup>
  - ▶ MGMT promoter methylation confers a survival advantage in glioblastoma and is used for risk stratification in clinical trials.<sup>23</sup>
  - ▶ MGMT promoter methylation is particularly useful in treatment decisions for elderly patients with high-grade gliomas (grades III–IV).<sup>24,25</sup>
  - ▶ Patients with glioblastoma that are not MGMT promoter methylated derive less benefit from treatment with temozolomide compared to those whose tumors are methylated.<sup>23</sup>

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)  
[References](#)

**BRAIN-F**  
**3 OF 8**



### PRINCIPLES OF BRAIN TUMOR PATHOLOGY MOLECULAR MARKERS

The following molecular markers are often used by neuropathologists to facilitate characterization of gliomas and/or by neuro-oncologists to guide treatment decisions:

#### **ATRX Mutation**

- **Recommendation:** ATRX mutation testing is strongly recommended but not required for glioma.
- **Description:** ATRX encodes a chromatin regulator protein. Loss of function mutations enable alternative lengthening of telomeres (ALT).<sup>26</sup>
- **Detection:** ATRX mutations can be detected by IHC for wild-type ATRX (loss of wild-type expression) and/or sequencing.<sup>27</sup>
- **Diagnostic value:** ATRX mutations in glioma are strongly associated with IDH mutations, and are nearly always mutually exclusive with 1p/19q codeletion.<sup>27</sup> ATRX deficiency, coupled with IDH mutation, is typical of astrocytoma. A lack of ATRX immunostaining in glioblastoma should trigger IDH1/2 sequencing if IDH1 R132H immunostaining is negative, due to the frequent co-occurrence of ATRX and IDH mutations.<sup>5,27</sup>

#### **TERT Mutation**

- **Recommendation:** TERT mutation testing is recommended but not required for gliomas.
- **Description:** TERT encodes the catalytic active site of telomerase, the enzyme responsible for maintaining telomere length in dividing cells. TERT mutations found in gliomas are located in its noncoding promoter region, and cause increased expression of the TERT protein.<sup>28</sup>
- **Detection:** TERT mutations can be detected by sequencing of the promoter region.<sup>29</sup>
- **Diagnostic value:** TERT mutations are almost invariably present in 1p/19q codeleted oligodendroglioma, and are found in most glioblastomas. TERT mutation, in combination with IDH mutation and 1p/19q codeletion, is characteristic of oligodendroglioma. Absence of TERT mutation, coupled with IDH mutation, designates astrocytoma.
- **Prognostic value:** In the absence of an IDH mutation, TERT mutations in diffusely infiltrative gliomas are associated with reduced overall survival compared to gliomas lacking TERT mutations.<sup>4,30,31</sup> Combined TERT and IDH mutations in the absence of 1p/19q codeletion is an uncommon event, but such tumors have a prognosis as favorable as gliomas with all three molecular alterations.<sup>4,30</sup>

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF BRAIN TUMOR PATHOLOGY MOLECULAR MARKERS**

The following molecular markers are often used by neuropathologists to facilitate characterization of gliomas and/or by neuro-oncologists to guide treatment decisions:

**H3F3A Mutation**

- **Recommendation:** *H3F3A* and *HIST1H3B* mutation testing is recommended in the appropriate clinical context.
- **Description:** The most common histone mutation in brain tumors, *H3K27M*, is caused by a lysine-to-methionine substitution in the *H3F3A* gene and inhibits the trimethylation of *H3.3* histone. *G34* mutations are more common in cortical gliomas in children.<sup>32-34</sup>
- **Detection:** Although a *K27M* histone antibody is available,<sup>35</sup> it is not 100% specific and interpretation can be difficult for non-experts. Therefore, screening by *H3F3A* and *HIST1H3B* sequencing is a viable alternative and the preferred approach, especially since it will also detect mutations in *G34*.
- **Diagnostic value:** Histone mutations most commonly occur in pediatric midline gliomas (eg, diffuse intrinsic pontine gliomas [DIPG]), although midline gliomas in adults can also contain histone mutations.<sup>36</sup> Their presence can be considered solid evidence of an infiltrative glioma, which is often helpful in small biopsies of midline lesions that may not be fully diagnostic with light microscopy or do not fully resemble infiltrative gliomas.<sup>32,33,36</sup>
- **Prognostic value:** *K27M* gliomas typically do not have *MGMT* promoter methylation, and the mutation is an adverse prognostic marker in children and adults. The *G34* mutation does not appear to have any prognostic significance once the diagnosis of a glioblastoma has been established.<sup>33,36,37</sup>

**BRAF Mutation**

- **Recommendation:** *BRAF* fusion and/or mutation testing is recommended in the appropriate clinical context.
- **Description:** Activating mutations in *BRAF*, most commonly the V600E variant seen in other cancers (eg, melanoma), are present in 60%–80% of supratentorial grade II–III pleomorphic xanthoastrocytomas (PXA), 30% of dysembryoplastic neuroepithelial tumors, 20% of grade I gangliogliomas, and 5% of grade I pilocytic astrocytomas (PA). Diffusely infiltrative gliomas can also harbor a *BRAF* mutation, especially in children. *BRAF* V600E has even been found in nonneoplastic cortical dysplasia. In contrast, activating *BRAF* fusions occur predominately in PA of the posterior fossa, although some supratentorial PA also have this fusion.<sup>38-40</sup>
- **Detection:** *BRAF* V600E is best detected by sequencing, and *BRAF* fusions can be detected with RNA-Seq or other PCR-based breakpoint methods that capture the main 16–9, 15–9, and 16–11 breakpoints between *BRAF* and its main fusion partner, *KIAA1549*. FISH is too unreliable to detect *BRAF* fusions.<sup>38</sup>
- **Diagnostic value:** The presence of a *BRAF* fusion is reliable evidence that the tumor is a pilocytic astrocytoma, provided the histology is compatible. *BRAF* V600E is more complicated, as it can occur in a variety of tumors over all four WHO grades and requires integration with histology.<sup>38</sup>
- **Prognostic value:** Tumors with *BRAF* fusions tend to be indolent, with occasional recurrence but only rare progression to lethality. *BRAF* V600E tumors show a much greater range of outcomes and need to be considered in context with other mutations and clinicopathologic findings (eg, *CDKN2A/B* deletion). *BRAF* V600E tumors may respond to *BRAF* inhibitors such as vemurafenib, but comprehensive clinical trials are still ongoing.<sup>41-43</sup>

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)  
[References](#)**BRAIN-F**  
**5 OF 8**



**PRINCIPLES OF BRAIN TUMOR PATHOLOGY MOLECULAR MARKERS**

The following molecular markers are often used by neuropathologists to facilitate characterization of gliomas and/or by neuro-oncologists to guide treatment decisions:

**RELA Fusion**

- **Recommendation:** RELA fusion testing is recommended in the appropriate clinical context.
- **Description:** Ependymomas arising in the supratentorium often contain activating fusions of RELA. Increased RELA activity leads to increased NF-kappa-B signaling and more aggressive behavior. This event is more common in children than in adults, and occurs only in the supratentorium, not the posterior fossa or spine.<sup>44,45</sup>
- **Detection:** The most common RELA fusion partner is C11orf95. This can be detected with RNA-Seq or a break-apart FISH probe set.<sup>46</sup>
- **Diagnostic value:** Detection of RELA fusion is not required for the diagnosis of ependymoma, as this entity is still diagnosed by light microscopy.
- **Prognostic value:** RELA fusion-positive ependymomas are now a distinct entity in the WHO classification of CNS tumors, as this subset of ependymomas tends to be far more aggressive than other supratentorial ependymomas.<sup>1,44,45,47</sup>

**Medulloblastoma Molecular Subtyping**

- **Recommendation:** Medulloblastoma testing should be referred to academic tertiary centers with expertise in this area.
- **Description:** Medulloblastomas are WHO grade IV tumors that predominantly arise from the cerebellum in pediatric patients, but can also occur in adults. The WHO committee on CNS tumors now recommends subclassification of these tumors into four distinct groups: i) WNT-activated; ii) Sonic hedgehog (SHH)-activated and *TP53*-mutant; iii) SHH-activated and *TP53*-wild type; and iv) non-WNT/non-SHH.<sup>1,48</sup>
- **Detection:** Virtually all WNT-driven medulloblastomas will contain mutations in either CTNNB1 or, less commonly, APC (the latter mutation may be germline if the patient has Turcot syndrome). Unlike in children, 50% of adult medulloblastomas with loss of 6q and positive nuclear catenin had no CTNNB1 mutations, pointing towards the possibility of alternative mechanisms of WNT pathway activation in adult medulloblastoma.<sup>49</sup> Adult and pediatric medulloblastomas are genetically distinct and require different algorithms for molecular risk stratification. WNT-driven tumors will also usually contain monosomy 6. 6q loss is not confined to WNT in adults; it is also described in SHH and Group 4. Monosomy 6 is a specific marker for pediatric WNT, but not for adult WNT.<sup>50</sup> Nuclear immunoreactivity for beta-catenin is a very useful way to identify WNT medulloblastomas, in conjunction with CTNNB1 sequencing and chromosome 6 FISH. Differentiating between WNT-activated, SHH-activated, and non-WNT/non-SHH tumors is best classified by expression arrays, DNA methylation arrays, or an immunohistochemistry panel composed of beta-catenin, GAB1, and YAP1. Because there are a variety of hotspots in *TP53*, gene sequencing is recommended in SHH-activated medulloblastomas.<sup>51-54</sup>
- **Diagnostic value:** None of the molecular markers associated with each medulloblastoma subtype is specific to medulloblastomas; the diagnosis of medulloblastoma is still made on the basis of light microscopy.
- **Prognostic value:** The most important aspect of medulloblastoma molecular diagnostics is that the WNT-activated subset has a markedly better prognosis relative to the other three subtypes, regardless of age at diagnosis. Among SHH-activated medulloblastomas, detection of *TP53* mutations is associated with more aggressive behavior, often in the setting of germline *TP53* mutations, wildtype SHH-activated medulloblastomas have a variable course, and are uncommon in adults.<sup>55-57</sup> Non-WNT/non-SHH medulloblastomas also show a variable course.<sup>1,48,55</sup> WNT tumors have worse prognosis in adults compared to children based on retrospective data.<sup>50</sup> 6q loss and positive nuclear catenin have no clear prognostic role in adult medulloblastomas.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**References**



**PRINCIPLES OF BRAIN TUMOR PATHOLOGY  
REFERENCES**

- <sup>1</sup>Louis DN, Perry A, Reifenberger G, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol* 2016;131:803-820.
- <sup>2</sup>Weller M, Pfister SM, Wick W, et al. Molecular neuro-oncology in clinical practice: a new horizon. *Lancet Oncol* 2013;14:e370-379.
- <sup>3</sup>Hegi ME, Stupp R. Withholding temozolomide in glioblastoma patients with unmethylated MGMT promoter--still a dilemma? *Neuro Oncol* 2015;17:1425-1427.
- <sup>4</sup>Eckel-Passow JE, Lachance DH, Molinaro AM, et al. Glioma Groups Based on 1p/19q, IDH, and TERT Promoter Mutations in Tumors. *N Engl J Med* 2015;372:2499-2508.
- <sup>5</sup>Cancer Genome Atlas Research N, Brat DJ, Verhaak RG, et al. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med* 2015;372:2481-2498.
- <sup>6</sup>Dubbink HJ, Atmodimedjo PN, Kros JM, et al. Molecular classification of anaplastic oligodendroglioma using next-generation sequencing: a report of the prospective randomized EORTC Brain Tumor Group 26951 phase III trial. *Neuro Oncol* 2016;18:388-400.
- <sup>7</sup>Horbinski C. What do we know about IDH1/2 mutations so far, and how do we use it? *Acta Neuropathol* 2013;125:621-636.
- <sup>8</sup>Hartmann C, Meyer J, Balss J, et al. Type and frequency of IDH1 and IDH2 mutations are related to astrocytic and oligodendroglial differentiation and age: a study of 1,010 diffuse gliomas. *Acta Neuropathol* 2009;118:469-474.
- <sup>9</sup>Yan H, Parsons DW, Jin G, et al. IDH1 and IDH2 mutations in gliomas. *N Engl J Med* 2009;360:765-773.
- <sup>10</sup>Sanson M, Marie Y, Paris S, et al. Isocitrate dehydrogenase 1 codon 132 mutation is an important prognostic biomarker in gliomas. *J Clin Oncol* 2009;27:4150-4154.
- <sup>11</sup>Hartmann C, Hentschel B, Wick W, et al. Patients with IDH1 wild type anaplastic astrocytomas exhibit worse prognosis than IDH1-mutated glioblastomas, and IDH1 mutation status accounts for the unfavorable prognostic effect of higher age: implications for classification of gliomas. *Acta Neuropathol* 2010;120:707-718.
- <sup>12</sup>Houillier C, Wang X, Kaloshi G, et al. IDH1 or IDH2 mutations predict longer survival and response to temozolomide in low-grade gliomas. *Neurology* 2010;75:1560-1566.
- <sup>13</sup>Jenkins RB, Blair H, Ballman KV, et al. A t(1;19)(q10;p10) mediates the combined deletions of 1p and 19q and predicts a better prognosis of patients with oligodendroglioma. *Cancer Res* 2006;66:9852-9861.
- <sup>14</sup>Burger PC, Minn AY, Smith JS, et al. Losses of chromosomal arms 1p and 19q in the diagnosis of oligodendroglioma. A study of paraffin-embedded sections. *Mod Pathol* 2001;14:842-853.
- <sup>15</sup>Labussiere M, Idbaih A, Wang XW, et al. All the 1p19q codeleted gliomas are mutated on IDH1 or IDH2. *Neurology* 2010;74:1886-1890.
- <sup>16</sup>Sahm F, Reuss D, Koelsche C, et al. Farewell to oligoastrocytoma: in situ molecular genetics favor classification as either oligodendroglioma or astrocytoma. *Acta Neuropathol* 2014;128:551-559.
- <sup>17</sup>van den Bent MJ, Brandes AA, Taphoorn MJ, et al. Adjuvant procarbazine, lomustine, and vincristine chemotherapy in newly diagnosed anaplastic oligodendroglioma: long-term follow-up of EORTC brain tumor group study 26951. *J Clin Oncol* 2013;31:344-350.
- <sup>18</sup>Cairncross G, Wang M, Shaw E, et al. Phase III Trial of Chemoradiotherapy for Anaplastic Oligodendroglioma: Long-Term Results of RTOG 9402. *J Clin Oncol* 2013;31:337-343.
- <sup>19</sup>Esteller M, Garcia-Foncillas J, Andion E, et al. Inactivation of the DNA-repair gene MGMT and the clinical response of gliomas to alkylating agents. *N Engl J Med* 2000;343:1350-1354.
- <sup>20</sup>Vlassenbroeck I, Califice S, Diserens AC, et al. Validation of real-time methylation-specific PCR to determine O6-methylguanine-DNA methyltransferase gene promoter methylation in glioma. *J Mol Diagn* 2008;10:332-337.
- <sup>21</sup>Xie H, Tubbs R, Yang B. Detection of MGMT promoter methylation in glioblastoma using pyrosequencing. *Int J Clin Exp Pathol* 2015;8:636-642.
- <sup>22</sup>Bady P, Sciuscio D, Diserens AC, et al. MGMT methylation analysis of glioblastoma on the Infinium methylation BeadChip identifies two distinct CpG regions associated with gene silencing and outcome, yielding a prediction model for comparisons across datasets, tumor grades, and CIMP-status. *Acta Neuropathol* 2012;124:547-560.
- <sup>23</sup>Hegi ME, Diserens AC, Gorlia T, et al. MGMT gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med* 2005;352:997-1003.
- <sup>24</sup>Malmstrom A, Gronberg BH, Marosi C, et al. Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised, phase 3 trial. *Lancet Oncol* 2012;13:916-926.
- <sup>25</sup>Wick W, Platten M, Meisner C, et al. Temozolomide chemotherapy alone versus radiotherapy alone for malignant astrocytoma in the elderly: the NOA-08 randomised, phase 3 trial. *Lancet Oncol* 2012;13:707-715.
- <sup>26</sup>Koschmann C, Calinescu AA, Nunez FJ, et al. ATRX loss promotes tumor growth and impairs nonhomologous end joining DNA repair in glioma. *Sci Transl Med* 2016;8:328ra28.
- <sup>27</sup>Reuss DE, Sahm F, Schrimpf D, et al. ATRX and IDH1-R132H immunohistochemistry with subsequent copy number analysis and IDH sequencing as a basis for an "integrated" diagnostic approach for adult astrocytoma, oligodendroglioma and glioblastoma. *Acta Neuropathol* 2015;129:133-146.
- <sup>28</sup>Arita H, Narita Y, Fukushima S, et al. Upregulating mutations in the TERT promoter commonly occur in adult malignant gliomas and are strongly associated with total 1p19q loss. *Acta Neuropathol* 2013;126:267-276.
- <sup>29</sup>Nikiforova MN, Wald AI, Melan MA, et al. Targeted next-generation sequencing panel (GlioSeq) provides comprehensive genetic profiling of central nervous system tumors. *Neuro Oncol* 2016;18:379-387.

**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**[Continued](#)

**PRINCIPLES OF BRAIN TUMOR PATHOLOGY**  
**REFERENCES**

- <sup>30</sup>Killela PJ, Reitman ZJ, Jiao Y, et al. TERT promoter mutations occur frequently in gliomas and a subset of tumors derived from cells with low rates of self-renewal. *Proc Natl Acad Sci U S A* 2013;110:6021-6026.
- <sup>31</sup>Arita H, Yamasaki K, Matsushita Y, et al. A combination of TERT promoter mutation and MGMT methylation status predicts clinically relevant subgroups of newly diagnosed glioblastomas. *Acta Neuropathol Commun* 2016;4:79.
- <sup>32</sup>Schwartzentruber J, Korshunov A, Liu XY, et al. Driver mutations in histone H3.3 and chromatin remodelling genes in paediatric glioblastoma. *Nature* 2012;482:226-231.
- <sup>33</sup>Sturm D, Witt H, Hovestadt V, et al. Hotspot mutations in H3F3A and IDH1 define distinct epigenetic and biological subgroups of glioblastoma. *Cancer Cell* 2012;22:425-437.
- <sup>34</sup>Wu G, Broniscer A, McEachron TA, et al. Somatic histone H3 alterations in pediatric diffuse intrinsic pontine gliomas and non-brainstem glioblastomas. *Nat Genet* 2012;44:251-253.
- <sup>35</sup>Bechet D, Gielen GG, Korshunov A, et al. Specific detection of methionine 27 mutation in histone 3 variants (H3K27M) in fixed tissue from high-grade astrocytomas. *Acta Neuropathol* 2014;128:733-741.
- <sup>36</sup>Meyronet D, Esteban-Mader M, Bonnet C, et al. Characteristics of H3 K27M-mutant gliomas in adults. *Neuro Oncol* 2017;19:1127-1134.
- <sup>37</sup>Khuong-Quang DA, Buczkowicz P, Rakopoulos P, et al. K27M mutation in histone H3.3 defines clinically and biologically distinct subgroups of pediatric diffuse intrinsic pontine gliomas. *Acta Neuropathol* 2012;124:439-447.
- <sup>38</sup>Horbinski C. To BRAF or not to BRAF: is that even a question anymore? *J Neuropathol Exp Neurol* 2013;72:2-7.
- <sup>39</sup>Qaddoumi I, Orisme W, Wen J, et al. Genetic alterations in uncommon low-grade neuroepithelial tumors: BRAF, FGFR1, and MYB mutations occur at high frequency and align with morphology. *Acta Neuropathol* 2016;131:833-845.
- <sup>40</sup>Chappe C, Padovani L, Scavarda D, et al. Dysembryoplastic neuroepithelial tumors share with pleomorphic xanthoastrocytomas and gangliogliomas BRAF(V600E) mutation and expression. *Brain Pathol* 2013;23:574-583.
- <sup>41</sup>Hawkins C, Walker E, Mohamed N, et al. BRAF-KIAA1549 fusion predicts better clinical outcome in pediatric low-grade astrocytoma. *Clin Cancer Res* 2011;17:4790-4798.
- <sup>42</sup>Horbinski C, Nikiforova MN, Hagenkord JM, et al. Interplay among BRAF, p16, p53, and MIB1 in pediatric low-grade gliomas. *Neuro Oncol* 2012;14:777-789.
- <sup>43</sup>Horbinski C, Hamilton RL, Nikiforov Y, Pollack IF. Association of molecular alterations, including BRAF, with biology and outcome in pilocytic astrocytomas. *Acta Neuropathol* 2010;119:641-649.
- <sup>44</sup>Parker M, Mohankumar KM, Punchihewa C, et al. C11orf95-RELA fusions drive oncogenic NF-kappaB signalling in ependymoma. *Nature* 2014;506:451-455.
- <sup>45</sup>Pajtl KW, Witt H, Sill M, et al. Molecular Classification of Ependymal Tumors across All CNS Compartments, Histopathological Grades, and Age Groups. *Cancer Cell* 2015;27:728-743.
- <sup>46</sup>Olsen TK, Panagopoulos I, Gorunova L, et al. Novel fusion genes and chimeric transcripts in ependymal tumors. *Genes Chromosomes Cancer* 2016;55:944-953.
- <sup>47</sup>Hubner JM, Kool M, Pfister SM, et al. Epidemiology, molecular classification and WHO grading of ependymoma. *J Neurosurg Sci* 2018;62(1):46-50.
- <sup>48</sup>Kool M, Korshunov A, Remke M, et al. Molecular subgroups of medulloblastoma: an international meta-analysis of transcriptome, genetic aberrations, and clinical data of WNT, SHH, Group 3, and Group 4 medulloblastomas. *Acta Neuropathol* 2012;123:473-484.
- <sup>49</sup>Korshunov A, Remke M, Werft W, et al. Adult and pediatric medulloblastomas are genetically distinct and require different algorithms for molecular risk stratification. *J Clin Oncol* 2010;28:3054-3060.
- <sup>50</sup>Remke M, Hielscher T, Northcott PA, et al. Adult medulloblastoma comprises three major molecular variants. *J Clin Oncol* 2011;29(19):2717-2723.
- <sup>51</sup>Northcott PA, Buchhalter I, Morrissy AS, et al. The whole-genome landscape of medulloblastoma subtypes. *Nature* 2017;547:311-317.
- <sup>52</sup>Ellison DW, Kocak M, Dalton J, et al. Definition of disease-risk stratification groups in childhood medulloblastoma using combined clinical, pathologic, and molecular variables. *J Clin Oncol* 2011;29:1400-1407.
- <sup>53</sup>Schwalbe EC, Williamson D, Lindsey JC, et al. DNA methylation profiling of medulloblastoma allows robust subclassification and improved outcome prediction using formalin-fixed biopsies. *Acta Neuropathol* 2013;125:359-371.
- <sup>54</sup>Taylor MD, Northcott PA, Korshunov A, et al. Molecular subgroups of medulloblastoma: the current consensus. *Acta Neuropathol* 2012;123:465-472.
- <sup>55</sup>Zhukova N, Ramaswamy V, Remke M, et al. Subgroup-specific prognostic implications of TP53 mutation in medulloblastoma. *J Clin Oncol* 2013;31:2927-2935.
- <sup>56</sup>Kool M, Jones DT, Jager N, et al. Genome sequencing of SHH medulloblastoma predicts genotype-related response to smoothened inhibition. *Cancer Cell* 2014;25:393-405.
- <sup>57</sup>Waszak SM, Northcott PA, Buchhalter I, et al. Spectrum and prevalence of genetic predisposition in medulloblastoma: a retrospective genetic study and prospective validation in a clinical trial cohort. *Lancet Oncol* 2018;19:785-798.

**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



### NCCN Categories of Evidence and Consensus

<b>Category 1</b>	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
<b>Category 2A</b>	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
<b>Category 2B</b>	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
<b>Category 3</b>	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

### NCCN Categories of Preference

<b>Preferred intervention</b>	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
<b>Other recommended intervention</b>	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
<b>Useful in certain circumstances</b>	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



# NCCN Guidelines Version 3.2020 Central Nervous System Cancers

## Discussion

This discussion corresponds to the NCCN Guidelines for Central Nervous System Cancers. Last updated on September 11, 2020.

## Table of Contents

<b>Overview .....</b>	<b>MS-2</b>
<b>Literature Search Criteria and Guidelines Update Methodology .....</b>	<b>MS-2</b>
<b>Tumor Types.....</b>	<b>MS-2</b>
<b>Principles of Management .....</b>	<b>MS-2</b>
<b>Treatment Principles .....</b>	<b>MS-3</b>
<b>Gliomas.....</b>	<b>MS-3</b>
Molecular Profiling for Gliomas.....	MS-4
Low-Grade Gliomas .....	MS-6
Anaplastic Gliomas and Glioblastomas.....	MS-11
<b>Intracranial and Spinal Ependymomas.....</b>	<b>MS-19</b>
Treatment Overview.....	MS-20
NCCN Recommendations .....	MS-21
<b>Adult Medulloblastoma .....</b>	<b>MS-22</b>
Treatment Overview.....	MS-23
NCCN Recommendations .....	MS-23
<b>Primary CNS Lymphomas .....</b>	<b>MS-24</b>
Treatment Overview.....	MS-25
NCCN Recommendations .....	MS-27

<b>Primary Spinal Cord Tumors.....</b>	<b>MS-30</b>
Treatment Overview .....	MS-30
NCCN Recommendations .....	MS-31
<b>Meningiomas .....</b>	<b>MS-31</b>
Imaging.....	MS-32
Treatment Overview .....	MS-32
NCCN Recommendations .....	MS-34
<b>Brain Metastases .....</b>	<b>MS-34</b>
Treatment Overview .....	MS-35
NCCN Recommendations .....	MS-42
<b>Leptomeningeal Metastases .....</b>	<b>MS-43</b>
Treatment Overview .....	MS-43
NCCN Recommendations .....	MS-44
<b>Metastatic Spinal Tumors.....</b>	<b>MS-46</b>
Treatment Overview .....	MS-46
NCCN Recommendations .....	MS-49
<b>References.....</b>	<b>MS-51</b>



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### Overview

In the year 2020, an estimated 23,890 people in the United States will be diagnosed with a malignant primary central nervous system (CNS) tumor, and these tumors will be responsible for approximately 18,020 deaths.<sup>1</sup> The incidence of primary brain tumors has been increasing over recent decades,<sup>2</sup> especially in older adults.<sup>3,4</sup> However, this growth in incidence rates may be explained by increased use of CT and MRI and changes in WHO classification of CNS tumors.<sup>2,5,6</sup>

### Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines for Central Nervous System Cancers, an electronic search of the PubMed database was performed to obtain key literature in the field of neuro-oncology, using the following search terms: {(brain OR spine OR spinal OR supratentorial OR cranial OR intracranial OR leptomeningeal) AND (cancer OR carcinoma OR tumor OR metastases OR lesion)] OR glioma OR astrocytoma OR oligodendroglioma OR glioblastoma OR ependymoma OR medulloblastoma OR (primary central nervous system lymphoma) OR meningioma}. The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.<sup>7</sup>

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Practice Guideline; Guidelines; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles and articles from additional sources deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications

ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available on the NCCN website ([www.NCCN.org](http://www.NCCN.org)).

### Tumor Types

The NCCN Guidelines for CNS Cancers focus on management of the following adult CNS cancers: pilocytic and infiltrative supratentorial astrocytomas, oligodendrogliomas, anaplastic gliomas and glioblastoma, ependymomas, medulloblastoma, brain metastases, leptomeningeal metastases, non-AIDS-related primary CNS lymphomas (PCNSLs), metastatic spinal tumors, meningiomas, and primary spinal cord tumors. These guidelines are updated annually to include new information or treatment philosophies as they become available. However, because this field continually evolves, practitioners should use all of the available information to determine the best clinical options for their patients.

### Principles of Management

Primary brain tumors are a heterogeneous group of neoplasms with varied outcomes and management strategies. Primary malignant brain tumors range from pilocytic astrocytomas, which are very uncommon, noninvasive, and surgically curable, to glioblastoma, the most common malignant brain tumor in adults, which is highly invasive and virtually incurable. Likewise, patients with metastatic brain disease may have rapidly progressive systemic disease or no systemic cancer at all. These patients may have one or dozens of brain metastases, and they may have a malignancy that is highly responsive or, alternatively, highly resistant to radiation therapy (RT) or chemotherapy. Because of this marked heterogeneity, the prognostic features and treatment options for primary and metastatic brain tumors must be carefully reviewed on an individual





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

basis and sensitively communicated to each patient. In addition, these CNS tumors are associated with a range of symptoms such as seizures, fatigue, psychiatric disorders, impaired mobility, neuro-cognitive dysfunction, difficulty speaking, and short-term memory problems, as well as complications such as intracerebral edema, endocrinopathies, and venous thromboembolism that can seriously impact patients' quality of life.

The involvement of an interdisciplinary team, including neurosurgeons, RT therapists, oncologists, neurologists, and neuroradiologists, is a key factor in the appropriate management of these patients. For any subtype of malignant brain tumors, the NCCN Panel strongly recommends brain tumor board or multidisciplinary review of each patient's case once the pathology is available. Further discussion of multidisciplinary care and allied services, as well as guidelines on medical management of various disease complications, can be found in the section, *Principles of Brain and Spine Tumor Management* in the algorithm.

### Treatment Principles

Several important principles guide surgical management and treatment with RT and systemic therapy for adults with brain tumors. Regardless of tumor histology, neurosurgeons generally provide the best outcome for their patients if they remove as much tumor as safely possible (ideally achieving a gross total resection [GTR]) and thereby provide sufficient representative tumor tissue to ensure an accurate diagnosis. Decisions regarding aggressiveness of surgery for primary brain tumors are complex and depend on the: 1) age and performance status (PS) of the patient; 2) proximity to “eloquent” areas of the brain; 3) feasibility of decreasing the mass effect with aggressive surgery; 4) resectability of the tumor (including the number and location of lesions); and 5) time since last surgery in patients with recurrent disease.<sup>8</sup> Further discussion can be found in the *Principles of Brain Tumor Surgery* in the algorithm. It is

recommended to consult neurosurgeons with extensive experience in the management of intracranial and spine neoplasms.

The surgical options include stereotactic biopsy, open biopsy, subtotal resection (STR), or GTR. The pathologic diagnosis is critical and may be difficult to accurately determine without sufficient tumor tissue. Review of the tumor tissue by an experienced neuropathologist is highly recommended. The *Principles of Brain Tumor Pathology* describe guiding principles for diagnosis of CNS tumor pathology, given the 2016 addition of molecular parameters to the WHO classification of CNS tumors.<sup>9</sup>

Radiation oncologists use several different treatment modalities in patients with primary brain tumors, including fractionated stereotactic RT and stereotactic radiosurgery (SRS). Standard fractionated external beam RT (EBRT) is the most common approach. Hypofractionated radiation is an appropriate option for select patients (ie, older adults and patients with a poor PS). RT for patients with primary brain tumors is administered within a limited field (covering tumor or surgical cavity and a small margin of adjacent brain tissue), while whole-brain RT (WBRT) and SRS are used primarily for treatment of brain metastases. The dose of RT administered varies depending on the pathology as seen in *Principles of Radiation Therapy for Brain and Spinal Cord*.

The information contained in the algorithms and principles of management sections are designed to help clinicians navigate through the complex management of patients with CNS tumors. Standard systemic therapy options for each tumor subtype are listed in the *Principles of Brain and Spinal Cord Tumor Systemic Therapy*; however, enrollment in a clinical trial is the preferred treatment for eligible patients.

### Gliomas

The NCCN Guidelines for CNS Cancers include recommendations for management of the following gliomas:<sup>9</sup>





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

- Grade I: pilocytic astrocytoma, pleomorphic xanthoastrocytoma, ganglioglioma, and subependymal giant cell astrocytoma
- Grade II: diffuse astrocytomas and oligodendrogliomas
- Grade III: anaplastic astrocytoma and oligodendroglioma
- Grade IV: glioblastoma

### Molecular Profiling for Gliomas

Integrated histopathologic and molecular characterization of gliomas should be standard practice. Molecular/genetic characterization complements standard histologic analysis, providing additional diagnostic and prognostic information that improves diagnostic accuracy and aids in treatment selection.

#### **Updated Classification of Gliomas Based on Histology and Molecular Features**

In 2016, the WHO classification for grade II–III gliomas was revised as follows: 1) oligodendrogliomas are now defined as tumors that have 1p19q codeletion and *IDH* mutation (unless molecular data are not available and cannot be obtained, in which case designation can be based on histology with appropriate caveats); 2) anaplastic gliomas were further subdivided according to *IDH* mutation status; 3) oligoastrocytoma is no longer a valid designation unless molecular data (1p19q codeletion and *IDH* mutation status) are not available and cannot be obtained.<sup>9</sup> Such tumors should be described as “oligoastrocytoma, not otherwise specified (NOS)” to indicate that the characterization of the tumor is incomplete. Very rare cases of concurrent, spatially distinct oligodendroglioma (1p19q codeleted) and astrocytoma (1p19q intact) components in the same tumor may also be labeled oligoastrocytoma.<sup>9</sup> It is important to note that correlations between the molecularly defined 2016 WHO categories and the histology-based 2007 WHO categories are limited and vary across studies.<sup>10–13</sup> Thus, the

change from 2007 WHO to 2016 WHO reclassified a significant proportion of gliomas.

Multiple independent studies on gliomas have conducted genome-wide analyses evaluating an array of molecular features (eg, DNA copy number, DNA methylation, protein expression) in large populations of patients with grade II–IV tumors.<sup>12,14,15</sup> Unsupervised clustering analyses, an unbiased method for identifying molecularly similar tumors, have been used to identify subgroups of gliomas with distinct molecular profiles.<sup>12,14,15</sup> Remarkably, further analysis has shown that these molecular subgroups could be distinguished based on only a handful of molecular features, including mutation of *IDH1/2* and 1p19q codeletion, biomarkers independently verified by many studies as hallmarks for distinguishing molecular subgroups in grade II–III gliomas.<sup>10–13,15–21</sup> Using these markers alone, the majority of grade II–III tumors can be divided into three molecular subtypes: 1) mutation of either *IDH1* or *IDH2* (*IDH*-mut) with 1p19q codeletion (1p19q codelet); 2) *IDH*-mut with no 1p19q codeletion or with isolated deletion of 1p or 19q; and 3) no mutation of *IDH1* or *IDH2* (*IDH* wild type; *IDH*-wt).<sup>12</sup> Multiple studies have shown that the 1p19q codeletion is strongly associated with *IDH* mutations, such that true whole-arm 1p19q codeletion in *IDH*-wt tumors is extremely rare.<sup>10,11,18,22,23</sup> In a tumor that is equivocal, the presence of an *IDH* mutation indicates at least a grade II diffusely infiltrative glioma.<sup>24</sup> Grade I non-infiltrative gliomas do not have *IDH* mutations.<sup>24</sup>

Other mutations commonly detected in gliomas can have diagnostic and prognostic value, such as those involving the histone chaperone protein, *ATRX*, which is most often found in grade II–III gliomas and secondary glioblastomas.<sup>25,26</sup> *ATRX* mutation is robustly associated with *IDH* mutations, and this combination is strongly suggestive of astrocytoma.<sup>27</sup> In contrast, *ATRX* mutation is nearly always mutually exclusive with 1p19q codeletion. Therefore, a glioma that has loss of normal *ATRX*



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

immunostaining is unlikely to be an oligodendroglioma. Mutations in the promoter region of the *telomerase reverse transcriptase* (*TERT*) gene occur frequently in glioblastomas and oligodendrogliomas.<sup>28,29</sup> *TERT* promoter mutations in gliomas are associated with 1p19q codeletion and *IDH* mutations in oligodendrogliomas.<sup>30</sup> Interestingly, they are also highly characteristic of *IDH*-wt and *ATRX* wild-type glioblastomas, especially those that contain amplification of epidermal growth factor receptor (*EGFR*).<sup>28,29</sup> *H3K27M* mutations in the histone-encoding *H3F3A* gene are mostly found in diffuse midline gliomas in both children and adults.<sup>31</sup> Patients with these *H3K27M* mutated gliomas tend to have a very poor prognosis regardless of histologic appearance, so they are classified as WHO grade IV.<sup>30,31</sup>

Analyses of large databases have also suggested a number of other molecular markers as being potential characteristic/prognostic features of specific subgroups.<sup>11,13,15,18,22,27</sup> Molecular features suggested as markers for subtyping grade II–III gliomas include mutations in *NOTCH1*, *CIC*, and *FUBP1*; mutation in *TP53* and/or overexpression of aberrant *TP53*; *PTEN* loss or promoter methylation; amplification of *EGFR*; and chromosome 7 gain, chromosome 10 loss.<sup>10,12,13,19,30</sup> Due to variability in results across studies, many of these molecular markers are not yet widely used to subclassify gliomas, although the 2020 version of the WHO classification of CNS tumors will include *CDKN2A/B* homozygous deletion as evidence of grade 4 status in *IDH* mutant astrocytomas, as indicated by a recent consensus statement.<sup>32</sup>

### Prognostic Relevance of Molecular Subgroups in Glioma

Numerous large studies of patients with brain tumors have determined that, among grade II–III gliomas, 1p19q codeletion correlates with greatly improved progression-free survival (PFS) and overall survival (OS).<sup>11,15,16,33–35</sup> Likewise, the presence of an *IDH* mutation is a strong favorable prognostic marker for OS in grade II–III gliomas.<sup>12</sup> Analyses

within single treatment arms showed that the *IDH* status is prognostic for outcome across a variety of postoperative adjuvant options. For example, in the NOA-04 phase III randomized trial in newly diagnosed anaplastic gliomas, *IDH* mutation was associated with improved PFS, longer time to treatment failure (TTF), and extended OS in each of the three treatment arms: standard RT (n = 160); combination therapy with procarbazine, lomustine, and vincristine (PCV; RT upon progression; n = 78); and temozolomide (TMZ; RT upon progression; n = 80).<sup>34</sup>

Multiple independent studies have shown that subdividing gliomas by molecular subtype, especially *IDH1/2* and 1p19q status, yields greater prognostic separation than subdivision based on histology (as defined by WHO 2007). These include very large studies covering multiple grades and histology-based subtypes of gliomas,<sup>12,15,33</sup> as well as smaller studies limited to 1 to 2 grades or histologic subtypes.<sup>11,36–38</sup> Multiple studies have also shown that, among patients with grade II–III gliomas, the *IDH*-mut plus 1p19q-codeletion group has the best prognosis, followed by *IDH*-mut without 1p19q codeletion; the *IDH*-wt group has the worst prognosis.<sup>11–13,33–35</sup> Analyses within single treatment arms have confirmed this trend in prognosis across a variety of postoperative adjuvant treatment options.<sup>11,34,35,38</sup> *TERT* mutations in patients with high-grade *IDH*-wt glioma are associated with shorter OS, compared to *IDH*-wt tumors without a *TERT* mutation.<sup>13,29,39</sup> However, a multivariate analysis of data from 291 patients with *IDH*-mut+1p19q-codeleted oligodendrogliomas showed that absence of a *TERT* mutation was associated with worse OS, compared to patients with *TERT*-mut oligodendrogliomas (HR, 2.72; 95% CI, 1.05–7.04; *P* = .04).<sup>40</sup>

MGMT (O-6-methylguanine-DNA methyltransferase) is a DNA repair enzyme that can cause resistance to DNA-alkylating drugs.<sup>41</sup> MGMT promoter methylation is associated with better survival outcomes in patients with high-grade glioma and is a predictive factor for response to



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

treatment with alkylating chemotherapy such as TMZ or lomustine,<sup>30,42-44</sup> even in older adult patients.<sup>45,46</sup> Tumors with *H3K27M* mutations are far less likely to be *MGMT* promoter methylated<sup>31</sup> and are associated with worse prognosis.<sup>47,48</sup> Patients whose glioblastomas contain *H3F3A* G34 mutations, however, may have relatively higher rates of *MGMT* promoter methylation, and do not have a worse prognosis than other *IDH*-wt glioblastomas.<sup>48,49</sup>

Most pilocytic astrocytomas in pediatric patients contain *BRAF* fusions or, less commonly, *BRAF* V600E mutations, especially those arising in the posterior fossa; such tumors are rarely high grade.<sup>50</sup> *BRAF* fusion is associated with better prognosis in pediatric low-grade astrocytoma.<sup>50-52</sup> The likelihood of a *BRAF* fusion in a pilocytic astrocytoma decreases with age.<sup>50</sup> The *BRAF* V600E mutation is present in most pleomorphic xanthoastrocytomas, though it has also been found in some other pediatric low-grade gliomas, such as gangliogliomas and dysembryoplastic neuroepithelial tumors,<sup>30,50,53</sup> as well as a small proportion of glioblastomas (especially epithelioid glioblastoma).<sup>54</sup> Retrospective studies have shown that *BRAF* V600E may be associated with increased risk of progression in pediatric low-grade gliomas,<sup>55</sup> but one study found that this association was not quite statistically significant ( $N = 198$ ;  $P = .07$ ).<sup>52</sup> Some studies have shown that tumors with a *BRAF* V600E mutation may respond to *BRAF* inhibitors such as vemurafenib,<sup>56-58</sup> but ongoing trials will further clarify targeted treatment options in the presence of a *BRAF* fusion or V600E mutation (eg, NCT03224767, NCT03430947). *BRAF* fusion and/or mutation testing are clinically indicated in patients with low-grade glioma.

### NCCN Molecular Testing Recommendations for Glioma

Recommendations for molecular testing of glioma tumors are provided in the *Principles of Brain Tumor Pathology* section in the algorithm. Based on studies showing that *IDH* status is associated with better prognosis in patients with grade II–III glioma,<sup>22,33,34,59</sup> the panel recommends *IDH*

mutation testing in patients with glioma. Immunohistochemistry can detect the most common *IDH* mutation, which is *IDH1* R132H. However, sequencing must be done to detect the less common *IDH1* mutations (eg, *IDH1* R132C) and *IDH2*. This sequencing should be done in the proper clinical context (eg, younger patients with non-enhancing gliomas). Patients with oligodendroglioma should also undergo 1p19q testing. However, since 1p19q codeletion is strongly associated with *IDH* mutation,<sup>22,23,60</sup> 1p19q testing is not necessary in tumors that are definitely *IDH*-wt, and tumors without an *IDH* mutation should not be regarded as 1p19q codeleted, even when results suggest otherwise. Mutation testing for *ATRX* and *TERT* are also recommended, given the diagnostic value of these mutations.<sup>25,27-29</sup> Screening for *H3K27M* mutations (*H3F3A* and *HIST1H3B* sequencing preferred) and *BRAF* fusion and/or mutation testing may be carried out as clinically indicated.

Grade III–IV gliomas should undergo testing for *MGMT* promoter methylation status, since *MGMT* promoter methylated tumors typically respond better to alkylating chemotherapy, compared to unmethylated tumors.<sup>42,45,46,61</sup> To date, there are no targeted agents that have shown improvement in OS in the treatment of glioblastoma. Nevertheless, molecular testing of glioblastomas is still encouraged by the panel, as patients with a detected driver mutation may be treated with a targeted therapy on a compassionate use basis, and these tests improve diagnostic accuracy and prognostic stratification. Detection of genetic or epigenetic alterations could also expand clinical trial options for a brain tumor patient.

### Low-Grade Gliomas

Low-grade gliomas (ie, pilocytic and diffusely infiltrative astrocytomas, oligodendrogliomas) are a diverse group of relatively uncommon malignancies classified as grade I and II under the WHO grading system.<sup>9</sup> Low-grade gliomas comprise approximately 5% to 10% of all CNS tumors.<sup>62</sup> Seizure is a common symptom (81%) of low-grade gliomas, and



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

is more frequently associated with oligodendrogliomas.<sup>63,64</sup> The median duration from onset of symptoms to diagnosis ranges from 6 to 17 months.

### Grade I Gliomas

Diffuse astrocytomas are poorly circumscribed and invasive, and most gradually evolve into higher-grade astrocytomas. Although these were traditionally considered benign, they can behave aggressively and will undergo anaplastic transformation within 5 years in approximately half of patients.<sup>65,66</sup> The most common non-infiltrative astrocytomas are pilocytic astrocytomas. Other grade I gliomas in which treatment recommendations are included in the NCCN Guidelines for CNS Cancers are pleomorphic xanthoastrocytoma, subependymal giant cell astrocytoma (SEGA), and ganglioglioma, though these grade I gliomas are uncommon. Pleomorphic xanthoastrocytomas are associated with favorable prognosis,<sup>67,68</sup> though mitotic index is associated with survival outcomes.<sup>68,69</sup> Gangliogliomas are commonly located in the temporal lobe, and the most significant predictors of survival are low tumor grade and younger age.<sup>70</sup>

SEGAs are typically located at the caudothalamic groove adjacent to the foramen of Monroe. Though they are generally slow-growing and histologically benign, they can also be associated with manifestations such as hydrocephalus, intracranial pressure, and seizures.<sup>71</sup> SEGAs can be distinguished from subependymal nodules by their characteristic serial growth.<sup>72</sup> These tumors occur in 5% to 20% of individuals with tuberous sclerosis complex (TSC).<sup>73-75</sup>

### Treatment

Grade I gliomas are usually curable by surgery alone. Indication for treatment of SEGAs is based on development of new symptoms or radiologic evidence of tumor growth.<sup>72</sup> Though surgery is sometimes a recommended option for SEGAs, many are in an area not amenable to resection, and recurrence may occur following resection.<sup>76,77</sup> Surgery may pose risks because of the frequent location of SEGAs near the foramen of

Monro, but in specialized centers, morbidity is acceptable, and surgical mortality is extremely low.<sup>78</sup>

There is some evidence that BRAF inhibitors, as well as a BRAF/MEK inhibitor combination, may be used for treatment of low-grade gliomas that are BRAF mutated. The phase II VE-BASKET study showed that vemurafenib was efficacious in BRAF-mutated low-grade gliomas, particularly PXA, with an overall response rate (ORR) of 42.9% (n = 7), median PFS of 5.7 months, and median OS not reached.<sup>58</sup> Another phase II trial including 10 patients with low-grade glioma showed that dabrafenib/trametinib was associated with an ORR of 56% (5 patients with a partial response and 4 patients with stable disease).<sup>79</sup> Case reports have demonstrated clinical activity for the combination BRAF/MEK inhibitor dabrafenib/trametinib in patients with *BRAF* V600E mutant glioma.<sup>80,81</sup>

Reducing or stabilizing the volume of SEGAs through systemic therapy has been investigated. A phase III trial showed that 78 patients with SEGA and TSC who received everolimus, an mTOR inhibitor, had at least a 50% reduction in tumor volume, compared to 39 patients who received a placebo (35% vs. 0%;  $P < .001$ ), and 6-month PFS was 100% versus 86%, respectively ( $P < .001$ ).<sup>82</sup> Analyses from a long-term follow-up showed that median duration of response was not reached, with response duration ranging from 2.1 months to 31.1 months.<sup>83</sup> Tumor volume reduction rates of 30% and 50% were maintained in patients in the everolimus arm for more than 3 years. This regimen was generally well-tolerated, with the most frequently reported grade 3 or 4 adverse events being stomatitis (8%) and pneumonia (8%). Everolimus has also been investigated in a phase II trial including 58 patients with recurrent grade II gliomas, with a 6-month PFS rate of 84%.<sup>84</sup> Medical therapy of SEGA, while effective, is a long-term commitment, unless it is being used short-term to facilitate surgical resection. Once mTOR inhibitor therapy is stopped, lesions typically recur, usually within several months, and





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

eventually reach pretreatment volume. The lesions will continue to grow unless therapy is reintroduced. Most patients tolerate long-term therapy with mTOR inhibitors quite well.<sup>85</sup>

### *NCCN Recommendations*

When possible, maximal safe resection is recommended for grade I gliomas, and the actual extent of resection should be documented with a T2-weighted or FLAIR MRI scan within 48 hours after surgery. Patients may be observed following surgery. If incomplete resection or biopsy, or if surgery was not feasible, then RT may be considered if there is significant tumor growth or if neurologic symptoms are present or develop. A BRAF/MEK inhibitor combination may be used for patients with *BRAF* V600E mutant low-grade glioma. Treatment with an mTOR inhibitor (eg, everolimus) should be considered for patients with SEGA,<sup>82,83</sup> though institutional expertise and patient preference should guide treatment decision-making for these rare tumors.<sup>72</sup>

### **Grade II Infiltrative Supratentorial Astrocytoma/Oligodendroglioma**

Radiographically, low-grade oligodendrogliomas appear well demarcated, occasionally contain calcifications, and do not often enhance with contrast. In histology, the typical “fried egg” appearance of these tumors is evident as a fixation artifact in paraffin but not in frozen sections. Grade II oligodendrogliomas have a much better 5-year survival rate (82.7%) than diffuse astrocytomas (51.6%).<sup>86</sup>

Factors prognostic for PFS or OS in patients with grade II gliomas include age, tumor diameter, tumor crossing midline, neurologic status or PS prior to surgery, and the presence of certain molecular markers (see section above on *Molecular Profiling for Gliomas*).<sup>11,16,87-92</sup> For example, *IDH1/2* mutation is associated with a favorable prognosis in patients with grade II and III gliomas,<sup>12,13,34</sup> supporting the emerging idea that molecular analysis should play a larger role in treatment decision-making, relative to histopathology.<sup>64</sup>

### *Treatment Overview*

#### **Surgery**

Surgery remains an important diagnostic and therapeutic modality. The primary surgical goals are maximal safe resection to delay progression and improve survival, relief of symptoms, and provision of adequate tissue for a pathologic diagnosis and grading. Needle biopsies are often performed when lesions are in deep or critical regions of the brain. Biopsy results can be misleading, because gliomas often have varying degrees of cellularity, mitoses, or necrosis from one region to another; thus, small samples can provide erroneous histologic grade or diagnosis.<sup>93,94</sup>

Surgical resection plays an important role in the management of low-grade gliomas. A systematic review showed that GTR was significantly associated with decreased mortality and lower risk of disease progression up to 10 years after treatment, compared to STR.<sup>95</sup> Because these tumors are relatively uncommon, published series generally include patients treated for decades, which introduces additional variables. For example, the completeness of surgical excision was based on the surgeon’s report in older studies. This approach is relatively unreliable when compared with assessment by modern postoperative imaging studies. Furthermore, many patients also received RT, and thus the net effect of the surgical procedure on outcome is difficult to evaluate. Two meta-analyses including studies of primary low-grade gliomas show that extent of resection is a significant prognostic factor for PFS and/or OS.<sup>96,97</sup> Maximal safe resection may also delay or prevent malignant progression<sup>97-99</sup> and recurrence.<sup>100</sup> Patients who undergo an STR, open biopsy, or stereotactic biopsy are, therefore, considered to be at higher risk for progression. GTR is also associated with improved seizure control compared to STR.<sup>97</sup>

Biological considerations also favor an attempt at a complete excision of a low-grade glioma. First, the tumor may contain higher-grade foci, which may not be reflected in a small specimen. Second, complete excision may



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

decrease the risk of future dedifferentiation to a more malignant tumor.<sup>101</sup> Third, removal of a large tumor burden may enhance the benefit of RT. As a result of these considerations, the general recommendation for treating a low-grade glioma is to first attempt as complete an excision of tumor as possible (based on postsurgical MRI verification) without compromising function. However, for tumors that involve eloquent areas, a total removal may not be feasible, and an aggressive approach could result in neurologic deficits. Residual tumor volume may also be a prognostic factor, with a randomized single institution study showing that the OS benefit of maximal safe resection was limited to patients with a residual tumor volume <15 cm<sup>3</sup>.<sup>102</sup>

### Adjuvant Therapy

A large meta-analysis, including data from phase 3 trials (EORTC 22844 and 22845,<sup>103,104</sup> and NCCTG 86-72-51<sup>90</sup>), confirmed that surgery followed by RT significantly improves PFS but not OS in patients with low-grade gliomas.<sup>105</sup> Early versus late postoperative RT did not significantly affect OS, however, suggesting that observation is a reasonable option for some patients with newly diagnosed gliomas.<sup>104</sup>

Final results of a phase 3 randomized clinical trial, RTOG 9802, which assessed the efficacy of adjuvant RT versus RT followed by 6 cycles of PCV in patients with newly diagnosed supratentorial WHO grade II gliomas and at least one of two risk factors for disease progression (STR or age ≥40 years)<sup>106</sup> showed significant improvements in both PFS and OS with the addition of PCV.<sup>107</sup> The median survival time increased from 7.8 years to 13.3 years ( $P = .02$ ), and the 10-year survival rate increased from 41% to 62%. It is important to note, however, that roughly three-quarters of the study participants had a Karnofsky Performance Status (KPS) score of 90 to 100, and the median age was around 40 years.<sup>106</sup> Exploratory analyses based on histologic subgroups showed a statistically significant improvement in OS for all subgroups except for patients with

astrocytoma.<sup>107</sup> Given that the study participants treated with PCV after RT experienced a significantly higher incidence of grade 3 or 4 adverse events (specifically neutropenia, gastrointestinal disorder, and fatigue),<sup>106,107</sup> PCV may be difficult to tolerate in patients who are older or with poor PS. A retrospective subgroup analysis suggests that the survival benefit with the addition of PCV was seen only in *IDH*-mut tumors; the *IDH*-wt subgroup did not appear to benefit from the chemotherapy.<sup>108</sup>

Combined treatment with RT plus TMZ is supported by a phase 2 multicenter trial (RTOG 0424) in patients with supratentorial WHO grade II tumors and additional risk factors (ie, age ≥40 years, astrocytoma, bi-hemispherical, tumor diameter ≥6 cm, neurologic function status >1).<sup>109</sup> However, since the historical controls included patients treated in an earlier time period using different RT protocols, prospective controlled trials are needed to determine whether treatment with TMZ concurrently and following RT is as efficacious as PCV following radiation. There are currently no phase III data to support the use of RT and TMZ over RT and PCV for the treatment of patients with newly diagnosed, high-risk, low-grade glioma. The phase 3 randomized EORTC 22033-26033 trial showed that PFS is not significantly different for adjuvant RT versus dose-dense TMZ in patients with resected or biopsied supratentorial grade II glioma and more than one risk factor ( $N = 477$ ).<sup>17</sup> However, analyses of OS have not yet been reported for this trial.

### Radiation Therapy

When RT is given to patients with low-grade gliomas, it is administered with restricted margins. A T2-weighted (occasionally enhanced T1) and/or FLAIR MRI scan is the best means for evaluating tumor extent, because these tumors enhance weakly or not at all. The clinical target volume (CTV) is defined by the FLAIR or T2-weighted tumor with a 1- to 2-cm margin. Every attempt should be made to decrease the RT dose outside the target volume. This can be achieved with 3-dimensional (3D) planning





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

or intensity-modulated RT (IMRT), with improved target coverage and normal brain/critical structure sparing often shown with IMRT.<sup>110,111</sup> The recommended dosing for postoperative RT is based on results from two phase 3 randomized trials showing that higher dose RT had no significant effect on OS or time to progression,<sup>90,103</sup> and on several retrospective analyses showing similar results.<sup>89,91,112</sup> Because higher doses offer no clear advantages, the CNS Panel recommends lower-dose RT (45–54 Gy) for treatment of low-grade gliomas (grades I/II), including high-risk cases. However, *IDH*-wt low-grade gliomas have similar survival only slightly better than *IDH*-wt glioblastomas.<sup>12</sup> Therefore, an RT dose of 59.4 to 60 Gy may be considered for this subset of patients with low-grade glioma. Preliminary data suggest that proton therapy could reduce the radiation dose to developing brain tissue and potentially diminish toxicities without compromising disease control.<sup>113</sup>

### Recurrent or Progressive Disease

Though the survival impact is unclear, surgery for recurrent disease in patients with low-grade glioma may reduce symptoms, provide tissue for evaluation, and potentially allow for molecular characterization of the tumor.<sup>114-117</sup> Maximal safe resection could play an important role for optimizing survival outcomes; a threshold value is unknown, but >90% extent of resection is suggested.<sup>117</sup> For patients without previous RT, results of the RTOG 9802 trial<sup>106,107</sup> support use of chemotherapy with RT. Data from phase II trials inform recommendations for chemotherapy treatment of patients with recurrent or progressive low-grade glioma.<sup>118-123</sup> Patients should be enrolled in clinical trials evaluating systemic therapy options.

### NCCN Recommendations

#### Primary and Adjuvant Treatment

For treatment recommendations for newly diagnosed grade II gliomas, the panel used the RTOG 9802<sup>106,107</sup> criteria for determining if a patient is

considered to be at low or high risk for tumor progression: patients are categorized as being at low risk if they are 40 years or younger and underwent a GTR; high-risk patients are older than 40 years of age and/or underwent an STR. However, the panel acknowledges that other prognostic factors have been used to guide adjuvant treatment choice in other studies of patients with low-grade glioma,<sup>124</sup> such as tumor size, presence of neurologic deficits, loss of *CDKN2A* homozygous deletion, and the *IDH* mutation status of the tumor.<sup>17,87</sup> If these other risk factors are considered, and treatment of a patient is warranted, then the panel recommends that the patient be treated as high-risk.

Patients with low-risk and low-grade glioma may be observed following surgery. Close follow-up is essential as over half of these patients will develop tumor progression within 5 years.<sup>92</sup> Following surgery, RT followed by PCV is a category 1 recommendation for patients with grade II glioma who are considered to be at high risk for tumor progression, based on the practice-changing results from the RTOG 9802 study,<sup>106,107</sup> as discussed above. There is currently a lack of prospective randomized phase 3 data for the use of radiation and TMZ in patients with low-grade glioma, but interim data from the phase III CATNON trial illustrate that there is a benefit from adjuvant TMZ in patients with newly diagnosed 1p19q non-codeleted anaplastic gliomas.<sup>125</sup> Therefore, RT followed by adjuvant TMZ is a category 2A option. Data from EORTC and NCIC studies, which included patients with glioblastoma, support RT with concurrent and adjuvant TMZ as an evidence-based regimen.<sup>126,127</sup> Therefore, this is also a category 2A option. Because PCV is generally a more difficult chemotherapy regimen to tolerate than TMZ, it may be reasonable to treat an elderly patient or a patient with multiple comorbidities with RT and TMZ instead of RT and PCV, but there are currently no data to show that doing so would result in similar improvement in OS.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Since the design of RTOG 9802<sup>106,107</sup> did not address whether all patients should be treated with RT followed by PCV immediately after a tissue diagnosis (an observation arm was not included for patients with high-risk glioma [defined as older than 40 years of age and/or underwent an STR]<sup>92</sup> in the study), observation after tissue diagnosis may be a reasonable option for some patients with high-risk grade II glioma who are neurologically asymptomatic or who have stable disease. However, close monitoring of such patients with brain MRI is important. Results from EORTC 22845, which showed that treatment with RT at diagnosis versus at progression did not significantly impact OS, provide rationale for observation in select cases with low-grade gliomas as an initial approach, deferring RT.<sup>104</sup> Long-term toxicity from radiation needs to be a consideration, especially for young patients with 1p19q codeletion, for whom there is slightly higher risk of radiation necrosis.<sup>128</sup>

### *Recurrence*

At the time of recurrence, surgery is recommended if resectable disease is present. Because recurrence on neuroimaging may be confounded by treatment effects, biopsy of unresectable disease should be considered to confirm recurrence. There is a propensity for low-grade gliomas to transform to higher-grade gliomas over time. Therefore, documenting the histopathologic transformation of a low-grade glioma to a high-grade glioma may also enable patients to have clinical trial opportunities, since most clinical trials in the recurrent setting are for patients with high-grade gliomas. Moreover, sampling of tumor tissue to confirm recurrence is encouraged to obtain tissue for next-generation sequencing, the results of which may inform treatment selection and/or clinical trial eligibility.

Surgery for recurrent disease may be followed by the following treatment options for patients previously treated with fractionated EBRT: 1) chemotherapy; 2) consideration of reirradiation with or without chemotherapy; and 3) palliative/best supportive care. Reirradiation is a

good choice if the new lesion is outside the target of previous RT or if the recurrence is small and geometrically favorable. For patients with low-risk features for whom GTR was achieved, observation with no further treatment may be considered.

Based on the strength of the RTOG 9802 results,<sup>106,107</sup> RT with chemotherapy is a treatment option for patients with recurrent or progressive low-grade gliomas who have not had prior RT. Options include RT + adjuvant PCV, RT + adjuvant TMZ, and RT + concurrent and adjuvant TMZ. RT alone is generally not the preferred treatment option except in select cases, such as a patient with a poor PS, or who does not want to undergo chemotherapy treatment. Chemotherapy alone (eg, TMZ, PCV, carmustine/lomustine) is also a treatment option for these patients, though this is a category 2B option based on less panel consensus.

### **Anaplastic Gliomas and Glioblastomas**

High-grade gliomas (defined as WHO grade III and IV gliomas) are the most common type of brain cancer, accounting for more than half of all malignant primary tumors of the CNS.<sup>86</sup> Whereas the prognosis for glioblastoma (grade IV glioma) is grim (5-year survival rates between 1%–19%, depending on age), outcomes for anaplastic gliomas (grade III gliomas) are typically better, depending on the molecular features of the tumor.<sup>62</sup> Challenges regarding treatment of glioblastoma include the inability of most systemic therapy agents to penetrate the blood-brain barrier (BBB) and heterogeneity among genetic drivers.<sup>129</sup>

High-grade astrocytomas diffusely infiltrate surrounding tissues and frequently cross the midline to involve the contralateral brain. Patients with these neoplasms often present with symptoms of increased intracranial pressure, seizures, or focal neurologic findings related to the size and location of the tumor and associated vasogenic edema. High-grade astrocytomas usually do not have associated hemorrhage or calcification



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

but can produce considerable edema and mass effect, and they enhance after the administration of intravenous contrast. Tumor cells have been found in peritumoral edema, which corresponds to the T2-weighted MRI abnormalities. Thus, this volume is frequently used to define RT treatment volumes.

It can be challenging to assess the results of therapy by MRI, because the extent and distribution of contrast enhancement, edema, and mass effect are a function of BBB integrity. Thus, factors that increase permeability of the BBB (such as surgery, RT, tapering of corticosteroids, and immunotherapies) can mimic tumor progression radiographically by increasing the presence of contrast enhancement and associated vasogenic edema. Furthermore, anti-VEGF therapy (ie, bevacizumab) suppresses vascular permeability and provides a radiographic appearance of a response, despite residual disease (pseudoresponse).<sup>130</sup>

Anaplastic oligodendrogliomas are relatively rare.<sup>86</sup> While these tumors can be confused with glioblastoma histopathologically, if molecular analysis detects that the tumor is 1p19q codeleted and *IDH1*-mut or *IDH2*-mut, then the tumor is considered to be an anaplastic oligodendroglioma.<sup>9</sup> This distinct subtype has a much better prognosis compared to other high-grade gliomas (anaplastic astrocytomas and glioblastomas).

### Treatment Overview

#### Surgery

The goals of surgery are to obtain a diagnosis, alleviate symptoms related to increased intracranial pressure or compression by tumor, increase survival, and decrease the need for corticosteroids. A meta-analysis including six studies with 1618 patients with glioblastoma showed that GTR is associated with superior OS and PFS, compared to incomplete resection and biopsy.<sup>131</sup> Unfortunately, the infiltrative nature of high-grade astrocytomas frequently renders GTR difficult. There are data suggesting

that resection of all fluid-attenuated inversion recovery (FLAIR) signal abnormalities in high-grade *IDH*-mut gliomas is associated with improved survival.<sup>132</sup> However, a newer and larger study did not find greater benefit of resection in *IDH*-mut tumors compared to *IDH*-wt high-grade gliomas.<sup>133</sup>

Unfortunately, nearly all high-grade gliomas recur. Re-resection at the time of recurrence may improve the outcome for select patients.<sup>134</sup> According to an analysis by Park et al,<sup>135</sup> tumor involvement in specific critical brain areas, poor KPS score, and large tumor volume ( $\geq 50$  cm<sup>3</sup>) were associated with unfavorable re-resection outcomes.

#### Radiation Therapy

Conformal RT (CRT) techniques, which include 3D-CRT and IMRT are recommended for performing focal brain irradiation. IMRT often will provide superior dosimetric target coverage and better sparing of critical structures than 3D-CRT.<sup>111</sup> Several randomized controlled trials conducted in the 1970s showed that radiation improved both local control and survival in patients with newly diagnosed high-grade gliomas.<sup>136,137</sup> Sufficient radiation doses are required to maximize this survival benefit. However, radiation dose escalation alone above 60 Gy has not been shown to be beneficial.<sup>138</sup> The recommended radiation dose for high-grade astrocytomas is 60 Gy in 2.0 Gy fractions or 59.4 Gy in 1.8 Gy fractions with an initial RT plan to 46 Gy in 2 Gy fractions or 45 to 50.4 Gy in 1.8 fractions, respectively, followed by a boost plan of 14 Gy in 2 Gy fractions or 9 to 14.4 Gy in 1.8 Gy fractions, respectively.<sup>138</sup>

Anaplastic oligodendrogliomas are conventionally treated with the same dose of radiation as high-grade astrocytomas; however, given the better prognosis in patients with anaplastic oligodendroglioma, radiation treatments are generally administered in a lower dose per fraction (1.8 Gy/fraction vs. 2.0 Gy/fraction) to theoretically decrease the risk of late side effects. Accordingly, as per trials such as RTOG 9813,<sup>59</sup> these gliomas are treated to 50.4 Gy in 1.8 Gy fractions for 28 fractions followed



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

by a five-fraction boost of 1.8 Gy/fraction to a total of 59.4 Gy. RT targets for high-grade gliomas are generated from a gross tumor volume (GTV), CTV, and planning target volume (PTV). The GTV encompasses any gross tumor remaining after maximal safe resection as well as the surgical cavity as determined by postoperative imaging. Strategies for GTV definition vary with respect to the inclusion of edema in an initial target volume. When edema is included in an initial phase of treatment, fields are usually reduced for the last phase of treatment. The CTV is an expansion of the GTV by adding an approximately 2-cm margin for grade III and IV gliomas (although smaller CTV expansions are supported in the literature and can be appropriate) to account for a non-enhancing tumor. The CTV is then expanded to a PTV to account for daily setup errors and image registration. The boost target volume will typically encompass only the gross residual tumor and the resection cavity.

Special attention has been given to determining the optimal therapy in older adults with glioblastoma, given their especially poor prognosis, often limited functional status, and increased risk of developing side effects. Overall, the approach in these patients has been to reduce treatment time while maintaining treatment efficacy. Roa et al randomized patients 60 years or older with a poor PS (KPS < 70) to 60 Gy in 30 fractions given over 6 weeks versus 40 Gy in 15 fractions given over 3 weeks and found no difference in survival between these two regimens.<sup>139</sup> However, fewer patients who received 40 Gy over a shorter time period required a post-treatment increase in corticosteroid dose, compared to the patients who received 60 Gy over the longer time period (23% vs. 49%, respectively;  $P = .02$ ). A subsequent study also supports using a regimen of 34 Gy in 10 fractions over 2 weeks in older adult patients.<sup>45</sup> Moreover, another study performed by Roa et al showed that an even shorter course of focal brain radiation consisting of 25 Gy in 5 fractions over 1 week is a reasonable alternative to 40 Gy in 15 fractions over 3 weeks in patients with newly diagnosed glioblastoma who have a poor prognosis (ie, patients who are

older adults and/or frail).<sup>140</sup> However, this was a small study that had some limitations, notably overly broad eligibility criteria and poorly defined non-inferiority margin.<sup>141,142</sup>

A randomized trial of hypofractionated RT (40 Gy given over 3 weeks) with concurrent and adjuvant TMZ versus hypofractionated RT alone in patients 65 years and older showed an improvement in median OS and PFS with the addition of concurrent and adjuvant TMZ (5-year OS of 9.8% vs. 1.9%, respectively; median OS of 14.6 months vs. 12.1 months, respectively; HR for mortality, 0.63; 95% CI, 0.53–0.75;  $P < .001$ ; 5-year PFS of 4.1% vs. 1.3%, respectively; HR, 0.56; 95% CI, 0.47–0.66;  $P < .001$ ).<sup>143</sup> The largest benefit was noted in patients with MGMT promoter methylation (see discussion of *Systemic Therapy for Glioblastoma*, below). Of note, a comparison of standard focal brain radiation (60 Gy given over 6 weeks) with concurrent and adjuvant TMZ versus hypofractionated radiation (40 Gy given over 3 weeks) with concurrent and adjuvant TMZ in elderly patients has not been performed in patients 65 years and older. Therefore, standard radiation (60 Gy given over 6 weeks) with concurrent and adjuvant TMZ (with or without alternating electric field therapy; see discussion of this treatment option below) is also a reasonable treatment option for an older adult patient who has a good PS and wishes to be treated aggressively. Ultimately, quality of life remains an important consideration in the optimal management of this patient population.

### Systemic Therapy

#### Anaplastic Oligodendroglioma

The addition of PCV to RT for the treatment of newly diagnosed anaplastic oligodendrogliomas is supported by results from two phase III trials, one which tested RT followed by PCV for 6 cycles (EORTC 26951<sup>144,145</sup>) and the other which assessed 4 cycles of dose-intensive PCV administered prior to RT (RTOG 9402<sup>35,146,147</sup>). Both studies compared the combination therapy to RT alone and found significant increases in median OS when





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

PCV was added to RT for the upfront management of 1p19q codeleted tumors.

The EORTC 26951 trial showed that, among the entire group of 368 histopathologically diagnosed study patients with anaplastic oligodendroglioma or anaplastic oligoastrocytoma, RT followed by 6 cycles of PCV significantly improved median PFS and OS (42.3 vs. 30.6 months; HR, 0.75; 95% CI, 0.60–0.95;  $P = .018$ ) compared with RT alone.<sup>145</sup> Moreover, in an exploratory subgroup analysis of the 80 patients whose tumors were 1p19q codeleted, the benefit was even more pronounced (OS not reached in the RT + PCV group vs. 112 months in the RT group; HR, 0.56; 95% CI, 0.31–1.03).<sup>22,144,145</sup>

RTOG 9402 randomized 291 patients with histopathologically diagnosed anaplastic oligodendroglioma or anaplastic oligoastrocytoma to treatment with an intensive PCV regimen followed by RT or RT alone.<sup>147</sup> In contrast to the EORTC 26951 study, no difference in median OS was observed between the two arms (4.6 years vs. 4.7 years; HR, 0.79; 95% CI, 0.60–1.04;  $P = .10$ ). However, an unplanned subgroup analysis of the 126 patients whose tumors were 1p19q codeleted found a doubling in median OS (14.7 vs. 7.3 years; HR, 0.59; 95% CI, 0.37–0.95;  $P = .03$ ) when PCV was added to RT as upfront treatment.

As would be predicted, in both studies toxicity was higher in the treatment arms that included PCV. In EORTC 26951, 70% of patients in the RT followed by PCV arm did not complete the planned six cycles of treatment.<sup>144,145</sup> In RTOG 9402, there was also a high rate of study treatment discontinuation and acute toxicities (mainly hematologic), including two early deaths attributed to PCV-induced neutropenia.<sup>146,147</sup> Given the similar efficacy results of the two studies, and the two deaths that occurred from the intensive PCV regimen in RTOG 9402, PCV administered after RT is optimal, as per EORTC 26951.

The phase III CODEL study was designed to assess the efficacy of TMZ for the treatment of newly diagnosed anaplastic oligodendrogliomas. The initial treatment arms were RT alone, RT + TMZ, and TMZ alone. Initial results showed that patients who received TMZ alone had significantly shorter PFS than patients treated with RT (either RT alone or with TMZ) (2.9 years vs. not reached, respectively; HR, 3.12; 95% CI, 1.26–7.69;  $P = .009$ ).<sup>148</sup> When the results of RTOG 9402 and EORTC 26951 were reported showing significant improvement in median OS with RT + PCV upfront, the CODEL study was redesigned to compare RT + PCV to RT + TMZ in patients with anaplastic oligodendroglioma as well as low-grade oligodendroglioma. This study is ongoing.

### **Anaplastic Astrocytoma**

The RTOG 9813 trial showed that RT with concurrent TMZ resulted in similar outcomes as RT with concurrent nitrosourea (either CCNU [lomustine] or BCNU [carmustine]) therapy in patients with newly diagnosed anaplastic astrocytomas, with perhaps slightly better PFS with TMZ (HR, 0.70; 95% CI, 0.50–0.98;  $P = .039$ ).<sup>59</sup> However, the toxicity of nitrosourea was significantly worse than for TMZ, and resulted in higher rates of discontinuation due to toxicity (79% vs. 40%, respectively;  $P < .001$ ). The ongoing CATNON phase 3 randomized trial is testing RT alone, as well as RT with adjuvant TMZ, concurrent TMZ, or both, in patients with newly diagnosed anaplastic astrocytoma. An initial interim analysis showed that adjuvant TMZ significantly improved PFS (HR, 0.62; 95% CI, 0.50–0.76) and OS (HR, 0.67; 95% CI, 0.51–0.88).<sup>125</sup> Median OS for the group of patients treated with post-RT TMZ had not been reached, but median OS at 5 years was 55.9% (95% CI, 47.2–63.8) with and 44.1% (36.3–51.6) without adjuvant TMZ. A second interim analysis showed that patients with *IDH*-mut anaplastic astrocytomas benefit from treatment with adjuvant TMZ (HR, 0.41; 95% CI, 0.27–0.64), but not those with tumors that are *IDH*-wt (HR, 1.05; 95% CI, 0.73–1.52).<sup>149</sup> There was also no definite benefit to concurrent TMZ in patients with *IDH*-mut anaplastic



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

astrocytomas (HR, 0.71; 95% CI, 0.35–1.42;  $P = .32$ ). However, the findings from the second interim analysis are currently available in abstract form only. Further follow-up and molecular analyses are ongoing.

### ***Glioblastoma***

Adjuvant involved-field RT with concurrent and adjuvant TMZ is the standard recommended treatment for patients with newly diagnosed glioblastoma and good PS based on the results of the phase III, randomized EORTC-NCIC study of 573 patients with newly diagnosed glioblastoma who were aged  $\leq 70$  years and had a WHO PS  $\leq 2$ .<sup>143</sup> Patients received either 1) daily TMZ administered concomitantly with postoperative RT followed by 6 cycles of adjuvant TMZ; or 2) RT alone. The chemoradiation arm resulted in a statistically better median survival (14.6 vs. 12.1 months) and 2-year survival (26.5% vs. 10.4%) when compared with RT alone. Final analysis confirmed the survival advantage at 5 years (10% vs. 2%).<sup>143</sup> However, the study design does not shed light on which component is responsible for the improvement: TMZ administered with RT, TMZ following RT, or possibly both.

The TMZ dose used in the EORTC-NCIC trial is 75 mg/m<sup>2</sup> daily concurrent with RT, then 150 to 200 mg/m<sup>2</sup> post-irradiation on a 5-day schedule every 28 days. Alternate schedules, such as a 75 to 100 mg/m<sup>2</sup> for 21 out of 28 days regimen or 50 mg/m<sup>2</sup> daily, have been explored in a phase II trial for newly diagnosed glioblastoma.<sup>150</sup> However, a comparison of the dose-intensive 21/28 and standard 5/28 schedules in the RTOG 0525 phase III study showed no difference in PFS, OS, or by MGMT methylation status with the post-radiation dose-intensive TMZ, compared to the standard post-radiation TMZ dose.<sup>151</sup> A pooled analysis of individual patient data from four randomized trials<sup>127,151-153</sup> of patients with newly diagnosed glioblastoma determined that treating with post-radiation TMZ beyond six cycles does not improve OS, even for patients whose tumors are MGMT promoter methylated.<sup>154</sup> A recent prospective, randomized phase II study

showed no improvement in 6-month PFS, PFS, or OS with continuing treatment with TMZ beyond 6 cycles, and doing so was associated with greater toxicity.<sup>155</sup>

### **MGMT Promoter Methylated Glioblastoma**

The presence of MGMT promoter methylation in glioblastoma is both a prognostic marker and a predictive one for response to treatment with alkylating agents. In the small (N = 31), single-arm phase II UKT-03 trial,<sup>156,157</sup> postoperative RT and TMZ combined with lomustine in patients with newly diagnosed glioblastoma resulted in a median OS of 34.3 months,<sup>156</sup> which compared favorably to the historical control data of 23.4 months in patients with MGMT promoter methylated tumors who were treated with RT and TMZ in the EORTC-NCIC trial.<sup>143</sup> Based on this improvement in survival with combination alkylating agents in patients with MGMT promoter methylated glioblastoma, the phase III CeTeG/NOA-09 trial randomized patients with newly diagnosed MGMT promoter methylated glioblastoma (aged 18–70 and KPS  $\geq 70$ ) to treatment with RT and lomustine + TMZ or RT and TMZ alone.<sup>158</sup> Analysis of the modified intent-to-treat population (N = 129) showed that OS was significantly improved in the TMZ + lomustine arm versus the TMZ arm (median OS of 48.1 months vs. 31.4 months, respectively;  $P = .049$ ). Of note, PFS was not significantly improved, which the investigators hypothesized could have been due to a higher incidence of pseudoprogression in the TMZ + lomustine arm. Grade 3 and 4 adverse events were only slightly higher in the TMZ + lomustine arm (59% vs. 51%, respectively), but the study was too small to adequately define the toxicity profile of RT with TMZ + lomustine. Analysis of health-related quality of life showed no significant differences between the study arms.<sup>159</sup>

### **Older Adults**

Building on the findings that hypofractionated RT alone has similar efficacy and is better tolerated compared to standard RT alone in older adults with





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

newly diagnosed glioblastoma, a phase III randomized trial with 562 newly diagnosed patients 65 years of age or older compared hypofractionated RT with concurrent and adjuvant TMZ to hypofractionated radiation alone. Patients in the combination therapy arm had better PFS (5.3 months vs. 3.9 months; HR, 0.50; 95% CI, 0.41–0.60;  $P < .001$ ) and median OS (9.3 months vs. 7.6 months; HR, 0.67; 95% CI, 0.56–0.80;  $P < .001$ ) compared to patients treated with hypofractionated RT alone.<sup>126</sup> The greatest improvement in median OS was seen in patients with MGMT promoter methylated tumors (13.5 months RT + TMZ vs. 7.7 months RT alone; HR, 0.53; 95% CI, 0.38–0.73;  $P < .001$ ). The benefit of adding TMZ to RT was smaller in patients with MGMT promoter unmethylated tumors and did not quite reach statistical significance (10.0 months vs. 7.9 months, respectively; HR, 0.75; 95% CI, 0.56–1.01;  $P = .055$ ;  $P = .08$  for interaction).

Two phase III studies in elderly newly diagnosed glioblastoma patients assessed treatment with TMZ alone versus radiation.<sup>45,46</sup> The Nordic trial randomized 291 patients aged 60 years and older with good PS across three treatment groups: TMZ, hypofractionated RT, or standard RT.<sup>45</sup> Patients older than 70 years had better survival with TMZ or hypofractionated RT compared to standard RT, and patients whose tumors were MGMT promoter methylated benefitted more from treatment with TMZ compared to patients with MGMT promoter unmethylated tumors (median OS 9.7 vs. 6.8 months; HR, 0.56; 95% CI, 0.34–0.93;  $P = .02$ ). The NOA-08 study assessed the efficacy of TMZ alone compared to standard RT in 373 patients aged 65 years and older.<sup>46</sup> TMZ was found to be non-inferior to standard RT; median OS was similar in both groups (8.6 months in the TMZ arm vs. 9.6 months in the standard RT arm; HR, 1.09; 95% CI, 0.84–1.42;  $P$  [non-inferiority] = .033). For patients whose tumors were MGMT promoter methylated, event-free survival was longer with TMZ treatment compared to standard RT (8.4 months vs. 4.6 months). Neither the Nordic trial nor the NOA-08 trial included a combination RT

and TMZ control arm, which is the treatment regimen typically offered to patients who are fit enough to tolerate it, regardless of age. Although radiation in combination with TMZ is recommended over single-modality therapy for newly diagnosed patients with glioblastoma who are older than 70 years of age and have good PS, the results of these two phase III studies support the recommendation that TMZ alone as initial therapy may be a reasonable option for those elderly patients who have MGMT promoter methylated tumors and would initially prefer to delay treatment with radiation.<sup>45,46</sup>

### ***Alternating Electric Field Therapy***

In 2015, the FDA approved alternating electric field therapy for the treatment of patients with newly diagnosed glioblastoma based on the results of the open-label phase III EF-14 clinical trial. This portable medical device generates low-intensity alternating electric fields to stop mitosis/cell division. In the EF-14 trial, 695 patients with newly diagnosed glioblastoma and good PS (KPS  $\geq 70$ ) were randomized to TMZ alone on a 5/28-day schedule or the same TMZ and alternating electric field therapy, following completion of standard focal brain radiation and daily TMZ.<sup>160</sup> The results of the study showed an improvement in median PFS (6.7 vs. 4.0 months, respectively; HR, 0.63; 95% CI, 0.52–0.76;  $P < .001$ ) and OS (20.9 vs. 16.0 months, respectively; HR, 0.63; 95% CI, 0.53–0.76;  $P < .001$ ) in patients who received TMZ plus alternating electric field therapy.<sup>161</sup> The number of adverse events was not statistically different between the two treatment groups except for a greater frequency of mild to moderate local skin irritation/itchiness in the patients treated with the alternating electric fields.<sup>162</sup> There was no increased frequency of seizures.<sup>163,164</sup> Based on the results of this study, concurrent treatment with adjuvant TMZ and alternating electric fields is a category 1 recommendation for newly diagnosed glioblastoma patients 70 years of age or younger who have a good PS. This is also considered a reasonable treatment option for patients older than 70 years of age with good PS and newly diagnosed



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

glioblastoma who are treated with standard focal brain radiation and concurrent daily TMZ.

### *Therapy for Recurrence*

Patients with malignant gliomas eventually develop tumor recurrence or progression. Surgical resection of locally recurrent disease is reasonable followed by treatment with chemotherapy. Unfortunately, there is no established second-line therapy for recurrent gliomas. If there has been a long time interval between stopping TMZ and development of tumor progression, it is reasonable to restart a patient on TMZ,<sup>165</sup> particularly if the patient's tumor is MGMT methylated. Similarly, a nitrosourea, such as carmustine or lomustine,<sup>166-169</sup> would be a reasonable second-line therapy, especially in a patient whose tumor is MGMT methylated. Although no studies of bevacizumab in patients with recurrent glioblastoma have demonstrated an improvement in survival, bevacizumab is FDA approved for the treatment of recurrent glioblastoma based on improvement in PFS.<sup>170-172</sup> Of note, improvement in PFS may be due to bevacizumab's ability to decrease BBB permeability (resulting in less contrast enhancement and vasogenic edema) rather than a true anti-tumor effect.<sup>173,174</sup> Treatment with regorafenib for recurrent glioblastoma is supported by the results of a randomized phase II trial in which OS was greater for patients randomized to receive regorafenib, compared to those who received lomustine (median OS of 7.4 months vs. 5.6 months, respectively; HR, 0.50; 95% CI, 0.33–0.75;  $P < .001$ ).<sup>175</sup> Of note, the median OS in the lomustine arm in this trial was lower than reported in other randomized phase II and III trials. A phase III study of regorafenib is being planned.

Other routes of chemotherapy delivery have been evaluated. Local administration of carmustine using a biodegradable polymer (wafer) placed intraoperatively in the surgical cavity has demonstrated a statistically significant improvement in survival for patients with recurrent

high-grade gliomas (31 vs. 23 weeks; adjusted HR, 0.67;  $P = .006$ ).<sup>176</sup> Patients who receive carmustine wafers are at greater risk for seizures and postoperative infections. When wafers are used, it is important to achieve a watertight dural closure and have sufficient use of steroids and antiepileptics in the perioperative period to prevent adverse events.<sup>177</sup> Clinicians and patients should be aware that treatment with the carmustine wafer may prevent participation in a clinical trial involving a locally delivered investigational agent.

Alternating electric field therapy is also FDA approved for treating recurrent glioblastoma based on the safety results of this medical device from the EF-11 clinical trial.<sup>178</sup> This phase III study randomized 237 patients with recurrent glioblastoma to alternating electric field therapy or the treating oncologist's choice of chemotherapy. The study did not meet its primary endpoint of demonstrating an improvement in survival in the cohort of patients treated with alternating electric field therapy. Although median OS was similar in both of the treatment arms (6.6 vs. 6 months), the study had not been powered for a non-inferiority determination. Due to lack of clear efficacy data for alternating electric field therapy in EF-11, the panel is divided about recommending it for the treatment of recurrent glioblastoma. Similarly, re-irradiation may be reasonable to consider for some recurrent glioblastoma patients, but the panel is also divided about this option. A systematic review including 50 non-comparative studies of 2095 patients with recurrent glioblastoma who were treated with re-irradiation showed pooled 6- and 12-month OS rates of 73% and 36%, respectively, and 6- and 12-month PFS rates of 43% and 17%, respectively.<sup>179</sup> Over half of the studies (29 out of 50) were rated as poor quality, indicating a need for better quality studies in this area. Further, there is no recommended dose or type of radiation used in the recurrent setting due to inconsistent trial design among these studies.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### ***NCCN Recommendations***

#### ***Primary Treatment***

When a patient presents with a clinical and radiologic picture suggestive of a high-grade glioma, neurosurgical input is needed regarding the feasibility of maximal safe resection. For first-line treatment of high-grade glioma, the NCCN Guidelines recommend maximal safe resection whenever possible. Use of intraoperative MRI and intraoperative fluorescence-guided surgery with 5-aminolevulinic acid (5-ALA) may potentially allow for more complete resection.<sup>180,181</sup> One exception is when CNS lymphoma is suspected; a biopsy should be performed before steroids are administered, and management should follow the corresponding pathway if the diagnosis is confirmed. When maximal resection is performed, the extent of tumor debulking should be documented with a postoperative MRI scan with and without contrast performed within 48 hours after surgery. Multidisciplinary consultation is encouraged once the pathology is available.

#### ***Adjuvant Therapy***

RT is generally recommended after maximal safe resection for the treatment of high-grade gliomas to improve local control and survival. For postoperative treatment of anaplastic gliomas in patients with good PS (KPS  $\geq 60$ ), combination therapy with focal brain radiation combined with PCV or TMZ are among the recommended options. For patients with anaplastic oligodendroglioma, RT plus PCV, given before or after RT, is preferred, based on the results of the RTOG 9402<sup>35,147</sup> and EORTC 26951 studies.<sup>144,145</sup> The panel advises administering PCV after RT as per EORTC 26951 instead of the dose-intensive PCV used prior to RT in the RTOG 9402 study<sup>147</sup> due to better patient tolerance. RT, with or without concurrent TMZ, followed by adjuvant TMZ is also a reasonable option,<sup>182</sup> particularly if it is predicted that the patient might have significant difficulty tolerating PCV due to age or coexisting medical conditions. The panel

awaits the results of CODEL to see if treatment with TMZ will be as efficacious as PCV in this patient population.

In the case of patients with anaplastic astrocytoma and anaplastic oligoastrocytoma (NOS) and good PS, RT, with or without concurrent TMZ and followed by adjuvant TMZ, is preferred based on the first interim analysis results of the CATNON trial showing improvement in survival with RT followed by 12 cycles of TMZ compared to RT alone.<sup>125</sup> However, for newly diagnosed anaplastic oligoastrocytoma patients, RT with PCV administered before or afterwards is also an acceptable treatment option.<sup>183,184</sup>

For patients with anaplastic gliomas and a poor PS (KPS  $< 60$ ), treatment options recommended in the NCCN Guidelines are limited to single-modality therapies due to concerns about the ability of these patients to tolerate the toxicity associated with combination regimens. Patients with a poor PS can be managed by RT (hypofractionation is preferred over standard fractionation), TMZ alone (considered for patients whose tumors are MGMT promoter methylated but is a category 2B option), or palliative/best supportive care.

For patients diagnosed with glioblastoma, the adjuvant options mainly depend on the patient's age, PS (as defined by KPS), and MGMT promoter methylation status.<sup>42,45,143,185</sup> Category 1 recommendations for patients aged 70 years and younger with a good PS, regardless of the tumor's MGMT methylation status, include standard brain RT plus concurrent and adjuvant TMZ with or without alternating electric field therapy. Because patients with newly diagnosed MGMT promoter unmethylated glioblastoma are likely to receive less benefit from TMZ, RT alone is included as a reasonable option, particularly if the patient is eligible to participate in a clinical trial, which omits the use of upfront TMZ.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Category 1 treatment recommendations for patients older than 70 years of age with newly diagnosed glioblastoma, a good PS, and MGMT promoter methylated tumors include hypofractionated brain RT plus concurrent and adjuvant TMZ<sup>126</sup> or standard brain RT plus concurrent and adjuvant TMZ and alternating electric field therapy. For those patients older than 70 years with newly diagnosed glioblastoma, a good PS, and with MGMT unmethylated or indeterminant tumors, hypofractionated brain radiation with concurrent and adjuvant TMZ<sup>126</sup> is preferred, but standard brain RT plus concurrent and adjuvant TMZ and alternating electric field therapy is also a reasonable option (category 1)<sup>160,161</sup> for those elderly patients who want to be treated as aggressively as possible. The complete list of recommendations that the panel did not consider category 1 can be found in the treatment algorithms for patients with glioblastoma who are older than 70 years.

Treatment recommendations for patients with newly diagnosed glioblastoma and KPS below 60 (regardless of age) include hypofractionated brain RT possibly with concurrent and adjuvant TMZ for patients aged 70 years or younger, TMZ alone (for patients with MGMT promoter methylated tumors), or palliative/best supportive care.

### *Follow-up and Recurrence*

Patients should be followed closely with serial brain MRI scans (at 2–8 weeks post-irradiation, then every 2–4 months for 3 years, then every 3–6 months indefinitely) after the completion of treatment for newly diagnosed disease. Scans may appear worse during the first 3 months or longer after completion of RT even though there may be no actual tumor progression.<sup>129</sup> This finding of “pseudoprogression” occurs more often in patients whose tumors are MGMT promoter methylated.<sup>186,187</sup> Early MRI scans allow for appropriate titration of corticosteroid doses based on the extent of mass effect and brain edema. Later scans are used to identify tumor recurrence. Early detection of recurrence is warranted, because

local and systemic treatment options are available for patients with recurrent disease. Biopsy, MR spectroscopy, MR perfusion, or brain PET/CT can be considered to try to determine if the changes seen on brain MRI are due to pseudoprogression or RT-induced necrosis versus actual disease progression.<sup>188,189</sup> RT-induced necrosis tends to be detected between 6 and 24 months following RT treatment.<sup>187</sup>

Management of recurrent tumors depends on the extent of disease and patient condition. The efficacy of current treatment options for recurrent disease remains poor; therefore, enrollment in a clinical trial, whenever possible, is preferred for the management of recurrent disease. Preferred chemotherapy options for recurrent disease include re-treatment with TMZ (if there has been a long interval between completion of adjuvant TMZ and development of recurrent disease),<sup>119,165,190-192</sup> carmustine/lomustine,<sup>166-169,193</sup> bevacizumab,<sup>170,194-199</sup> regorafenib,<sup>175</sup> and PCV.<sup>120,200,201</sup> A patient with a poor PS should receive palliative/best supportive care.

### **Intracranial and Spinal Ependymomas**

Ependymomas constitute up to 1.9% of adult CNS tumors and 5.7% of pediatric CNS tumors.<sup>86</sup> In adults, ependymomas occur more often in the spinal canal than in the intracranial compartment (supratentorial and posterior fossa). These tumors can cause hydrocephalus and increased intracranial pressure, mimic brainstem lesions, cause multiple cranial nerve palsies, produce localizing cerebellar deficits, and cause neck stiffness and head tilt if they infiltrate the upper portion of the cervical cord.<sup>202,203</sup>

RELA activating fusions occur in about 19% of patients with ependymomas.<sup>204</sup> They occur only in supratentorial ependymomas and not in those in the posterior fossa or spinal canal; they are also more likely to occur in children than in adults.<sup>204</sup> Ependymomas with RELA activating fusions are more likely to be advanced and aggressive than RELA fusion-





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

negative ependymomas, with a greater likelihood of being grade II or III, and with shorter PFS and OS.<sup>204,205</sup> In the revised 2016 WHO classification system, RELA fusion-positive ependymoma is now designated as a subtype,<sup>9</sup> raising the potential for RELA fusion testing of ependymomas based on the clinical context.

This section focuses on adult spinal and intracranial ependymal tumors including grade I (subependymomas or myxopapillary ependymomas), grade II differentiated (classic ependymomas), and grade III (anaplastic ependymomas) tumors.

### Treatment Overview

#### *Surgery*

There is a paucity of robust studies addressing the role of surgery in this uncommon disease, but multiple case series have reported that patients with totally resected tumors tend to have the best survival for both low- and high-grade ependymomas.<sup>206-210</sup> Grade I subependymomas are non-infiltrative and can often be cured by resection alone. For myxopapillary ependymomas, complete resection of the mass without capsular violation (marginal en bloc resection) can be curative.<sup>211</sup> In a retrospective analysis by Rodriguez et al,<sup>212</sup> patients who underwent surgery had a better outcome than those who did not (HR, 1.99;  $P < .001$ ). Supratentorial ependymomas generally have a poorer prognosis than their infratentorial counterparts, because a greater proportion of supratentorial lesions are of high grade.

#### *Radiation Therapy*

The survival benefits of RT following surgery have been established for anaplastic ependymomas and suboptimally resected tumors, although much of the data are derived from pediatric patients. Rodriguez et al<sup>212</sup> reviewed over 2400 cases of ependymomas in the SEER database and reported that patients with partially resected tumors who do not receive RT

have a poorer prognosis than those who are treated with RT (HR, 1.75;  $P = .024$ ). The short-term and 10-year survival rate after RT reached over 70% and 50%, respectively.<sup>213-215</sup> The value of RT is more controversial for differentiated ependymomas,<sup>207,216</sup> with data demonstrating improved survival mainly for subtotally resected tumors.<sup>208,212</sup> Emerging data show poor survival rates in patients with supratentorial non-anaplastic ependymoma who do not receive RT following GTR.<sup>217</sup> Further, much of the data supporting observation following surgical resection are based on retrospective studies.<sup>218-220</sup> Given the availability of highly CRT modalities and the relatively lower level of concerns for late effects of RT in adults (vs. children), RT is recommended as the standard adjuvant treatment approach in these patients until high-quality evidence supporting observation alone becomes available.

In the past, the standard practice was to irradiate the entire craniospinal axis or administer WBRT. However, studies have demonstrated that: 1) local recurrence is the primary pattern of failure; 2) spinal seeding is uncommon in the absence of local failure; 3) the patterns of failure are similar in patients with high-grade tumors who are treated with local fields or craniospinal axis irradiation; and 4) spinal metastases may not be prevented by prophylactic treatment.<sup>221-223</sup> Prophylactic craniospinal RT or WBRT does not lead to improvement in survival compared to conformal regional RT with higher doses in modern studies of non-disseminated disease.<sup>209,216,224</sup>

Typical craniospinal irradiation scheme includes 36 Gy in 1.8 Gy fractions to the whole brain and spine, followed by limited-field irradiation to spine lesions to 45 Gy. For intracranial ependymomas, the primary brain site should receive a total of 54 to 59.4 Gy in 1.8 to 2.0 Gy fractions. PTV of margin of 3 to 5 mm is typically added to the CTV. Tolerance of the cauda equina is in the range of 54 to 60 Gy.<sup>225,226</sup> Therefore, a boost to gross



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

intracranial metastatic sites (respecting normal tissue tolerances) may be considered.

For spinal ependymomas, patients could receive local RT to 45 to 54 Gy in 1.8 Gy fractions, with higher doses up to 60 Gy being reasonable for spinal tumors below the conus medullaris. These dosing recommendations are consistent with those for primary spinal cord tumors. However, it is important to note that retrospective analyses have shown that adjuvant RT does not consistently improve disease outcomes in patients with these tumors.<sup>227,228</sup>

Proton beam craniospinal irradiation may be considered when clinically appropriate and when toxicity is a concern. SRS has been used as a boost after EBRT or to treat recurrence with some success, although data on long-term results are still lacking.<sup>229-231</sup>

### **Systemic Therapy**

Studies regarding the role of chemotherapy have largely been in the setting of pediatric ependymomas; the role of chemotherapy in the treatment of ependymomas in adult patients remains poorly defined. No study has demonstrated a survival advantage with the addition of chemotherapy to RT in newly diagnosed tumors. However, chemotherapy is sometimes considered as an alternative to palliative/best supportive care or RT in the recurrence setting. Possible options include platinum-based regimens (cisplatin or carboplatin),<sup>232,233</sup> etoposide,<sup>234,235</sup> nitrosourea-based regimens (lomustine or carmustine),<sup>233</sup> bevacizumab,<sup>236</sup> and temozolomide.<sup>237</sup> The combination of lapatinib, a tyrosine kinase inhibitor (TKI), and dose-dense TMZ has been evaluated in a phase II trial in patients with recurrent grade I–III ependymoma, with preliminary results reported only in abstract form.<sup>238</sup>

## **NCCN Recommendations**

### ***Primary and Adjuvant Treatment***

In general, when feasible, management of rare tumors such as ependymomas should begin with a timely and early consultation with centers of neuro-oncologic expertise. Whenever possible, maximal safe resection should be attempted with contrast-enhanced brain image verification within 48 hours after surgery. Spine MRI, if not done prior to surgery, should be delayed by at least 2 to 3 weeks after surgery to avoid post-surgical artifacts. If maximal resection is not feasible at diagnosis, STR or biopsy (stereotactic or open) should be performed. Due to the established relationship between the extent of resection and outcome, multidisciplinary review and re-resection (if possible) should be considered if MRI shows that initial resection is incomplete. For spinal myxopapillary ependymomas, en bloc resection without capsule violation is recommended whenever feasible.

The adjuvant treatment algorithm depends on the extent of surgical resection, histology, and staging by craniospinal MRI and cerebrospinal fluid (CSF) cytology. For spinal ependymomas, brain MRI should be obtained to determine if these are drop metastases from a primary brain lesion. CSF dissemination develops in up to 15% of intracranial ependymomas. Lumbar puncture for CSF cytology, which is indicated when there is clinical concern for meningeal dissemination, should be done following spine MRI and, if not done prior to surgery, should be delayed at least 2 weeks after surgery to avoid a false-positive result. Lumbar puncture may be contraindicated in some cases (for example, if there is increased intracranial pressure and risk of herniation).

RT is the appropriate postoperative management for patients with negative findings for tumor dissemination on MRI scans and CSF analysis. Patients with grade I spinal ependymomas that have been totally resected may not require adjuvant RT, as the recurrence rate tends to be low. For





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

patients who have undergone maximum safe resection for low-grade intracranial ependymoma with no signs of dissemination on MRI and CSF analysis, adjuvant RT may be considered. RT is also an adjuvant treatment option for patients with myxopapillary ependymoma who had an STR or if capsule violation occurred, even if CSF cytology is negative. Craniospinal RT is recommended when MRI spine or CSF results reveal metastatic disease, regardless of histology and extent of resection.

### **Follow-up and Recurrence**

Follow-up of ependymoma depends on tumor grade and the location and extent of the disease. For localized disease, contrast-enhanced brain and spine MRI (if initially positive) should be done 2 to 3 weeks postoperatively and then every 3 to 4 months for one year. The interval can then be extended to every 4 to 6 months in the years 2 through 4, every 6 to 12 months for years 5 through 10, then as clinically indicated depending on the physician's concern regarding the extent of disease, histology, and other relevant factors. If tumor recurrence in the brain or spine is noted on one of these scans, restaging by brain and spine MRI as well as CSF analysis is necessary. More frequent MRI scans may also be indicated indefinitely for close follow-up in this setting. Resection is recommended if possible.

Upon disease progression or recurrence, treatment options depend on extent of disease, imaging and CSF findings, and prior treatment. For patients not previously irradiated, treatment with RT or consideration of SRS in appropriate cases for localized recurrence (negative MRI scan and CSF results), or craniospinal RT, when there is evidence of neuraxis metastasis, is recommended. For patients who have received prior RT treatment, clinical trials, chemotherapy, or palliative/best supportive care (in the setting of poor functional status) are the treatment options for those with evidence of recurrence with or without metastasis based on imaging and CSF findings. Patients who have received prior RT, are in good

functional status, and do not show evidence of neuraxis metastatic disease should be considered for enrollment in a clinical trial. Re-irradiation and chemotherapy may also be considered for these patients, as clinically appropriate.

### **Adult Medulloblastoma**

Though medulloblastoma is the most common brain tumor in children, it also can occur in adults,<sup>239</sup> though it makes up only 1% of CNS tumors in adults.<sup>240</sup> These tumors are often located in the cerebellar hemisphere<sup>241</sup> and can be broken into distinct molecular subtypes: WNT-activated, SHH-activated, and non-WNT/non-SHH (ie, Group 3 and Group 4).<sup>9,239,242</sup> Subtype analysis continues to evolve.<sup>243</sup> Adult medulloblastoma tends to be different genomically from pediatric medulloblastoma, including differing prognostic markers.<sup>244</sup> 6q loss is a prognostic marker in pediatric medulloblastoma, but not in adult medulloblastoma.<sup>245</sup> Tumors activated by SHH signaling are common in adult medulloblastoma.<sup>239,245,246</sup> Metastatic disease is less common in adult medulloblastoma than in children. It tends to occur in patients with non-WNT, non-SHH disease.<sup>247</sup> Tumors activated by WNT signaling are associated with good OS outcomes ( $P < .001$ ), based on a sample of patients with medulloblastoma that included children, infants, and adults, though trends were not statistically significant in analysis including only adults ( $n = 65$ ).<sup>239</sup> Somatic CTNNB1 mutations are very common in WNT-activated tumors; germline APC mutations occur in these tumors as well but are less common.<sup>248</sup> In patients with tumors activated by SHH signaling, prognosis is poor for those with tumors that are *TP53*-mut, compared to those with SHH-activated tumors that are not *TP53*-mut, even when controlling for histology, sex, presence of distantly metastatic disease, and age.<sup>249</sup> Therefore, the WHO further classifies SHH-mut medulloblastoma as *TP53*-mut and *TP53*-wt.<sup>9,250</sup>



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### Treatment Overview

#### ***Surgery***

Evidence in adult patients is meager for this rare disease and there are no randomized trial data, but there is general consensus that surgical resection should be the routine initial treatment to establish diagnosis, relieve symptoms, and maximize local control. Complete resection can be achieved in half of the patients<sup>251-253</sup> and is associated with improved survival.<sup>251,254</sup> When viewed by molecular subtype, near-total resection (<1.5 cm residual) and GTR produced equivalent OS for SHH, WNT, and Group 3 patients.<sup>255</sup> In addition, surgical placement of a ventriculoperitoneal shunt can be used to treat hydrocephalus.

#### ***Radiation Therapy***

Adjuvant RT following surgery is the current standard of care, although most studies are based on the pediatric population. The conventional dose is 30 to 36 Gy of craniospinal irradiation and a boost to a total of 54 to 55.8 Gy to the primary brain site.<sup>251,254</sup> Data from pediatric trials support use of a lower craniospinal dose of 23.4 Gy, combined with chemotherapy, while maintaining 54 to 55.8 Gy to the posterior fossa.<sup>256-258</sup> A randomized pediatric trial for standard-risk patients treated with radiation alone found an increased relapse risk with dose reduction.<sup>259</sup> A multicenter study including 70 adults with nonmetastatic medulloblastoma showed that reduced-dose craniospinal irradiation (23.4 or 35.2 Gy with a boost of 55.2 Gy to the fossa posterior) with maintenance chemotherapy is feasible.<sup>260</sup> It is reasonable to consider proton beam for craniospinal irradiation where available, as it is associated with less toxicity.<sup>261</sup> SRS demonstrated safety and efficacy in a small series of 12 adult patients with residual or recurrent disease.<sup>262</sup> Concomitant chemotherapy (vincristine) is typically omitted in adults given potential for severe toxicity.

#### ***Systemic Therapy***

The use of post-irradiation chemotherapy to allow RT dose reduction is becoming increasingly common especially for children,<sup>256,257</sup> but optimal use of adjuvant chemotherapy is still unclear for adult patients.<sup>253,263-266</sup>

Neoadjuvant therapy has not shown a benefit in pediatric or adult patients.<sup>267-269</sup> It is used in infants to defer radiation. A phase III study that enrolled more than 400 patients between 3 and 21 years of age with average-risk disease to receive post-irradiation cisplatin-based chemotherapy regimens recorded an encouraging 86% 5-year survival.<sup>258</sup>

In the setting of recurrence, several regimens are in use in the recurrence setting, most of which include etoposide.<sup>270-272</sup> Temozolomide has also been used in this setting.<sup>119,273</sup> High-dose chemotherapy in combination with autologous stem cell transplantation is a feasible strategy for patients who have had good response with conventional-dose chemotherapy, although long-term control is rarely achieved.<sup>272,274</sup> SHH-pathway inhibitors that have been evaluated in phase II trials including adults with recurrent medulloblastoma include vismodegib<sup>275</sup> and sonidegib.<sup>276</sup> Patients in these trials with SHH-activated disease were more likely to respond than patients with non-SHH disease.<sup>275,276</sup>

### NCCN Recommendations

#### ***Primary Treatment***

MRI scan is the gold standard in the assessment of medulloblastoma. The typical tumor shows enhancement and heterogeneity. Diffusion-weighted abnormalities are also characteristic of medulloblastoma. Fourth ventricular floor infiltration is a common finding related to worse prognosis.<sup>263,265,266</sup> Multidisciplinary consultation before treatment initiation is advised. Maximal safe resection is recommended wherever possible. Contrast-enhanced brain MRI should be performed within 48 hours following surgery, but spinal MRI should be delayed by 2 to 3 weeks. Because of the propensity of medulloblastoma to CSF seeding, CSF



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

sampling after spine imaging via lumbar puncture is also necessary for staging. Molecular profiling is recommended, as identification of clinically relevant medulloblastoma subtypes (eg, SHH-activated) may encourage opportunities for clinical trial enrollment. Medulloblastoma should be staged according to the modified Chang system using information from both imaging and surgery.<sup>277,278</sup>

### **Adjuvant Therapy**

Patients should be stratified according to recurrence risk for planning of adjuvant therapy (reviewed by Brandes et al<sup>279</sup>). The NCCN Panel agrees that patients with large cell or anaplastic medulloblastoma, disease dissemination, unresectable tumors, or residual tumors greater than 1.5 cm<sup>2</sup> postsurgery are at heightened risk. These patients should undergo irradiation of the neuraxis and chemotherapy. Collection of stem cells before RT may be considered on the condition that RT is not delayed for potential future autologous stem cell reinfusion at disease progression. For patients at average risk, craniospinal RT with or without chemotherapy or reduced-dose craniospinal RT with chemotherapy followed by post-irradiation chemotherapy are viable options.

### **Recurrence and Progression**

There are no robust data supporting an optimal follow-up schedule for medulloblastoma. Panel recommendations include brain MRI every 3 months for the first 2 years, every 6 to 12 months for 5 to 10 years, then every 1 to 2 years or as clinically indicated. If recurrent disease is detected on these scans, CSF sampling is also required, and concurrent spine imaging should be performed. Bone scans, contrast-enhanced CT scans of the chest, abdomen, and pelvis, and bone marrow biopsies may be considered as indicated.

Maximal safe resection should be attempted for recurrent medulloblastoma if symptomatic and there is no evidence of dissemination. Additional options include systemic therapy alone, RT

alone (including SRS), and chemoradiation. High-dose chemotherapy with autologous stem cell rescue may be considered for patients showing no evidence of disease after conventional reinduction chemotherapy. Patients with metastases should be managed by systemic therapy or best supportive care, which can include palliative RT. In very select cases, intrathecal chemotherapy might be utilized.

### **Primary CNS Lymphomas**

PCNSL accounts for approximately 3% of all neoplasms and 4% to 6% of all extranodal lymphomas.<sup>280</sup> It is an aggressive form of non-Hodgkin lymphoma that develops within the brain, spinal cord, eye, or leptomeninges without evidence of systemic involvement. The overall incidence of PCNSL in immunocompetent patients is 0.47 per 100,000 person-years, with higher incidence in males than in females and an increasing incidence with age.<sup>280</sup> The greatest increase in incidence has been reported in older adults with 1.8 per 100,000 patient-years reported in patients aged 65 years or older and 1.9 in patients aged 75 years or older, indicating that, in immunocompetent patients, PCNSL is a disease of older adults.<sup>280,281</sup> Non-immunosuppressed patients have a better prognosis than AIDS-related cases,<sup>282</sup> and survival of this group has improved over the years with treatment advances.<sup>283,284</sup> For more guidance on treatment of patients with PCNSL who are living with HIV, see the NCCN Guidelines for Cancer in People Living with HIV (available at [www.NCCN.org](http://www.NCCN.org)).

Pathologically, PCNSL is an angiocentric neoplasm composed of a dense monoclonal proliferation of lymphocytes, usually diffuse large B cells.<sup>285</sup> More than 90% of these primary CNS diffuse large B-cell lymphoma cases are of the activated B-cell–like (ABC) subtype.<sup>286</sup> The tumor is infiltrative and typically extends beyond the primary lesion, as shown by CT or MRI scans, into regions of the brain with an intact BBB.<sup>286</sup> The brain parenchyma is involved in more than 90% of all PCNSL patients, and the



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

condition can be multifocal in more than 50% of cases. Leptomeningeal involvement may occur, either localized to adjacent parenchymal sites or in diffuse form (that is, positive cytology) in up to 30% of patients. Ocular involvement may develop independently in 10% to 20% of patients. Patients with PCNSL can present with various symptoms because of the multifocal nature of the disease. In a retrospective review of 248 immunocompetent patients, 43% had mental status changes, 33% showed signs of elevated intracranial pressure, 14% had seizures, and 4% suffered visual symptoms at diagnosis.<sup>287</sup>

PCNSL occurs in about 7% to 15% of patients with post-transplant lymphoproliferative disorders (PTLD)<sup>288-291</sup> and is associated with poor prognosis.<sup>290,292,293</sup> PTLDs are a heterogeneous group of lymphoid neoplasms associated with immunosuppression following solid organ transplantation (SOT) or allogeneic hematopoietic stem cell transplantation (HSCT).<sup>294-296</sup> For guidance on managing transplant recipients, see the Post-Transplant Lymphoproliferative Disorders sub-algorithm in the NCCN Guidelines for Diffuse Large B-Cell Lymphoma (available at [www.NCCN.org](http://www.NCCN.org)).

### Treatment Overview

#### **Steroid Administration**

Steroids can rapidly alleviate signs and symptoms of PCNSL and improve PS. However, as these drugs are cytolytic, they can significantly decrease enhancement and size of tumors on CT and MRI scans as well as affect the histologic appearance. In the absence of significant mass effect, it is recommended that steroids be withheld or used judiciously until diagnostic tissue can be obtained if PCNSL is suspected.

#### **Stereotactic Biopsy**

In contrast to the principles previously outlined for invasive astrocytomas and other gliomas, the surgical goals for PCNSL are different, with the

main goal being establishment of diagnosis under minimal risk of morbidity. Currently, most authors recommend biopsy rather than resection.<sup>297</sup> This approach stems from the fact that data do not demonstrate a survival advantage for patients who have had a complete resection or extensive STR when compared with those who have had only a stereotactic biopsy. In addition, STR is associated with risk for postoperative neurologic deficits.<sup>287</sup>

#### **Systemic Therapy**

Methotrexate is the most effective agent against PCNSL. It is commonly used in combination with other drugs such as procarbazine, vincristine, cytarabine, rituximab, and temozolomide.<sup>298-311</sup> High doses of intravenous methotrexate are necessary (3.5 g/m<sup>2</sup> or higher) to overcome the BBB and achieve therapeutic levels in the CSF. Intrathecal methotrexate can be useful where CSF cytology yields positive findings and when patients cannot tolerate systemic methotrexate at 3.5 g/m<sup>2</sup> or higher. Other intrathecal chemotherapy options in this setting include cytarabine<sup>312</sup> and rituximab.<sup>313</sup> Phase II trials in the United States and Europe have shown that high-dose chemotherapy with autologous stem cell transplantation following high-dose methotrexate-based chemotherapy is feasible and well-tolerated, with little evidence of neurotoxicity.<sup>307,314-321</sup>

Renal dysfunction induced by high-dose methotrexate therapy is a potentially lethal medical emergency due to heightened toxicities resulting from a delay in methotrexate excretion. Early intervention with glucarpidase, a recombinant bacterial enzyme that provides an alternative route for methotrexate clearance, has shown efficacy in rapidly reducing plasma concentrations of methotrexate and preventing severe toxicity.<sup>322,323</sup>

It has become clear that consolidative therapy can result in significant and sometimes lethal neurotoxic effects from consolidation RT, especially in patients older than 60 years of age.<sup>301,324,325</sup> Complete response to





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

chemotherapy ranges from 42% to 61%, with OS ranging between 14 and 55 months. A number of phase II trials have adopted the approach of chemotherapy without planned RT.<sup>298,301,326-330</sup> However, a high fraction of patients who have forgone initial RT—typically older or with significant comorbidities—may fail to achieve complete response to chemotherapy. Studies investigating the efficacy of methotrexate-based regimens as induction therapy for patients with PCNSL have utilized WBRT, including reduced WBRT following cytarabine as consolidation treatment.<sup>300-302</sup>

There are currently no conclusive prospective data published comparing consolidation with high-dose chemotherapy regimens or high-dose chemotherapy with autologous stem cell transplantation versus maintenance therapy or observation, and there are different approaches at different institutions. Consolidation with high-dose chemotherapy and autologous stem cell transplant is frequently considered for fitter patients. Eligibility criteria used in the respective trials that studied these regimens need to be carefully considered when considering this approach, and referral to centers with subspecialty expertise in PCNSL should be considered.

Cytarabine combined with etoposide as high-dose consolidation therapy following induction treatment with methotrexate, temozolomide, and rituximab was evaluated in the multicenter Alliance 50202 trial.<sup>331</sup> This protocol was feasible and generally well-tolerated, with one treatment-related death.

High-dose chemotherapy with autologous stem cell transplantation in the relapsed/refractory setting has been tested with some success in two phase 2 European trials,<sup>332,333</sup> although evidence of its advantage over conventional treatment is lacking. The German Cooperative PCNSL Study Group evaluated the safety and efficacy of rituximab, high-dose cytarabine, and thiopeta followed by autologous stem cell transplantation in 39 patients with relapsed or refractory PCNSL with previous high-dose

methotrexate-based treatment.<sup>333</sup> A complete response was achieved in 56% of the patients. Out of the remaining patients, only one had progressive disease (18% of the patients had a partial response or stable disease). However, median OS was not reached, with a 2-year OS rate of 56.4%. Median PFS was 12.4 months, with a 2-year PFS rate of 46%. A phase 2 trial from France evaluated the efficacy of high-dose cytarabine and etoposide followed by autologous stem cell transplantation in 43 patients with relapsed or refractory PCNSL with previous high-dose methotrexate-based treatment.<sup>332</sup> Out of the 27 patients who completed autologous stem-cell rescue, median OS was 58.6 months (2-year OS was 69%) and median PFS was 41.1 months (2-year PFS was 58%).

High-dose chemotherapy and autologous stem cell transplantation as part of initial treatment has now been explored in several trials. High complete response rates and 2-year PFS have been demonstrated.<sup>307,334</sup> Whether high-dose chemotherapy and autologous stem cell rescue provides any additional benefit over consolidative conventional-dose chemotherapy or not is being investigated in two trials currently in progress. Consolidative conventional dose chemotherapy (NCTNA51101, MATRIX)<sup>335</sup> or consolidative WBRT (ANOCEF-GOELAMS, IELSG32)<sup>336</sup> have resulted in equivalent 2-year PFS in randomized phase II trials. Toxicities differ and might be a basis for individual patient selection. Of note, longitudinal neurocognitive assessment in the IELSG32 study showed persistent neurocognitive impairment in the consolidative WBRT group, but not in the high-dose chemotherapy group. The extent to which the patient selection inherent in high-dose chemotherapy trials underlies these favorable outcomes remains to be determined.

Unfortunately, even for patients who initially achieved complete response, half will eventually relapse. Re-treatment with high-dose methotrexate may produce a second response in patients who achieved complete response with prior exposure.<sup>337</sup> Rituximab as well as ibrutinib may be used in



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

combination with high-dose methotrexate retreatment.<sup>338</sup> Several other regimens, including ibrutinib,<sup>339,340</sup> rituximab,<sup>341</sup> TMZ with or without rituximab,<sup>342-345</sup> lenalidomide with or without rituximab,<sup>346</sup> high-dose cytarabine,<sup>347</sup> pomalidomide,<sup>348</sup> and pemetrexed<sup>349</sup> have also shown activity in the relapsed/refractory disease setting, but none has been established as a standard of care.

### **Radiation Therapy**

Historically, WBRT was the treatment standard to cover the multifocal nature of the disease. The majority of studies demonstrated the limitation of high-dose RT and led to the recommended dose of 24 to 36 Gy in 1.8 to 2.0 Gy fractions to the whole brain, without a boost.<sup>300,302,350-353</sup> Although RT alone is useful for initial tumor control, frequent and rapid relapse of the disease led to a short OS of 12 to 17 months.<sup>282,352</sup> This dismal outcome has prompted the addition of pre-irradiation methotrexate-based combination chemotherapy in later studies. This approach yields impressive response rates of up to 94% and improved OS ranging from 33 to 60 months.<sup>300-302,310,324,325,350,354,355</sup> However, excessive grade 3 and 4 hematologic toxicity (up to 78%) as well as RT-induced delayed neurotoxicity (up to 32%) sometimes leading to deaths are primary concerns, although most of these studies utilized a high RT dose of greater than or equal to 40 Gy. Of note, younger patients (aged <60 years) consistently fare better, and there is a higher incidence of late neurotoxic effects in older patients, but significant neurotoxicity can also occur in younger adults.

Thiel and colleagues<sup>356</sup> conducted a randomized, phase III, non-inferiority trial of high-dose methotrexate plus ifosfamide with or without WBRT in 318 patients with PCNSL. There was no difference in OS (HR, 1.06; 95% CI, 0.80–1.40;  $P = .71$ ), but the primary hypothesis (0.9 non-inferiority margin) was not proven. Patients who received WBRT had a higher rate of neurotoxicity than those who did not (49% vs. 26%). The panel currently

recommends that patients receiving WBRT because they are not candidates for chemotherapy should receive a dose of 24 to 36 Gy with a boost to gross disease, for a total dose of 45 Gy.

Although WBRT alone is seldom sufficient as primary treatment and is primarily reserved for patients who cannot tolerate multimodal treatment, it may be a reasonable treatment option for patients not suitable for other systemic therapies or clinical trials. Results from a phase II trial showed that reduced-dose WBRT (23.4 Gy in 1.8 Gy/fraction) following a complete response to induction chemotherapy was associated with disease response and long-term control, as well as low neurotoxicity.<sup>357</sup> When administered after chemotherapy failure, WBRT has shown response rates reaching nearly 75%.<sup>358</sup> Median PFS was 9.7 months overall, 57.6 months in patients who had achieved a CR with WBRT, and 9.7 months in patients with a PR. For patients who had a less than complete response to chemotherapy, a dosing schedule consistent with that used for induction treatment may be used, followed by a limited field to gross disease, or focal RT to residual disease.

### **NCCN Recommendations**

#### **Initial Evaluation**

Neuroradiologic evaluation is important in the diagnosis of PCNSL and to evaluate the effectiveness of subsequent therapy. With MRI, the tumor is often isointense or hypointense on T1- and T2-weighted images and enhances frequently.<sup>359</sup> In addition, restricted diffusion can be seen in the area of the enhancing abnormality on diffusion-weighted imaging sequences. On a CT scan, PCNSL is usually isodense or hyperdense compared to the brain and enhances in most cases. Hallmark features include a periventricular distribution, ring enhancement, multiple lesions, and a smaller amount of edema than might otherwise be expected from a similar-sized metastatic tumor or glioma. If contrast-enhanced brain MRI (or contrast-enhanced CT if MRI is contraindicated) suggests PCNSL,





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

clinicians are advised to hold the use of steroids if possible before diagnosis is established, since the imaging and histologic features of PCNSL can be profoundly affected by these drugs.

Patients with an enhancing brain lesion consistent with PCNSL should receive a biopsy (if lesion is amenable to biopsy), as this is the most direct and rapid route to achieve a pathologic diagnosis. Because the role of maximal surgical resection is limited to alleviating symptoms of raised intracranial pressure or preventing herniation,<sup>287</sup> stereotactic biopsy is generally preferred to minimize invasiveness.<sup>297</sup> Even with molecular marker testing, however, a biopsy can occasionally be falsely negative, particularly if the patient had been treated previously with steroids. Thus, if a biopsy is nondiagnostic, the panel recommends that the steroids be tapered and that the patient be followed closely, both clinically and radiographically. If and when the lesion recurs, there should be a prompt repeat CSF evaluation or rebiopsy before the initiation of steroids. If, on the other hand, no definitive diagnosis of lymphoma is made from biopsy and the patient has not received steroid therapy, workup for other diagnoses (for example, inflammatory processes) or repeat CSF evaluation/rebiopsy is recommended.

### ***Evaluation for Extent of Disease***

Once the diagnosis of PCNSL is established, the patient should undergo a thorough staging workup detailed by The International PCNSL Collaborative Group.<sup>297</sup> This workup involves a complete CNS evaluation including imaging of the entire neuraxis (MRI of the spine with contrast). If possible, this should be done before CSF analysis is attempted to avoid post-lumbar puncture artifacts that can be mistaken for leptomeningeal disease on imaging.

A lumbar puncture with evaluation of CSF (15–20 mL of spinal fluid) should be considered, if it can be done safely and without concern for herniation from increased intracranial pressure, and if it will not delay

diagnosis and treatment. A delay in treatment may compromise patient outcomes.<sup>331</sup> Caution should be taken in patients who are anticoagulated, thrombocytopenic, or who have a bulky intracranial mass. CSF analysis should include flow cytometric analysis, CSF cytology, and cell count. The yield for a positive diagnostic test can be increased by the use of molecular markers of monoclonality, such as an immunoglobulin gene rearrangement.

Since disease is sometimes detected in the retina and optic nerve, a full ophthalmologic exam should be done, which should include a slit-lamp eye examination. In some cases, the diagnosis of lymphoma is made by vitrectomy; in this case, flow cytometric analysis is recommended. In addition, blood work (CBC and chemistry panel) and a contrast-enhanced body CT or PET/CT<sup>360</sup> are required to rule out systemic involvement. Elevated lactate dehydrogenase (LDH) serum level is associated with worse survival in patients with PCNSL,<sup>361,362</sup> and LDH should be evaluated as part of the workup for this disease. Bone marrow biopsy is a category 2B option that may be considered. In men older than 60 years of age, testicular ultrasound may be considered (category 2B). In these patients, regular testicular examination is encouraged. If both testicular examination and CT or PET/CT imaging are negative, then testicular ultrasound may not be necessary.

An HIV blood test should also be performed, because both prognosis and treatment of patients with HIV-related PCNSL may be different than that of patients who are otherwise immunocompetent. HIV-positive patients should receive highly active retroviral therapy in addition to their cancer therapy.

### ***Newly Diagnosed Disease***

Induction treatment should be initiated as soon as possible following confirmation of diagnosis. The International PCNSL Collaborative Group has published treatment response criteria for complete response,



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

unconfirmed complete response, partial response, progressive disease, and relapsed disease.<sup>297</sup> Given the dramatic effect of steroids on symptom relief, they are commonly administered concurrently with workup. A high-dose methotrexate-containing regimen is the recommended induction treatment. In the case of methotrexate-induced renal dysfunction, consider glucarpidase to aid clearance. Non-methotrexate-based regimens may be used if the patient cannot tolerate methotrexate, usually those with impaired renal function.

If a patient is found to have malignant uveitis, orbital RT may be considered because of poor penetration of systemic chemotherapy into the uveal fluid. However, there are reports of clearance of ocular lymphoma in patients who were treated with systemic high-dose methotrexate.<sup>298</sup> Therefore, for a patient with PCNSL who has asymptomatic ocular involvement, a reasonable strategy is to delay RT to the globe in order to see if high-dose methotrexate is effective. Referral to a neuro-ophthalmologist or ophthalmologic oncologist for intraocular injection of chemotherapy (category 2B) is also an option.

WBRT may be used in patients who are not candidates for chemotherapy. For a patient treated with WBRT, consideration of intra-CSF chemotherapy plus focal spinal RT are treatment options if the lumbar puncture or spinal MRI are positive. Intrathecal chemotherapy options include methotrexate, cytarabine, and rituximab.

Treatment following induction high-dose methotrexate-based therapy depends on disease response.<sup>297</sup> Given the rarity of this disease, there are few high-quality studies to inform treatment decision-making. For patients who have a complete or unconfirmed complete response, consolidation therapy options that may be considered include high-dose chemotherapy (carmustine/thiotepa or thiotepa/busulfan/cyclophosphamide [TBC]) with stem cell rescue<sup>307,314-320</sup> or low-dose WBRT. However, WBRT in this setting may increase neurotoxicity,<sup>356,363</sup> especially in patients older than

60 years.<sup>301,324,325</sup> High-dose cytarabine with or without etoposide is also a consolidation treatment option for patients who had a complete response to induction high-dose methotrexate-based therapy (this regimen may also be considered in patients who do not have a complete response).<sup>300-302,331</sup> If there is not a complete or unconfirmed complete disease response following induction therapy, it is recommended to pursue another systemic therapy or WBRT in order to rapidly induce a response, diminish neurologic morbidity, and optimize quality of life. Best supportive care is another option for patients with residual disease following methotrexate-based treatment who are not candidates for other reasonable rescue therapies.

### ***Relapsed or Refractory Disease***

Patients should be followed using brain MRI. Imaging of the spine and CSF sampling may be done as clinically indicated for patients with spine disease. If there is ocular involvement, ophthalmologic exams may also be carried out.

For patients who are treated with prior WBRT and ultimately relapse, they may consider further chemotherapy (systemic and/or intrathecal), focal reirradiation, or palliative/best supportive care.

For patients who were initially treated with high-dose methotrexate-based chemotherapy but did not receive WBRT, the decision about whether to use other systemic therapy or proceed to RT at the time of relapse depends on the duration of response to initial chemotherapy. If a patient had experienced a relatively long-term response of one year or more, then treating either with the same (in most cases, high-dose methotrexate-based therapy) or another regimen is reasonable. However, for patients who either have no response or relapsed within a very short time after systemic therapy, recommendations include WBRT, switching to a different chemotherapy regimen, or involved-field RT with or without



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

chemotherapy.<sup>358</sup> In either case, palliative/best supportive care remains an option.

High-dose chemotherapy with stem cell rescue may also be considered as treatment for relapsed/refractory disease in patients who did not previously receive this treatment (ie, patients who were treated with high-dose methotrexate-based therapy or with WBRT) (category 2B). Regardless of primary treatment received, stem cell rescue should only be used for relapsed/refractory disease if there is a complete or partial response to re-induction high-dose chemotherapy.

For patients previously treated with high-dose chemotherapy with stem cell rescue, retreatment may be considered if there was a previous disease response and if time to relapse was at least one year. For patients who did not have a response to high-dose chemotherapy with stem cell rescue, and the time to relapse was less than one year, treatment options include RT to the whole brain or to the involved field. Regardless of time to relapse, using a different systemic therapy regimen (without stem cell rescue) and best supportive care are also options.

As there is no uniform standard of care for the treatment of refractory or relapsed PCNSL, participation in clinical trials is encouraged.

### Primary Spinal Cord Tumors

Spinal tumors are classified according to their anatomic location as extradural, intradural-extramedullary, and intradural-intramedullary. Extradural tumors are primarily due to metastatic disease and are discussed in the section *Metastatic Spinal Tumors*. This section focuses on intradural primary spinal tumors.

Primary spinal cord tumors are a histologically diverse set of diseases that represent 2% to 4% of all primary CNS tumors. The overall incidence is 0.74 per 100,000 person-years with a 10-year survival rate of 64%.<sup>364</sup>

Extramedullary lesions, most commonly benign meningiomas, account for 70% to 80% of spinal cord tumors.<sup>365</sup> Astrocytomas (more prevalent in children) and ependymomas (more prevalent in adults) are the most common intramedullary tumors. Clinicians are advised to refer to the corresponding sections in these guidelines for further details regarding these subtypes, as intracranial and spinal lesions are biologically similar.

Individuals with type I neurofibromatosis, type II neurofibromatosis, and von Hippel-Lindau syndrome are predisposed to form, respectively, spinal astrocytomas, spinal peripheral nerve sheath tumors, spinal ependymomas, and intramedullary hemangioblastomas.

Since 70% of primary spinal cord tumors are low-grade and slow-growing,<sup>364</sup> it is common for patients to suffer from pain for months to years before diagnosis. Pain that worsens at night is a classic symptom for intramedullary lesions. Progressive motor weakness occurs in half of the patients, and patients may experience sensory loss with late autonomic dysfunction (incontinence).

### Treatment Overview

#### Observation

Many asymptomatic primary tumors of the spinal cord, especially grade I meningiomas and peripheral nerve sheath tumors, follow an indolent course and can be followed by observation without immediate intervention.

#### Surgery

Surgery is the preferred primary treatment when the tumor is symptomatic and amenable to surgical resection. For lesions that are radiographically well defined, such as ependymoma, WHO grade I astrocytoma, hemangioblastoma, schwannoma, and WHO grade I meningioma, potentially curative, maximal, safe resection is the goal. En bloc total resection yielded excellent local control rates of more than 90%.<sup>366-369</sup>



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

GTR is seldom feasible with grade II or higher astrocytomas because they are infiltrative and poorly circumscribed. In a study of 202 patients with intramedullary tumors, over 80% of grade I astrocytomas were completely resected, while total resection was achieved in only 12% of grade II tumors.<sup>370</sup> Nevertheless, Benes et al<sup>371</sup> conducted a review of 38 studies on spinal astrocytomas and concluded that maximal safe resection should be attempted whenever possible based on reports of survival benefit.

### **Radiation Therapy**

RT is not recommended as the primary therapy without surgery and unknown histology because of the potential for limited response and low RT tolerance of the spinal cord. It is also not advisable following GTR of certain histologies, as select spinal cord tumors that can be excised completely have a low local recurrence rate.

A large retrospective analysis including more than 1700 patients with primary spinal gliomas found an association between RT and worse cause-specific survival and OS, although there may be a bias that patients who received RT had more adverse factors.<sup>372</sup> The role of adjuvant RT following incomplete excision or biopsy remains controversial.<sup>371,373,374</sup> One exception is primary spinal myxopapillary ependymoma, for which postoperative RT has been demonstrated to reduce the rate of tumor progression.<sup>375,376</sup> On the other hand, EBRT is considered a viable option at disease progression or recurrence. SRS has also shown safety and efficacy in several patient series.<sup>377-379</sup>

### **Systemic Therapy**

Unfortunately, evidence on efficacious chemotherapeutic agents for primary spinal cord tumors is too scant for specific recommendations. The panel agrees that chemotherapy should be an option where surgery and RT fail, but there is no consensus on the best regimen. Chemotherapy is best given in the setting of a clinical trial.

### **NCCN Recommendations**

MRI imaging is the gold standard for diagnosis of spinal cord lesions. However, CT myelogram may be used for diagnosis in patients for whom MRI is contraindicated. Asymptomatic patients may be observed (especially for suspected low-grade) or resected, while all symptomatic patients should undergo some form of surgery. The surgical plan and outcome are influenced by whether a clear surgical plane is available.<sup>380</sup> Whenever possible, maximal safe resection should be attempted, with a spine MRI 2 to 3 weeks following surgery to assess the extent of the resection. Postoperative adjuvant RT is appropriate if symptoms persist after incomplete resection or biopsy, or for patients with myxopapillary ependymoma that has been incompletely resected. Patients should be managed according to the pathology results (see *Low-Grade Pilocytic and Infiltrative Astrocytomas & Oligodendrogliomas, Anaplastic Gliomas and Glioblastomas, and Intracranial and Spinal Ependymomas*). Those diagnosed with hemangioblastoma should consider screening for von Hippel-Lindau syndrome including neuraxis imaging.<sup>381</sup>

All patients should be followed by sequential MRI scans, with a greater frequency in patients with high-grade tumors. At progression or recurrence, re-resection is the first choice. If this is not feasible, conventional EBRT or SRS is the next option. Chemotherapy is reserved for cases where both surgery and RT are contraindicated. Specific chemotherapy regimens are dependent on primary tumor type.

### **Meningiomas**

Meningiomas are extra-axial CNS tumors arising from the arachnoid cap cells in the meninges. They are most often discovered in middle-to-late adult life, and have a female predominance. The annual incidence for males and females reported by the Central Brain Tumor Registry of the United States (CBTRUS) are 1.8 and 3.4 per 100,000 people, respectively.<sup>382</sup> In a review of 319 cases using the WHO grading scale,





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

92% of meningiomas are grade I (benign), 6% are grade II (atypical), and 2% are grade III (malignant).<sup>383</sup> Small tumors are often asymptomatic, incidental findings.<sup>384</sup> Seizure is a common presenting symptom occurring in 27% of patients.<sup>385</sup>

### Imaging

Brain imaging with contrast-enhanced CT or MRI is the most common method of diagnosing, monitoring, and evaluating response to treatment (review by Campbell et al<sup>386</sup>). The CT scan best reveals the chronic effects of slowly growing mass lesions on bone remodeling. Calcification in the tumor (seen in 25%) and hyperostosis of the surrounding skull are features of an intracranial meningioma that can be easily identified on a non-contrast CT scan. Nonetheless, MRI reveals a number of imaging characteristics highly suggestive of meningioma, and in stereotactic RT articles, MR has been used to operationally define pathology. These MR findings include a tumor that is dural-based and isointense with gray matter, demonstrates prominent and homogeneous enhancement (>95%), has frequent CSF/vascular cleft(s), and often has an enhancing dural tail (60%). However, approximately 10% to 15% of meningiomas have an atypical MRI appearance mimicking metastases or malignant gliomas. In particular, secretory meningiomas may have a significant amount of peritumoral edema. Cerebral angiography is occasionally performed, often for surgical planning, as meningiomas are vascular tumors prone to intraoperative bleeding. In some instances preoperative embolization is helpful for operative hemostasis management. Angiographic findings consistent with a meningioma include a dual vascular supply with dural arteries supplying the central tumor and pial arteries supplying the tumor periphery. A “sunburst effect” may be seen due to enlarged and multiple dural arteries, and a prolonged vascular stain or so-called “blushing” can be seen, which results from intratumoral venous stasis and expanded intratumoral blood volume.

Meningiomas are also known to have high somatostatin receptor density, which allows for the use of octreotide brain scintigraphy to help delineate extent of disease and to pathologically define an extra-axial lesion.<sup>387-389</sup> Octreotide imaging with radiolabeled indium or, more recently, gallium may be particularly useful in distinguishing residual tumor from postoperative scarring in subtotally resected/recurrent tumors.

### Treatment Overview

#### Observation

Studies that examined the growth rate of incidental meningiomas in otherwise asymptomatic patients suggested that many asymptomatic meningiomas may be followed safely with serial brain imaging until either the tumor enlarges significantly or becomes symptomatic.<sup>390,391</sup> These studies confirm the tenet that many meningiomas grow very slowly and that a decision not to operate is justified in selected asymptomatic patients. As the growth rate is unpredictable in any individual, repeat brain imaging is mandatory to monitor an incidental asymptomatic meningioma.

#### Surgery

The treatment of meningiomas is dependent upon both patient-related factors (ie, age, PS, medical comorbidities) and treatment-related factors (ie, reasons for symptoms, resectability, goals of surgery). Most patients diagnosed with surgically accessible symptomatic meningioma undergo surgical resection to relieve neurologic symptoms. Complete surgical resection may be curative and is therefore the treatment of choice, if feasible. Both the tumor grade and the extent of resection impact the rate of recurrence. In a cohort of 581 patients, 10-year PFS was 75% following GTR but dropped to 39% for patients receiving STR.<sup>392</sup> Short-term recurrences reported for grade I, II, and III meningiomas were 1% to 16%, 20% to 41%, and 56% to 63%, respectively.<sup>393-395</sup> The Simpson classification scheme that evaluates meningioma surgery based on extent of resection of the tumor and its dural attachment (grades I–V in



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

decreasing degree of completeness) correlates with local recurrence rates.<sup>396</sup> First proposed in 1957, it is still being widely used by surgeons today.

### **Radiation Therapy**

Safe GTR is sometimes not feasible due to tumor location. In this case, STR followed by adjuvant EBRT has been shown to result in long-term survival comparable to GTR (86% vs. 88%, respectively), compared to only 51% with incomplete resection alone.<sup>397</sup> Of 92 patients with grade I tumors, Soyuer and colleagues found that RT following STR reduced progression compared to incomplete resection alone, but has no effect on OS.<sup>398</sup> Conformal fractionated RT (eg, 3D-CRT, IMRT, VMAT, proton therapy) may be used in patients with grade I meningiomas to spare critical structures and uninvolved tissue.<sup>399</sup>

Because high-grade meningiomas have a significant probability of recurrence even following GTR,<sup>400</sup> postoperative high-dose EBRT (>54 Gy) has become the accepted standard of care for these tumors to improve local control.<sup>401</sup> Initial results of the phase II RTOG 0539 trial showed that patients with high-risk meningioma treated with IMRT (60 Gy in 30 fractions) had a 3-year PFS rate of 58.8%.<sup>402</sup> High risk was defined as new or recurrent grade III, recurrent grade II, or new grade II with STR. Since new and recurrent tumors were grouped together, this study does not provide clarification on the appropriate role of RT following GTR in patients with newly diagnosed WHO grade II disease, and the role of post-GTR RT in these cases remains controversial.

Technical advances have enabled stereotactic administration of RT by linear accelerator (LINAC), Leksell Gamma Knife™, or CyberKnife™ radiosurgery. The use of stereotactic RT (either single fraction or fractionated) in the management of meningiomas continues to evolve. Advocates have suggested this therapy in lieu of EBRT for small (<35 mm), recurrent, or partially resected tumors. In addition, it has been used

as primary therapy in surgically inaccessible tumors (ie, base-of-skull meningiomas) or in patients deemed poor surgical candidates because of advanced age or medical comorbidities. Nonrandomized and retrospective studies show that stereotactic RT is associated with excellent tumor control and good survival outcomes, particularly in grade I tumors, indicating that this treatment is effective as primary and second-line treatment for meningiomas smaller than 3.5 cm.<sup>403-407</sup> However, optimal dosing has not been determined.

### **Systemic Therapy**

For meningiomas that recur despite surgery and/or RT, or are not amenable to treatment with surgery or RT, systemic therapies are often considered. Due to the rarity of these patients requiring systemic therapy, large randomized trials are lacking. Historical estimates of 6-month PFS rates in these patients range from 0% to 29%.<sup>408</sup> Smaller studies support the use of targeted therapy including somatostatin analogues in select cases.<sup>409,410</sup> Recently, studies investigating anti-angiogenic therapies in meningioma have demonstrated improved results.

A prospective, multicenter, nonrandomized, phase II trial evaluating the safety and efficacy of sunitinib in 36 heavily pretreated patients with refractory meningioma showed a 6-month PFS rate of 42%, with a median PFS rate of 5.2 months and a median OS rate of 24.6 months.<sup>411</sup> However, toxicities were considerable, with 60% of patients experiencing grade 3 or higher toxicity.

Retrospective data support the use of bevacizumab for patients with recurrent meningioma, especially for patients with symptoms driven by RT necrosis, with a 6-month PFS rate of 43.8% for recurrent surgery and radiation-refractory grade II and III meningioma with bevacizumab monotherapy.<sup>412,413</sup> In a phase II trial evaluating the efficacy and safety of bevacizumab combined with everolimus for recurrent meningioma ( $N = 17$ ), stable disease was reported in 88% of patients, with no complete or





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

partial responses reported.<sup>414</sup> The median PFS and OS rates were 22.0 months and 23.8 months, respectively, with 18-month PFS and OS rates of 57% and 69%, respectively. Treatment was discontinued in 22% of patients due to toxicity.

### NCCN Recommendations

#### *Initial Treatment*

Meningiomas are typically diagnosed by brain MRI. Surgery or octreotide scan may be considered for confirmation. For treatment planning, multidisciplinary panel consultation is encouraged. Patients are stratified by the presence or absence of symptoms and the tumor size. Most asymptomatic patients with small tumors ( $\leq 3$  cm) are best managed by observation; otherwise, patients should undergo surgical resection whenever possible. Non-surgical candidates should undergo RT.

Regardless of tumor size and symptom status, all patients with surgically resected grade III meningioma (even after GTR) should receive adjuvant RT to enhance local control. For patients with grade II meningioma, postoperative RT is recommended for incomplete resection; in the case of complete resection in these patients, postoperative RT may be considered, although this treatment strategy remains controversial. Patients with grade I meningioma may be observed following surgery, though postoperative RT may be considered in patients with symptomatic disease. SRS may be used in lieu of conventional RT as adjuvant or primary therapy in asymptomatic cases.

#### *Follow-up and Recurrence*

In the absence of data, panelists have varying opinions on the best surveillance scheme and clinicians should follow patients based on individual clinical conditions. Generally, malignant or recurrent meningiomas are followed more closely than grade I and II tumors. A typical schedule for low-grade tumors is MRI every 3 months in year 1,

then every 6 to 12 months for another 5 years. After 5 years, imaging may be done every 1 to 3 years as clinically indicated.

Upon detection of recurrence, the lesion should be resected whenever possible, followed by RT. Non-surgical candidates should receive RT. Chemotherapy is reserved for patients with an unresectable recurrence refractory to RT. Observation is an option if there is no clinical indication for treatment at recurrence.

### Brain Metastases

Metastases to the brain are the most common intracranial tumors in adults and may occur up to 10 times more frequently than primary brain tumors. Population-based data reported that about 8% to 10% of patients with cancer are affected by symptomatic metastatic tumors in the brain.<sup>415,416</sup> Based on autopsy studies, brain metastases have been shown to be present in 25% of patients with cancer.<sup>417</sup>

As a result of advances in diagnosis and treatment, many patients improve with proper management and do not die of progression of these metastatic lesions. Primary lung cancers are the most common source,<sup>418</sup> although melanoma has a high predilection to spread to the brain.<sup>419</sup> Diagnosis of CNS involvement is increasing in patients with breast cancer as therapy for metastatic disease is improving.<sup>420</sup>

Nearly 80% of brain metastases occur in the cerebral hemispheres, an additional 15% occur in the cerebellum, and 5% occur in the brainstem.<sup>421</sup> These lesions typically follow a pattern of hematogenous spread to the gray-white junction where the relatively narrow caliber of the blood vessels tends to trap tumor emboli. The majority of cases have multiple brain metastases evident on MRI scans. The presenting signs and symptoms of metastatic brain lesions are similar to those of other mass lesions in the brain, such as headache, seizures, and neurologic impairment.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### Treatment Overview

#### ***Surgery***

Despite advances in surgical technique, surgery alone for brain metastases is not sufficient for achieving local control.<sup>422,423</sup> The objectives of surgery for brain metastasis include retrieval of tissue for diagnosis, reduction of mass effect, and improvement of edema.<sup>424</sup> To promote local control following resection of a brain metastasis, adjuvant RT represents an acceptable treatment strategy, discussed further below. Randomized trials reported in the 1990s demonstrated an OS benefit with surgical resection for patients with single brain metastases. In a study of 48 patients, Patchell et al<sup>425</sup> demonstrated that surgery followed by WBRT compared with WBRT alone improved OS (40 vs. 15 weeks in WBRT arm;  $P < .01$ ) and functional dependence (38 vs. 8 weeks;  $P < .005$ ), as well as decreased recurrence (20% vs. 52%;  $P < .02$ ). Similarly, combined surgery and WBRT led to longer survival and functional independence compared to WBRT alone in another randomized study by Vecht and colleagues ( $n = 63$ ).<sup>426</sup> A third study of 84 patients found no difference in survival between the two strategies; however, patients with extensive systemic disease and lower performance level were included, which likely resulted in poorer outcomes in the surgical arm.<sup>427</sup>

#### ***Stereotactic Radiosurgery***

SRS offers an excellent minimally invasive ablative treatment option for brain metastases. Patients undergoing SRS avoid the risk of surgery-related morbidity, and SRS is generally preferred over surgery for patients with small, asymptomatic lesions that do not require surgery and for patients with lesions that are not surgically accessible.<sup>424</sup> Late side effects of SRS such as symptomatic edema and RT necrosis are relatively uncommon, but may be observed at higher rates when treating larger lesions.<sup>428</sup>

The role of stereotactic SRS alone for limited brain metastases has been established by multiple phase III randomized trials comparing SRS alone to SRS plus WBRT.<sup>429-432</sup> Collectively, these studies demonstrate comparable OS and superior cognitive preservation and quality of life with SRS alone compared to SRS plus WBRT. The role of SRS for patients with multiple metastases has also continued to expand. A prospective trial of 1194 patients found no differences in OS or neurologic mortality with SRS for 2 to 4 versus 5 to 10 brain metastases.<sup>433</sup> A number of analyses have suggested that total volume of brain metastases and the rate of developing new brain metastases may be more important prognostic factors for OS than the number of discrete brain metastases.<sup>434-437</sup> Taken together, patients with multiple lesions but a low total volume of disease, as well as those with relatively indolent rates of developing new CNS lesions, can represent suitable candidates for SRS. Additionally, patients with a favorable histology of the primary tumor (such as breast cancer) or controlled primary tumors can often benefit from SRS regardless of the number of brain metastases present.<sup>438,439</sup> While brain metastases arising from small cell lung cancer have historically been treated with WBRT, an international retrospective study suggested that SRS may be suitable in some cases.<sup>440</sup> Some brain metastases of radio-resistant primary tumors such as melanoma and renal cell carcinoma have also been shown to achieve good local control with SRS.<sup>441</sup> Other predictors of longer survival with SRS include younger age, good PS, and primary tumor control.<sup>434,438,439,442</sup> However, there are a number of contemporary series supporting SRS in patients with a poor prognosis, with poor KPS, or who are older.<sup>443-446</sup>

Maximal marginal doses for SRS use should be based on tumor volume and range from 15 to 24 Gy when treating lesions with a single fraction of SRS.<sup>429,433,447</sup> Multi-fraction SRS may be considered for larger tumors, with the most common doses being 27 Gy in 3 fractions and 30 Gy in 5 fractions.<sup>448-450</sup> Contouring guidelines have been published elsewhere.<sup>451</sup>



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

In the recurrence setting, several patient series have demonstrated local control rates greater than 70% with SRS for patients with good PS and stable disease who have received prior WBRT.<sup>452-455</sup>

Postoperative SRS also represents an important strategy to improve local control after resection of brain metastases. After resection alone, the rates of local recurrence are relatively high, and have been reported in the range of 50% at 1 to 2 years in prospective trials. Postoperative SRS to the surgical cavity is supported by a randomized phase III trial including 132 patients with resected brain metastases (1–3 lesions). This trial demonstrated that postoperative SRS was associated with a higher 12-month local recurrence-free rate compared to no postoperative treatment (72% vs. 43%, respectively; HR, 0.46; 95% CI, 0.24–0.88;  $P = .015$ ).<sup>422</sup> A separate randomized phase III trial comparing postoperative SRS with postoperative WBRT demonstrated similar OS and better cognitive preservation with a strategy of postoperative SRS, despite superior CNS control outcomes with WBRT.<sup>456</sup>

### **Whole-Brain Radiation Therapy**

Historically, WBRT was the mainstay of treatment for metastatic lesions in the brain. Although the role of WBRT has diminished over the last several decades, WBRT continues to play a role in the modern era, primarily in clinical scenarios where SRS and surgery are not feasible or indicated (eg, diffuse brain metastases). The standard dosing for WBRT is 30 Gy in 10 fractions or 37.5 Gy in 15 fractions. For patients with poor prognosis, 20 Gy in 5 fractions may also be used.

The impact of WBRT in addition to SRS has been evaluated in multiple randomized controlled studies.<sup>429-432,457</sup> A 2018 Cochrane meta-analysis of randomized controlled trials found that the addition of WBRT to SRS alone was associated with better brain control, no differences in OS, and worse neurocognitive outcomes or quality of life in several trials.<sup>458</sup> The randomized phase III EORTC 22952 trial failed to show an OS benefit

from WBRT following resection or SRS, compared to observation,<sup>432</sup> even in subgroup analyses including only patients with controlled extracranial disease and a favorable prognostic score.<sup>459</sup> Overall, for patients treated with SRS for brain metastases, the routine addition of WBRT is not recommended due to increased cognitive and quality-of-life toxicity and the lack of an OS benefit.

The randomized phase III non-inferiority QUARTZ trial compared WBRT to optimal supportive care in patients with non-small cell lung cancer (NSCLC) who were not candidates for SRS, due to various factors including age, PS, and extent of disease. No differences in OS or quality of life were observed with WBRT versus optimal supportive care, which suggests that this population may derive minimal benefit from WBRT.<sup>460</sup> Moreover, as noted above, a number of studies support SRS for older patients and those with poor prognosis who have historically received WBRT.<sup>443-446,461</sup> The optimal treatment strategy of brain metastases for patients with a poor prognosis is highly individualized and may call for best supportive care, WBRT, SRS, or trials of CNS-active systemic agents depending on the clinical scenarios.

In light of the well-characterized deleterious cognitive effects of WBRT,<sup>430,431,456</sup> a number of trials have evaluated strategies to promote cognitive preservation in patients with brain metastases including investigation of neuroprotective agents, anatomical avoidance strategies, and deferral of WBRT in favor of alternate strategies such as SRS or trials of CNS-active systemic agents. In patients undergoing WBRT for brain metastases, the RTOG 0614 ( $N = 554$ ) compared concurrent and adjuvant memantine, an N-methyl-D-aspartate receptor antagonist, to placebo. Memantine was well-tolerated in patients receiving WBRT for brain metastases, and the rates of toxicity were similar to patients receiving placebo.<sup>462</sup> There was possibly less decline in episodic memory (HVL-T-R Delayed Recall) in the memantine arm compared to placebo at 24 weeks



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

( $P = .059$ ). The memantine arm had significantly longer time to cognitive decline (HR, 0.78; 95% CI, 0.62–0.99;  $P = .01$ ), and the probability of cognitive function failure at 24 weeks was 54% in the memantine arm and 65% in the placebo arm. However, for most cognitive endpoints, no significant differences were observed between memantine and placebo, despite numerical trends that generally favored the memantine arm. For patients with a good prognosis, memantine may be considered during WBRT, as well as after treatment for as long as 6 months.

To evaluate an anatomic-avoidance strategy to promote cognitive preservation, the nonrandomized phase II RTOG-0933 trial showed that reduced radiation dose to the hippocampal neural stem-cell compartment was associated with a smaller decline in recall ( $P < .001$ ), compared to a historical control.<sup>463</sup> Based on these results, the phase III NRG-CC001 trial evaluated WBRT with memantine with or without hippocampal avoidance.<sup>464</sup> There were no significant differences in survival outcomes. However, risk of cognitive failure was significantly lower in the hippocampal avoidance arm than in the control arm (HR, 0.76; 95% CI, 0.60–0.98;  $P = .03$ ). For patients without tumor in or around the hippocampus, hippocampal-sparing WBRT may be preferred in select patients (eg, those with good prognosis).

In the postoperative setting, phase 3 trials have evaluated the role of WBRT after surgical resection of brain metastases. Patchell conducted a study that randomized 95 patients with single intracranial metastases to surgery with or without adjuvant WBRT.<sup>465</sup> Postoperative RT was associated with a dramatic reduction in tumor recurrence (18% vs. 70%;  $P < .001$ ) and likelihood of neurologic deaths (14% vs. 44%;  $P = .003$ ). OS, a secondary endpoint, showed no difference between the arms. The aforementioned EORTC 22952 trial randomized patients treated with local therapy (surgery or SRS) to observation versus WBRT.<sup>432</sup> Patients randomized to WBRT were found to have superior brain disease control

and less death from neurological causes, but inferior quality of life and no differences in OS.<sup>432,466</sup> The NCCTG N107C/CEC-3 randomized phase III trial included 194 patients with resected brain metastases randomized to either postoperative SRS or WBRT.<sup>456</sup> Although there was no significant difference between the treatment arms for OS, cognitive deterioration at 6 months was less frequent in the SRS arm than in the WBRT arm (52% vs. 85%, respectively;  $P < .001$ ), and cognitive deterioration-free survival was also greater for postoperative SRS compared to WBRT (median 3.7 months vs. median 3.0 months; HR, 0.47; 95% CI, 0.35–0.63;  $P < .001$ ). In another phase III trial, 215 patients with 1 to 3 brain metastases from melanoma were randomized to either WBRT or observation following local treatment with surgery or SRS.<sup>467</sup> Though local failure rate was significantly lower in the WBRT arm (20.0% vs. 33.6%, respectively;  $P = .03$ ), there were no significant differences between the study arms for intracranial failure, OS, and deterioration in performance status. Further, grade 1 to 2 toxicity during the first 2 to 4 months was more frequently reported in the WBRT arm.

### Systemic Therapy

Many tumors that metastasize to the brain are not chemosensitive or have already been heavily pretreated with organ-specific effective agents. Poor penetration through the BBB is an additional concern.<sup>419</sup> However, there are increasing numbers of systemic treatment options with demonstrated activity in the brain, and it is now reasonable to treat some of these patients (ie, those with asymptomatic brain metastases) with systemic therapy upfront instead of upfront SRS or WBRT.

Specific recommended regimens for brain metastases are based on effective treatment of the primary tumor (see below). However, there is also an increasing number of “basket” studies that evaluate the efficacy of targeted therapy options for a specific mutation or biomarker, regardless of tumor type. For example, the TRK inhibitors larotrectinib and entrectinib





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

were found to be active in patients with brain metastases from *NTRK* gene fusion-positive solid tumors.<sup>468,469</sup>

As CNS-active systemic agents are changing paradigms for the management of brain metastases, it is important to acknowledge that there is a paucity of prospective data to characterize optimal strategies regarding radiation and systemic therapy combinations or sequencing. When considering a trial of upfront systemic therapy alone for brain metastases, a multidisciplinary discussion between medical and radiation oncology is recommended. Ongoing CNS surveillance with brain MRIs is essential to allow early interventions in cases of progression or inadequate response.

### *Melanoma*

Rapid advancements in melanoma have produced effective systemic options for metastatic disease.<sup>470,471</sup> These include multiple immunotherapy options. Two phase II trials support the use of a combination of the immunotherapy agents ipilimumab and nivolumab for patients with asymptomatic untreated brain metastases from melanoma.<sup>472,473</sup> In one of these trials, which was conducted in Australia, intracranial responses were observed in 46% of patients who received this combination, with a complete response observed in 17% ( $n = 79$ ), and median duration of response was not reached at the time of publication (median 14 months of follow-up).<sup>472</sup> In the second trial, CheckMate 204, the intracranial response was 57%, with a complete response of 26% ( $N = 94$ ), with median duration of intracranial response also not having been reached at time of publication (median 14 months of follow-up).<sup>473</sup> In both of these trials, grade 3 or 4 treatment-related adverse events occurred in just over half of the patients evaluated.<sup>472,473</sup> Results from the Australian trial also suggest there may be a role for nivolumab monotherapy for patients with asymptomatic untreated brain metastases ( $n = 27$ ), with an intracranial response rate of 20%.<sup>472</sup> For patients with asymptomatic

untreated lesions, the response rate for patients who received ipilimumab/nivolumab was better than for nivolumab monotherapy. This trial also evaluated nivolumab monotherapy for a small number of patients for whom local therapy failed ( $n = 16$ ), but the intracranial response rate was low (6%). A nonrandomized phase II study supports ipilimumab monotherapy for patients with small asymptomatic brain metastases from melanoma ( $n = 51$ ), with a CNS disease control rate of 24% (no complete responses).<sup>474</sup> Most of the patients in this study had received previous systemic or local treatment. Nivolumab monotherapy is a reasonable treatment option for a carefully monitored patient whose goal is to avoid radiation.

The anti-PD-1 antibody pembrolizumab is also supported for treatment of both untreated and progressive brain metastases from melanoma, based on early results of a phase II trial showing a CNS ORR of 22% ( $n = 18$ ).<sup>475</sup> Long-term follow-up from this trial showed a CNS response in 26% of the sample ( $N = 23$ ), with four complete responses.<sup>476</sup> In patients who had a CNS response, these responses were ongoing at 24 months in all of the patients. Median PFS and OS were 2 months and 17 months, respectively. Grade 3–4 treatment-related adverse events were minimal. Despite data showing that brain metastases can respond to immune checkpoint inhibitors, the data do not yet provide any robust comparison of these agents from treatment of brain metastases from melanoma.

There is also evidence that brain metastases from melanoma can respond to BRAF/MEK inhibitor combination therapy. The nonrandomized phase II COMBI-MB trial demonstrated clinical benefit and acceptable toxicity for the combination of the BRAF inhibitor dabrafenib with the MEK inhibitor trametinib in 125 patients with brain metastases from *BRAF* V600-mut melanoma.<sup>477</sup> Among the patients with asymptomatic brain metastases, an intracranial response was observed in 58% of those with untreated metastases and in 56% of those with previously treated metastases. In



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

patients with symptomatic brain metastases, an intracranial response was observed in 59%. Use of the BRAF inhibitor vemurafenib for patients with both newly diagnosed and previously treated brain metastases from *BRAF* V600-mut melanoma is supported by nonrandomized studies.<sup>478,479</sup>

Although there are no published prospective studies on the combination of vemurafenib and cobimetinib for patients with brain metastases from melanoma, there is high-quality evidence that, for distantly metastatic melanoma, combination therapy with vemurafenib and cobimetinib is associated with improved outcomes, compared with vemurafenib monotherapy.<sup>480,481</sup> A case series showed that the BRAF/MEK inhibitor combination encorafenib/binimetinib showed good CNS penetration.<sup>482</sup> Prospective randomized trials are needed to determine which *BRAF*-directed therapy options provide the best results in patients with brain metastases from melanoma.

### *Lung Cancer*

Systemic treatment options for patients with brain metastases from NSCLC include immunotherapy agents and targeted therapies for cancer that is anaplastic lymphoma kinase (ALK) rearrangement-positive and EGFR mutation-positive.

### **PD-1/PD-L1 Inhibitors**

A phase II trial showed a 33% response rate for pembrolizumab in 18 patients with brain metastases from PD-L1-positive NSCLC.<sup>475</sup> Pooled analyses from a phase II trial<sup>483</sup> and two phase III trials<sup>484,485</sup> showed that nivolumab for patients with previously treated brain metastases from NSCLC is well-tolerated, though results from these analyses are currently only reported in abstract form.<sup>486</sup> Nivolumab for patients with brain metastases from NSCLC is also supported by results from a retrospective multi-institutional study.<sup>487</sup>

### **ALK Inhibitors**

At time of diagnosis, brain metastases are present in 24% of patients with ALK rearrangement-positive NSCLC.<sup>488</sup> Crizotinib inhibits *ALK* rearrangements, *ROS1* rearrangements, and some MET TKIs. Crizotinib does demonstrate some CNS activity,<sup>489</sup> but the response and control rates appear to be clearly lower than newer generation ALK inhibitors.

In a randomized phase III trial, the ALK inhibitor alectinib was compared to crizotinib in 303 patients with advanced ALK rearrangement-positive NSCLC and no previous systemic therapy treatment.<sup>490</sup> Brain metastases were reported in 40.3% of the sample. Among these patients, a CNS response was observed in 81% of patients in the alectinib arm (8 complete responses) and 50% of patients in the crizotinib arm (1 complete response). The median duration of intracranial response in these 122 patients was 17.3 months in the alectinib arm and 5.5 months in the crizotinib arm. Pooled analyses from two phase II studies<sup>491,492</sup> including patients with ALK rearrangement-positive NSCLC that progressed on crizotinib showed that alectinib was associated with a good objective response rate and excellent disease control in patients with brain metastases.<sup>493</sup> Patients who did not receive previous brain RT seemed to have a better response to alectinib than patients with previous RT, but the sample size for these analyses was small.

In a similar randomized phase III trial, brigatinib, another ALK inhibitor, was compared to crizotinib in 275 patients with locally advanced or metastatic ALK rearrangement-positive NSCLC and no previous systemic therapy treatment.<sup>494</sup> Among patients with brain metastases ( $n = 90$ ), an intracranial response was more likely in the brigatinib arm than in the crizotinib arm (67% vs. 17%, respectively; OR, 13.00; 95% CI, 4.38–38.61). Complete intracranial responses were observed in 16 patients who received brigatinib and 2 patients who received crizotinib. Twelve-month survival without intracranial disease progression was greater in the brigatinib arm than in the crizotinib arm (67% vs. 21%, respectively; HR,





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

0.27; 95% CI, 0.13–0.54). Brigatinib treatment in patients with brain metastases from ALK rearrangement-positive NSCLC and disease progression on crizotinib is supported by the phase II ALTA trial, which showed an intracranial response rate of 67%.<sup>495</sup> Median intracranial PFS was 12.8 months in these patients. A dosing schedule of 180 mg once daily with a 7-day lead-in at 90 mg was used to reduce the chance of early-onset moderate to severe pulmonary adverse events.

The ALK inhibitor ceritinib was evaluated in a phase I trial including 246 patients with ALK rearrangement-positive NSCLC.<sup>496</sup> About half the sample had brain metastases ( $n = 124$ ). Retrospective analyses were used to evaluate intracranial response in these patients. Disease control rate was 78.9% in patients not previously treated with an ALK inhibitor and 65.3% in patients with previous ALK inhibitor treatment. However, most of these patients had received RT to the brain. Therefore, it is difficult to draw conclusions regarding the contribution of RT versus ceritinib to disease control rates in these patients.

In general, the panel prefers second- and third-generation ALK inhibitors for patients with brain metastases from ALK rearrangement-positive NSCLC, based on better activity profiles. A phase II trial in which the third-generation ALK/ROS1 TKI lorlatinib was evaluated in patients with ALK-positive NSCLC previously treated with an ALK TKI showed that this agent may be active against CNS metastases.<sup>497</sup>

### Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors

Some treatment options for patients with advanced NSCLC that harbor EGFR-TKI-sensitizing mutations have been evaluated and are now available.

Older-generation EGFR-TKIs have demonstrated some CNS activity. Gefitinib for treatment of patients with CNS metastases from NSCLC is supported by phase II studies.<sup>498,499</sup> Pulsatile erlotinib is supported by a

phase I study including patients with untreated CNS metastases from EGFR-sensitizing mutation-positive NSCLC.<sup>500</sup> Afatinib treatment was evaluated in patients with CNS metastasis from NSCLC and with disease progression following platinum-based chemotherapy and either erlotinib or gefitinib ( $n = 100$ ).<sup>501</sup> Cerebral response was observed in 35% of these patients, and disease control was observed in 66%.

In a randomized phase III FLAURA trial, the EGFR-TKI osimertinib was compared to a different EGFR-TKI (gefitinib or erlotinib) in 556 patients with previously untreated EGFR-sensitizing mutation-positive NSCLC.<sup>502</sup> CNS metastases were reported in 20.9% of the sample. Median PFS was greater for these patients in the osimertinib arm than in the standard EGFR-TKI arm (15.2 months vs. 9.6 months, respectively; HR, 0.47; 95% CI, 0.30–0.74;  $P < .001$ ). Preplanned exploratory analyses including 41 patients with at least one measureable CNS lesion showed a CNS ORR of 91% in the osimertinib arm, compared to 68% in the EGFR-TKI arm, but this difference did not reach statistical significance (OR, 4.6; 95% CI, 0.9–34.9;  $P = .066$ ).<sup>503</sup> Twenty-three percent of patients in the osimertinib arm had a complete CNS response, compared to none of the patients in the EGFR-TKI arm. CNS disease control rate did not significantly differ between the study arms in patients with at least one measureable CNS lesion.

Osimertinib has also been evaluated in the randomized phase III AURA3 trial, in which it was compared to pemetrexed with platinum-based therapy in 419 patients with *T790M* mutation-positive advanced NSCLC that progressed after first-line EGFR-TKI therapy.<sup>504</sup> CNS metastases were reported in 34.4% of the sample. Median PFS was greater for these patients in the osimertinib arm than in the pemetrexed/platinum arm (8.5 months vs. 4.2 months, respectively; HR, 0.32; 95% CI, 0.21–0.49). Preplanned analyses including 46 patients with at least one measureable CNS lesion showed a significantly greater CNS ORR for the osimertinib



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

arm than in the pemetrexed/platinum arm (70% vs. 31%, respectively; OR, 5.13; 95% CI, 1.44–20.64;  $P = .015$ ).<sup>505</sup> CNS disease control rate was 93% in the osimertinib arm, compared to 63% in the pemetrexed/platinum arm. Median CNS duration of response was also longer in the patients who received osimertinib. Pooled analyses from two phase II studies<sup>506,507</sup> including patients with *T790M*-positive advanced NSCLC that progressed following treatment with EGFR-TKI therapy showed a CNS ORR of 54% and disease control rate of 92%.<sup>508</sup> Median CNS duration of response and median PFS were not reached.

### Other Systemic Therapy Options

A phase I/II study of topotecan plus WBRT has shown a 72% response rate in 75 patients with brain metastases.<sup>509</sup> Unfortunately, a follow-up phase III trial including only patients with brain metastases from lung cancer was closed early due to slow accrual.<sup>510</sup>

### Breast Cancer

Capecitabine combined with a number of agents has been evaluated in patients with brain metastases from HER2-positive breast cancer. Capecitabine combined with the TKI lapatinib for patients with brain metastases from HER2-positive breast cancer is supported by a systematic review and pooled analysis showing an ORR of 29.2%, a disease control rate of 65.1%, and a 2-year OS rate of 33.4%.<sup>511</sup> A phase II study supports use of capecitabine combined with the TKI neratinib in patients with CNS metastases from HER2-positive breast cancer.<sup>512</sup> CNS metastases in most of the patients were previously treated with surgery or RT. Results from this study helped inform development of the phase III NALA trial, in which patients with HER2-positive metastatic breast cancer who received at least 2 lines of HER2-directed therapy were randomized to receive capecitabine and neratinib or capecitabine and lapatinib ( $N = 621$ ).<sup>513</sup> Patients in the capecitabine/neratinib arm had superior PFS compared to those in the capecitabine/lapatinib arm (HR, 0.76; 95% CI,

0.63–0.93;  $P = .006$ ), though there was no OS advantage. Further, patients who received capecitabine/neratinib were less likely to have required intervention for symptomatic CNS metastases than patients in the capecitabine/lapatinib arm (22.8% vs. 29.2%, respectively;  $P = .043$ ). In the HER2CLIMB phase III trial, patients with HER2-positive metastatic breast cancer who were previously treated with HER2-directed therapy ( $N = 612$ ) were randomized to receive trastuzumab and capecitabine combined with either the TKI tucatinib or a placebo.<sup>514</sup> Among the patients with brain metastases at baseline (47.5% of the sample), both PFS (HR, 0.46; 95% CI, 0.31–0.67) and OS (HR, 0.58; 95% CI, 0.40–0.85) were superior in the tucatinib arm. The estimated 1-year PFS was 24.9% for these patients who received tucatinib, compared to 0% in patients who received the placebo, with duration of PFS being 7.6 months and 5.4 months, respectively. Based on these study results, the FDA approved tucatinib in combination with trastuzumab and capecitabine in 2020 for patients with advanced unresectable or metastatic HER2-positive breast cancer (including patients with brain metastases) who were previously treated with HER2-directed therapy. Capecitabine monotherapy treatment in patients with brain metastases from breast cancer is supported by a phase I trial<sup>515</sup> and case reports.<sup>516-519</sup>

In a randomized phase II trial evaluating paclitaxel combined with neratinib, compared to trastuzumab combined with paclitaxel, in patients with untreated metastatic HER2-positive breast cancer, incidence of symptomatic or progressive CNS events were significantly lower in the neratinib arm (8.3% vs. 17.3%, respectively; HR, 0.48; 95% CI, 0.29–0.79;  $P = .002$ ).<sup>520</sup> Though patients with asymptomatic CNS metastases at baseline were eligible to participate in this trial, they comprised only 3.8% of the study sample, limiting the conclusions that can be drawn about the efficacy of this regimen for these patients.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

A study of high-dose methotrexate in patients mostly with breast cancer achieved disease control in 56% of patients.<sup>521</sup> The use of cisplatin and etoposide monotherapies and combination therapy in patients with brain metastases from breast cancer is supported by nonrandomized studies published in the 1990s.<sup>522-524</sup>

### NCCN Recommendations

#### **Workup**

Patients who present with a single mass or multiple lesions on MRI or CT imaging suggestive of metastatic cancer to the brain, and who do not have a known primary, require a careful systemic workup with chest x-ray or CT with contrast, abdominal or pelvic CT with contrast, or other tests as indicated. Whole-body PET/CT may be considered. If no other readily accessible tumor is available for biopsy, a stereotactic or open biopsy resection is indicated to establish a diagnosis.

#### **Treatment for Limited Metastatic Lesions**

The panel defines “limited” brain metastases as patients for whom SRS represents an effective alternative to WBRT, but with more cognitive protection.<sup>433</sup> Because brain metastases are often managed by physicians from multiple disciplines, the NCCN Panel encourages multidisciplinary consultation prior to treatment for optimal planning.

Surgical resection may be considered in select cases (eg, for management of mass effect or other symptoms; for tumors >3 cm that are surgically accessible; if there is no other readily accessible tumor to be biopsied). For patients with newly diagnosed or stable systemic disease, treatment options include SRS (preferred) and WBRT. When patients are managed with SRS, NCCN does not recommend the routine addition of WBRT, as this approach has been consistently associated with cognitive deterioration and no difference in survival.<sup>430</sup> The management of patients with disseminated systemic disease or poor prognosis should be

individualized and may include strategies of best supportive care, WBRT, SRS, or a trial of CNS-active systemic agents; multidisciplinary evaluation is encouraged.

In patients with systemic cancers with options for CNS-active systemic therapies, (eg, ALK or EGFR mutations in NSCLC; *BRAF* mutations in metastatic melanoma), upfront systemic therapy alone may be considered in carefully selected, asymptomatic patients. When considering a trial of upfront systemic therapy alone for brain metastases, NCCN recommends a multidisciplinary discussion between medical and radiation oncologists and ongoing CNS surveillance with brain MRIs to allow for early interventions in cases of progression or inadequate response.

Patients should be followed with brain MRI every 2 to 3 months for 1 to 2 years and then every 4 to 6 months indefinitely. Closer follow-up every 2 months may be particularly helpful for patients treated with SRS or systemic therapy alone.<sup>431</sup> Evaluation of potential disease recurrence can be confounded by treatment effects of SRS. Tumor sampling may be indicated to discern recurrence versus treatment effect in some cases. Upon detection of recurrent disease, prior therapy clearly influences the choice of further therapies. Patients with recurrent CNS disease should be assessed for local versus systemic disease, because therapy will differ. For local recurrences, patients who were previously treated with surgery only can receive the following options: 1) surgery with consideration of SRS or RT to the surgical bed; 2) single-dose or fractionated SRS; 3) WBRT; or 4) systemic therapy. However, patients who previously received WBRT generally should not undergo WBRT at recurrence due to concern regarding neurotoxicity. If the patient had previous SRS with a durable response for >6 months, reconsider SRS if imaging or biopsy supports active tumor and not necrosis. Repeat SRS to a prior location is a category 2B recommendation.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

If isolated CNS disease progression occurs in the setting of limited systemic treatment options and poor PS, management of brain metastases should be individualized and may include best supportive care, WBRT, SRS, and CNS-active systemic agents. WBRT re-irradiation is generally discouraged due to toxicity to cognition and quality of life and should be administered only in highly selected circumstances.

### ***Treatment for Extensive Metastatic Lesions***

Patients diagnosed with extensive metastatic lesions should generally be treated with WBRT or SRS as primary therapy. For WBRT dosing, the standard regimens are 30 Gy in 10 fractions or 37.5 Gy in 15 fractions. For patients with poor neurologic performance, a more rapid course of RT can be considered (20 Gy, delivered in 5 fractions). SRS may be considered in select patients, particularly those with good PS and low overall tumor volume. Some patients may be eligible for upfront systemic therapy treatment. Palliative neurosurgery may also be considered if a lesion is causing a life-threatening mass effect, hemorrhage, or hydrocephalus.

After WBRT or SRS, patients should have a repeat contrast-enhanced MRI scan every 2 to 3 months for 1 to 2 years, then every 4 to 6 months indefinitely. Treatment for recurrences are individualized and may include best supportive care, surgery, WBRT, SRS, or a trial of CNS-active systemic therapy; multidisciplinary review is recommended. Repeat WBRT is generally discouraged due to toxicity to cognition and quality of life and should only be administered in highly selected circumstances.

### **Leptomeningeal Metastases**

Leptomeningeal metastasis or neoplastic meningitis refers to the multifocal seeding of the leptomeninges by malignant cells. It is known as leptomeningeal carcinomatosis or carcinomatous meningitis when these cells originate from a solid tumor. When it is related to a systemic lymphoma, it is called lymphomatous meningitis, and when associated

with leukemia, it is termed leukemic meningitis. Leptomeningeal metastasis occurs in approximately 5% of patients with cancer.<sup>525</sup> This disorder is being diagnosed with increasing frequency as patients with cancer live longer with improved systemic therapeutics and as neuroimaging studies improve. Most cases arise from breast cancers, lung cancers, and melanoma, which has the highest rate of leptomeningeal spread.<sup>526,527</sup>

Tumor cells gain access to the leptomeninges by hematogenous dissemination, lymphatic spread, or direct extension. Once these cells reach the CSF, they are disseminated throughout the neuraxis by the constant flow of CSF. Infiltration of the leptomeninges by any malignancy is a serious complication that results in substantial morbidity and mortality. Cranial nerve palsies, headaches, focal deficits from cortical disturbances, mental changes, and motor weakness are among the most common presenting symptoms.<sup>525</sup> The median survival of patients diagnosed with this disorder is typically <3 months with death resulting from progressive neurologic dysfunction, but survival may be extended by early detection and intervention.<sup>526,527</sup>

### **Treatment Overview**

The goals of treatment in patients with leptomeningeal metastases are to improve or stabilize the patient's neurologic symptoms and to prolong survival.<sup>528</sup> Unfortunately, there is a lack of standard treatments due to meager evidence in literature. Because treatment is largely palliative, aggressive chemotherapy should only be given to patients most likely to benefit (see *Patient Stratification*).

### ***Radiation Therapy***

RT is mainly given for symptom alleviation, CSF flow correction, or for debulking to facilitate chemotherapy.<sup>527,529,530</sup> SRS may be an option for





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

patients with focal leptomeningeal disease, particularly in the setting of focal disease causing CSF flow disruption.<sup>531</sup>

### ***Surgery***

The role of neurosurgery for leptomeningeal metastases is mainly limited to intraventricular catheter and subcutaneous reservoir placement for drug administration.<sup>532</sup> This is preferred over lumbar punctures because of improved drug delivery, safety, superior pharmacokinetics, lower inter-patient variability, and patient comfort.<sup>533</sup>

### ***Systemic Therapy***

Some systemically administered agents are able to reach the leptomeninges, while others do not traverse the blood CSF barrier. Intrathecal (intra-CSF) chemotherapy can be used to address particularly non-bulky leptomeningeal disease, although it is important to note that it is an effective treatment for brain parenchymal disease. Some drugs have good CNS penetration, particularly organ-specific targeted therapies or systemically administered chemotherapies given in high doses.<sup>528</sup> Intrathecal therapy can involve either administration via a lumbar puncture or intraventricular injections via an Ommaya reservoir. However, both intra-CSF therapy and high-dose systemic therapy are associated with significant toxicity or complications and are therefore generally restricted to patients with good PS.

Agents used for intra-CSF therapy are often histology-specific and, because they are directly injected into the CSF, have good drug bioavailability. The panel included intrathecal options deemed appropriate based on moderate benefit: methotrexate<sup>534-536</sup>; cytarabine<sup>535,537,538</sup>; thiotepe<sup>536,539</sup>; rituximab for lymphoma<sup>540</sup>; topotecan<sup>541</sup>; etoposide<sup>542</sup>; and trastuzumab for HER2-positive breast cancer.<sup>543</sup> Interferon alfa was removed as an intra-CSF chemotherapy option in 2020 due to discontinuation.

Breast cancers<sup>521,544</sup> and lymphomas<sup>537,545</sup> are also particularly responsive to high-dose methotrexate. In addition, osimertinib and weekly pulse erlotinib have been used for metastatic NSCLC with EGFR sensitizing mutations [exon 19 deletion or exon 21 L858R mutation only for erlotinib (category 2B)].<sup>546-549</sup>

## **NCCN Recommendations**

### ***Patient Evaluation***

Patients present with signs and symptoms ranging from injury to nerves that traverse the subarachnoid space, direct tumor invasion of the brain or spinal cord, alteration of the local blood supply, obstruction of normal CSF flow pathways leading to increased intracranial pressure, or interference with normal brain function. Patients should have a physical examination with a careful neurologic evaluation. MRI of the brain and spine should also be performed for accurate staging, particularly if the patient is a candidate for active treatment. A definitive diagnosis is most commonly made by CSF analysis via lumbar puncture if it is safe for the patient. The CSF protein is typically increased, and there may be a pleocytosis or decreased glucose levels and ultimately positive CSF cytology for tumor cells. Assessment of circulating tumor cells increases the sensitivity of tumor cell detection in CSF.<sup>550-552</sup> This assessment is now CLIA-approved in some states and should be done when it is available. CSF cytology testing has approximately 50% sensitivity with the first lumbar puncture, and up to 90% sensitivity after repeated CSF analyses in affected patients.<sup>530</sup> Clinicians should be aware that lumbar punctures may be contraindicated in patients with anticoagulation, thrombocytopenia, or bulky intracranial disease. In these cases, suspicious CSF biochemical results combined with suggestive clinical and/or radiologic features should be taken into consideration. Although a positive CSF cytology in patients with solid tumors is virtually always diagnostic, reactive lymphocytes from infections (for example, herpes zoster infection) can often be mistaken for malignant lymphocytes.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### **Patient Stratification**

Once the diagnosis has been established, the patient's overall status should be carefully assessed to determine how aggressively the carcinomatous or lymphomatous meningitis should be treated. Unfortunately, this disease is most common in patients with advanced, treatment-refractory systemic malignancies for whom treatment options are limited. In general, fixed neurologic deficits (such as cranial nerve palsies or paraplegia) do not resolve with therapy, although encephalopathies may improve dramatically. As a result, patients should be stratified into "poor-risk" and "good-risk" groups. The poor-risk group includes patients with KPS below 60; multiple, serious, major neurologic deficits; extensive systemic disease with few treatment options; bulky CNS disease; and neoplastic meningitis related to encephalopathy. The good-risk group includes patients with KPS greater than or equal to 60, no major neurologic deficits, minimal systemic disease, and reasonable systemic treatment options. Many patients fall in between these two groups, and clinical judgment will dictate how aggressive their treatment should be.

### **Treatment**

Patients in the poor-risk group are usually offered palliative/supportive care measures, though patients considered good-risk may also receive palliative/best supportive care if they do not desire further treatment. Fractionated EBRT to neurologically symptomatic sites (eg, to the whole brain for increased intracranial pressure or to the lumbosacral spine for a developing cauda equina syndrome) can be considered to temporarily improve function.

Chemotherapy (systemic or intrathecal) is recommended for patients considered good-risk. These patients may also receive SRS, WBRT, or involved-field RT to neurologically symptomatic or painful sites and to areas of bulky disease identified on neuroimaging studies. Craniospinal RT may also be considered, but only in highly select patients given the

substantial toxicity and resultant bone marrow suppression that can limit future cancer-directed therapies.

CSF flow abnormalities are common in patients with neoplastic meningitis, and these often lead to increased intracranial pressure. Administering chemotherapy into the ventricle of a patient with a ventricular outlet obstruction increases the patient's risk for leukoencephalopathy. In addition, the agent administered may not reach the lumbar subarachnoid space where the original CSF cytology was positive if there are flow obstructions. Therefore, a CSF flow scan should be carried out if there are concerns about a CSF flow blockage (eg, a patient with hydrocephalus) prior to administration of intrathecal systemic therapy. If significant flow abnormalities are seen, fractionated EBRT can be administered to the sites of obstruction before repeating a CSF flow scan. High-dose systemically administered methotrexate remains an option for patients with breast cancer or lymphoma, as normal CSF flow is not required to reach cytotoxic concentrations.

The patient should be reassessed clinically and with a repeat CSF cytology. Cytology should be sampled from the lumbar spine, if possible, or via an intraventricular port. Neuraxis imaging with MRI is recommended if CSF cytology was initially negative and can also be considered for sites that were previously positive on a radiograph.

If negative cytology is achieved after induction, continue the induction chemotherapy for another month before switching to maintenance intrathecal chemotherapy. The CSF cytology status should be followed every 4 to 8 weeks. If the patient is clinically stable or improving after induction and there is no clinical or radiologic evidence of progressive leptomeningeal disease, the patient should receive another 4 weeks of "induction" intrathecal chemotherapy or should consider switching intrathecal drugs for 4 weeks. This regimen should be followed by maintenance therapy and monthly cytology if the cytology has converted





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

to negative or is improving (still positive) while the patient is clinically stable.

### **Progressive Disease**

If the patient's clinical status is deteriorating from progressive leptomeningeal disease or if the cytology is persistently positive, the clinician has several options: 1) RT to symptomatic sites; 2) systemic chemotherapy; or 3) palliative or best supportive care.

### **Metastatic Spinal Tumors**

Bone metastases are a growing problem among patients with cancer due to increasing life expectancy, with the spine being the most frequently affected site. Spinal metastases primarily arise from breast, lung, prostate, and renal cancers.<sup>553,554</sup> Extradural lesions account for about 95% of spinal tumors, mostly in the thoracic region.

Some patients are found to have vertebral involvement as an asymptomatic, incidental finding. However, for most affected patients, pain is the primary presenting symptom preceding neurologic dysfunction. Three types of pain have been classically defined. Local pain due to tumor growth is often described as a constant, deep aching that improves with steroid medications. Mechanical back pain varies with movement and position and is attributed to structural spinal instability. While seldom responsive to steroids, mechanical pain can be alleviated by surgical stabilization. Radicular pain is a sharp or stabbing sensation that occurs when nerve roots are compressed by the tumor. Patients may experience any one or a combination of these types of pain.

Spinal cord compression is the most debilitating complication of spine metastases. It affects 5% to 10% of all patients with cancer, with more than 20,000 cases diagnosed each year in the United States.<sup>555</sup> The majority of patients initially complain of progressive radicular pain.<sup>556</sup> This is followed by neurologic symptoms such as motor weakness and sensory

loss, and may even include autonomic bladder dysfunction. If left untreated, neurologic deficits rapidly progress to paralysis. Unfortunately, a study of 319 patients with cord compression revealed significant delay in the report of initial pain (3 months) as well as diagnosis (2 months) that can lead to irreversible spinal cord damage.<sup>557</sup> Therefore, it is paramount that the clinician watches for early suspicious signs and establishes prompt diagnosis by spine MRI. Once diagnosed, spinal cord compression is considered a medical emergency; intervention should be implemented immediately to prevent further neurologic decline.

### **Treatment Overview**

Dissemination to the spinal column is largely incurable. Therefore, the goals of treatment are palliation and improvement of quality of life through preservation of neurologic function, pain relief, and stabilization of mechanical structure. One exception is slow-growing cancers (mainly renal cell carcinoma) with solitary spinal metastasis, for which surgery may achieve possible cure.<sup>558</sup> Patients with spine metastases require care from a multidisciplinary team, including neurosurgeons; orthopedic surgeons; radiologists and interventional radiologists; and specialists in pain management; care of the bowel, bladder, and back; and ambulatory support.

The type and aggressiveness of the primary tumor often dictates the choice of treatment, as different cancers have varying sensitivities to systemic therapy and RT. In addition, patient characteristics including PS and comorbidities will determine whether they can tolerate surgery and, if so, which surgical technique should be used.

### **Surgery**

There is general consensus that a patient should have a life expectancy of at least 3 months to be a surgical candidate. Paraplegia for over 24 hours is a strong relative contraindication due to low chances of improvement



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

when prolonged neurologic deficits exist before surgery.<sup>559</sup> Patients with hematologic malignancies should also be excluded, as they are best managed by RT or chemotherapy. Because estimation of life expectancy can be difficult, several groups have developed prognostic scoring systems to help predict surgical outcomes.<sup>560-563</sup>

Modern surgical techniques enable surgeons to achieve 360° decompression of the spinal cord, and stabilization can be performed concomitantly, if required. The development of a plethora of spinal implants composed of high-quality materials such as titanium greatly improves reconstruction outcome. The surgical approach—anterior, posterior, or combined/circumferential—is primarily determined by disease anatomy.<sup>564,565</sup>

Sundaresan and colleagues<sup>558</sup> reported favorable results using a variety of surgical approaches on 80 patients with solitary spine metastases. Both pain and mobility were improved in the majority of patients. OS reached 30 months, with 18% of patients surviving 5 years or more. The best outcome was observed in patients with kidney and breast cancers.

Surgery followed by adjuvant EBRT has emerged as a highly effective approach in relieving spinal cord compression and restoring function, especially for solid tumors. A meta-analysis including 24 surgery cohort studies and four RT studies found that patients are twice as likely to regain ambulatory function after surgery than RT alone.<sup>566</sup> However, data also revealed significant surgery-related mortality (6.3%) and morbidity (23%). In another review of literature from 1964 to 2005, anterior decompression with stabilization plus RT was associated with superior outcome over RT alone or laminectomy, achieving 75% mean improvement in neurologic function. However, high surgical mortality rate (mean 10%) was also reported.<sup>567</sup>

To date, only one relevant randomized trial has been reported.<sup>568</sup> Approximately 100 patients with metastatic spinal compression were randomized to surgery plus postoperative RT or RT alone. Compared to the RT group, significantly more patients in the surgery group regained walking ability (84% vs. 57%;  $P = .001$ ) and for a longer period of time (median 122 days vs. 13 days;  $P = .003$ ). The impressive results were obtained with strict eligibility criteria. The study excluded patients with radiosensitive tumors, neurologic deficits for 24 hours, multiple spinal tumors, lesions only compressing spinal roots, and prior RT to the vertebrae. Although studies demonstrated high efficacy of surgery, the formidable complications related to surgery cannot be overlooked. Using the Nationwide Inpatient Sample all-payer database, Patil et al<sup>569</sup> reviewed data of more than 26,000 patients who had undergone surgery for spinal metastases. The in-hospital mortality and complication rates were 5.6% and 22%, respectively. The most common complications were pulmonary (6.7%) and hemorrhages or hematomas (5.9%). Clearly, careful individual patient selection based on life expectancy and overall health is warranted.

### **Radiation Therapy**

Traditionally, EBRT has been the main form of treatment for spinal metastases. In the modern surgery era, RT alone is often not sufficient in achieving decompression or stabilization (see above), but it is routinely used as adjuvant therapy following surgery as it is difficult to obtain wide negative margins. Given the potential impact of RT on wound healing, most studies posed an interval of 1 to 3 weeks between resection and subsequent RT.<sup>570</sup>

An excellent response to RT alone for spinal compression was reported by Marazano and colleagues.<sup>571</sup> Three hundred patients were randomized to a short-course (8 Gy x 2 days) or split-course (5 Gy x 3 days; 3 Gy x 5 days) schedule. After RT, 35% of nonambulatory patients regained walking ability, and pain relief was recorded in 57% of patients with a



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

median survival of 4 months. Efficacy of RT was highly dependent on the histology: 70% of patients with nonambulatory breast cancer recovered mobility compared to only 20% of hepatocellular carcinoma patients. In general, solid tumors are considered either moderately radiosensitive (eg, breast and prostate cancers) or radioresistant (eg, melanoma; osteosarcomas; cancers of the thyroid, colon, and kidney).<sup>572</sup> On the other hand, hematologic malignancies such as lymphomas and multiple myelomas are highly responsive to RT. Hence, RT alone is routinely utilized as therapy for these cancers, even in the presence of cord compression.

Where there is no compression, fracture, or instability, EBRT is effective in achieving local control as primary treatment. A mean 77% local control rate from seven retrospective studies including 885 patients was found in a systematic review by Gerszten and colleagues.<sup>572</sup> RT is also a mainstay of palliative treatment for patients with poor PS, significant comorbidities, and/or limited life expectancy (<3–4 months). Klimo's meta-analysis, including 543 patients treated by RT, revealed pain control rates of 54% to 83%.<sup>566</sup> Unlike surgery, RT has no immediate significant treatment-related complications and very few local recurrences. However, it increases surgical complications as it impairs wound healing.

Stereotactic radiation approaches (SRS or stereotactic body RT [SBRT]) allow precise high-dose targeting in one or two fractions while minimizing exposure of the surrounding cord.<sup>573</sup> This is especially important in pre-irradiated patients. A review including 59 publications with 5655 patients who received SRS for spinal metastases showed 1-year local control rates of 80% to 90% for newly diagnosed disease, 80% following surgery, and 65% for previously irradiated disease.<sup>574</sup> Single-institution reports suggest that SRS is safe and offers more durable response than conventional therapy,<sup>572,575,576</sup> and results of the phase II RTOG 0631 trial demonstrated the feasibility of SRS for these patients.<sup>577</sup> The phase III component of this

trial is ongoing and is comparing single-dose stereotactic RT of 16 Gy to single-dose EBRT of 8 Gy in patients with one to three spinal metastases (NCT00922974). Consensus guidelines should be followed for stereotactic radiation planning and delivery.<sup>573,578,579</sup> Reasonable dosing schedules for the postoperative setting have been published by Redmond et al.<sup>579</sup>

### **Vertebral Augmentation**

Percutaneous vertebroplasty and kyphoplasty involve injection of cement (polymethyl methacrylate) into the vertebral body. Vertebroplasty is a direct injection, while kyphoplasty involves inserting a balloon that provides a cavity for the injection. These vertebral augmentation procedures immediately reinforce and stabilize the column, thereby relieving pain and preventing further fractures.<sup>580</sup> They are suitable in poor surgical candidates with painful fractures, but are relatively contraindicated in the case of spinal cord compression because they do not achieve decompression. Symptomatic complications occur in up to 8% of patients (mostly with vertebroplasty), including embolization of the cement and local metastasis along the needle tract.

### **Systemic Therapy**

Corticosteroids remain a routine initial prescription for patients presenting with cord compression, with a number of theoretical benefits including anti-inflammation, reduction in edema, short-term neurologic function improvement, and enhanced blood flow. However, the preference between high-dose (96 mg daily) and low-dose (10–16 mg daily) is still unclear.<sup>581-583</sup>

Chemotherapy has a limited role in metastatic spinal tumors except for chemosensitive tumors such as lymphoma, myeloma, and germ cell tumors. Agents efficacious for the primary tumor are used.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

### NCCN Recommendations

#### Workup

Initial workup depends on the presence or absence of symptoms. Patients with an incidental, asymptomatic, metastatic lesion confirmed by systemic imaging can be observed with MRI. However, biopsy and further treatment of an incidental lesion are indicated if treatment of the patient is altered as a result of treatment of the incidental lesion. In the absence of symptoms, it is not mandatory to obtain a spinal MRI for every incidental metastatic lesion seen on surveillance bone scans. The alternate category involves severe or new back pain. Increasing intensity, duration, and changes in the character of pain should trigger an evaluation with an MRI study, even in patients with pre-existing degenerative spine conditions. Immediate spinal MRI is warranted in the occurrence of neurologic symptoms including weakness, paresthesias, and bladder or bowel incontinence. Contrast can be used to highlight and further evaluate any focal abnormality. The MRI can be used to image the entire spine or a focal area of interest. If the patient is unable to have an MRI, then a CT myelogram is recommended.

A normal neurologic examination implies that there is no spinal radiculopathy or myelopathy correlating with the patient's symptoms. In this case, other causes should be considered (eg, leptomeningeal disease). An abnormal neurologic examination includes motor abnormalities, sphincter abnormalities, and/or sensory deficits attributable to a dysfunction of nerve root(s) and/or the spinal cord. Therefore, detection of radiculopathy, myelopathy, or cauda equina syndrome is indicative of an abnormal examination. However, reflex asymmetry and/or presence of pathologic reflexes, as well as sensory deficits of a stocking/glove distribution are excluded.

#### Treatment

Once metastatic vertebral involvement is diagnosed, treatment is based on whether the patient is suffering from spinal cord compression, fracture, or spinal instability. In the presence of multiple metastatic spinal tumors, the one causing the patient's main symptoms is addressed first. Additional tumors can be treated at a later point according to the algorithm.

Radiographic spinal cord compression implies deformation of the spinal cord because of epidural tumor, retropulsed bone fragment, or both. It should be noted that epidural tumor may occupy part of the spinal canal with or without partial obliteration of CSF around the spinal cord. Those cases are excluded because there is no cord deformation. For tumors occurring below L1, any canal compression of 50% or more should be considered of equal importance as spinal cord compression. Patients with radiographic cord compression should start on dexamethasone (10–100 mg) to alleviate symptoms. Decompressive surgery (concomitant stabilization if indicated) and adjuvant RT is the preferred treatment (category 1) where there is spinal instability and no surgical contraindication. Primary EBRT alone is appropriate for patients with radiosensitive cancers (hematologic malignancies) and without evidence of spinal instability. Many fractionation schemes are available (8 Gy in 1 fraction, 20 Gy in 5 fractions, or 30 Gy in 10 fractions); the most common is a total of 30 Gy in 3-Gy daily fractions for 10 days.<sup>584,585</sup> Tolerance at the spinal cord and/or nerve root must be considered in determining dose. Primary chemotherapy is also an option for chemo-responsive tumors in the absence of clinical myelopathy. In general, a treatment interval of at least 6 months is recommended.

Metastases to the spine without cord compression include the presence of tumor in the vertebral body, pedicle(s), lamina, transverse, or spinous process. It can also include epidural disease without cord deformation. Patients in this category should be assessed for fractures and spinal





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

instability. Because the criteria for spinal destabilization secondary to tumor remain unclear, consultation by a surgeon is recommended. *Spinal instability* is grossly defined as the presence of significant kyphosis or subluxation (deformity) or of significantly retropulsed bone fragment. Not every pathologic fracture implies unstable structure. The degree of kyphosis or subluxation compatible with instability depends on the location of the tumor in the spine. The cross-sectional area of the vertebral body unaffected by the tumor and the patient's bone mineral density are additional factors affecting stability. In addition, vertebral body involvement is more important than dorsal element involvement with regard to stability. Circumferential disease as well as junctional and contiguous tumor location should be taken into account when assessing spinal stability. If fracture or instability is detected, the patient should undergo surgical stabilization or minimally invasive vertebral augmentation to relieve pain. These procedures should be followed by adjuvant RT to obtain local control.

If no fracture or instability is found, EBRT is the treatment of choice. Stereotactic RT may be appropriate in select cases of limited disease. Other alternatives are chemotherapy for responsive tumors, or surgery plus adjuvant RT in select cases. Patients experiencing intractable pain or rapid neurologic decline during RT should consider surgery, or stereotactic RT is an option if oligometastases are present or if the disease is radioresistant. Neurologic deterioration is apparent when the patient's neurologic examination is becoming worse on a daily basis and the patient's ambulatory status is threatened. Intractable pain means that pain is not controlled with oral analgesics or that the patient cannot tolerate the medication due to side effects.

### **Progression and Recurrence**

Follow-up involves MRI or CT imaging within 1 to 3 months post-treatment, then every 3 to 4 months for 1 year, then as clinically indicated.

Upon detection of progression or recurrence on imaging scans, management strategy is based on previous treatment. Patients who underwent prior RT or surgery plus adjuvant RT may consider surgery or re-irradiation to the recurred area. Stereotactic RT may be appropriate for select patients. Clinicians should plan 6 months or more between treatments in consideration of tolerance of the spine and its nerve roots. Retreatment dose should be limited to no more than 10 Gy to the surface of the spinal cord. In patients who were previously treated with chemotherapy, surgery may be indicated depending upon the degree of spinal stability/cord compression. RT may also be considered.





### References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin* 2020;70:7-30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31912902>.
2. Jukich PJ, McCarthy BJ, Surawicz TS, et al. Trends in incidence of primary brain tumors in the United States, 1985-1994. *Neuro Oncol* 2001;3:141-151. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11465394>.
3. Deorah S, Lynch CF, Sibenaller ZA, Ryken TC. Trends in brain cancer incidence and survival in the United States: Surveillance, Epidemiology, and End Results Program, 1973 to 2001. *Neurosurg Focus* 2006;20:E1. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16709014>.
4. Pouchieu C, Gruber A, Berteaud E, et al. Increasing incidence of central nervous system (CNS) tumors (2000-2012): findings from a population based registry in Gironde (France). *BMC Cancer* 2018;18:653. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29898691>.
5. Caldarella A, Crocetti E, Paci E. Is the incidence of brain tumors really increasing? A population-based analysis from a cancer registry. *J Neurooncol* 2011;104:589-594. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21327863>.
6. Arora RS, Alston RD, Eden TO, et al. Are reported increases in incidence of primary CNS tumours real? An analysis of longitudinal trends in England, 1979-2003. *Eur J Cancer* 2010;46:1607-1616. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20194015>.
7. U.S. National Library of Medicine-Key MEDLINE® Indicators. Available at: [http://www.nlm.nih.gov/bsd/bsd\\_key.html](http://www.nlm.nih.gov/bsd/bsd_key.html). Accessed July 24, 2014.
8. Sawaya R, Hammoud M, Schoppa D, et al. Neurosurgical outcomes in a modern series of 400 craniotomies for treatment of parenchymal tumors. *Neurosurgery* 1998;42:1044-1055; discussion 1055-1046. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9588549>.
9. Louis DN, Perry A, Reifenberger G, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol* 2016;131:803-820. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27157931>.
10. Yan H, Parsons DW, Jin G, et al. IDH1 and IDH2 mutations in gliomas. *N Engl J Med* 2009;360:765-773. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19228619>.
11. Houillier C, Wang X, Kaloshi G, et al. IDH1 or IDH2 mutations predict longer survival and response to temozolomide in low-grade gliomas. *Neurology* 2010;75:1560-1566. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20975057>.
12. Cancer Genome Atlas Research Network, Brat DJ, Verhaak RG, et al. Comprehensive, integrative genomic analysis of diffuse lower-grade gliomas. *N Engl J Med* 2015;372:2481-2498. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26061751>.
13. Eckel-Passow JE, Lachance DH, Molinaro AM, et al. Glioma groups based on 1p/19q, IDH, and TERT promoter mutations in tumors. *N Engl J Med* 2015;372:2499-2508. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26061753>.
14. Wiestler B, Capper D, Sill M, et al. Integrated DNA methylation and copy-number profiling identify three clinically and biologically relevant groups of anaplastic glioma. *Acta Neuropathol* 2014;128:561-571. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25008768>.
15. Weller M, Weber RG, Willscher E, et al. Molecular classification of diffuse cerebral WHO grade II/III gliomas using genome- and transcriptome-wide profiling improves stratification of prognostically distinct patient groups. *Acta Neuropathol* 2015;129:679-693. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25783747>.
16. Nitta M, Muragaki Y, Maruyama T, et al. Proposed therapeutic strategy for adult low-grade glioma based on aggressive tumor resection. *Neurosurg Focus* 2015;38:E7. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25599276>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

17. Baumert BG, Hegi ME, van den Bent MJ, et al. Temozolomide chemotherapy versus radiotherapy in high-risk low-grade glioma (EORTC 22033-26033): a randomised, open-label, phase 3 intergroup study. *Lancet Oncol* 2016;17:1521-1532. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27686946>.

18. Sanson M, Marie Y, Paris S, et al. Isocitrate dehydrogenase 1 codon 132 mutation is an important prognostic biomarker in gliomas. *J Clin Oncol* 2009;27:4150-4154. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19636000>.

19. Sahm F, Reuss D, Koelsche C, et al. Farewell to oligoastrocytoma: in situ molecular genetics favor classification as either oligodendroglioma or astrocytoma. *Acta Neuropathol* 2014;128:551-559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25143301>.

20. Pai T, Epari S, Desai S, et al. Histological spectrum of oligodendroglial tumors: Only a subset shows 1p/19q codeletion. *Neurol India* 2017;65:113-120. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28084254>.

21. Neumann JE, Dorostkar MM, Korshunov A, et al. Distinct histomorphology in molecular subgroups of glioblastomas in young patients. *J Neuropathol Exp Neurol* 2016;75:408-414. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26975364>.

22. Dubbink HJ, Atmodimedjo PN, Kros JM, et al. Molecular classification of anaplastic oligodendroglioma using next-generation sequencing: a report of the prospective randomized EORTC Brain Tumor Group 26951 phase III trial. *Neuro Oncol* 2016;18:388-400. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26354927>.

23. Labussiere M, Idbaih A, Wang XW, et al. All the 1p19q codeleted gliomas are mutated on IDH1 or IDH2. *Neurology* 2010;74:1886-1890. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20427748>.

24. Horbinski C. What do we know about IDH1/2 mutations so far, and how do we use it? *Acta Neuropathol* 2013;125:621-636. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23512379>.

25. Jiao Y, Killela PJ, Reitman ZJ, et al. Frequent ATRX, CIC, FUBP1 and IDH1 mutations refine the classification of malignant gliomas. *Oncotarget* 2012;3:709-722. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22869205>.

26. Leeper HE, Caron AA, Decker PA, et al. IDH mutation, 1p19q codeletion and ATRX loss in WHO grade II gliomas. *Oncotarget* 2015;6:30295-30305. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26210286>.

27. Reuss DE, Sahm F, Schrimpf D, et al. ATRX and IDH1-R132H immunohistochemistry with subsequent copy number analysis and IDH sequencing as a basis for an "integrated" diagnostic approach for adult astrocytoma, oligodendroglioma and glioblastoma. *Acta Neuropathol* 2015;129:133-146. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25427834>.

28. Arita H, Narita Y, Fukushima S, et al. Upregulating mutations in the TERT promoter commonly occur in adult malignant gliomas and are strongly associated with total 1p19q loss. *Acta Neuropathol* 2013;126:267-276. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23764841>.

29. Killela PJ, Reitman ZJ, Jiao Y, et al. TERT promoter mutations occur frequently in gliomas and a subset of tumors derived from cells with low rates of self-renewal. *Proc Natl Acad Sci U S A* 2013;110:6021-6026. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23530248>.

30. Kristensen BW, Priesterbach-Ackley LP, Petersen JK, Wesseling P. Molecular pathology of tumors of the central nervous system. *Ann Oncol* 2019;30:1265-1278. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31124566>.

31. Meyronet D, Esteban-Mader M, Bonnet C, et al. Characteristics of H3 K27M-mutant gliomas in adults. *Neuro Oncol* 2017;19:1127-1134. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28201752>.

32. Brat DJ, Aldape K, Colman H, et al. cIMPACT-NOW update 5: recommended grading criteria and terminologies for IDH-mutant



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

astrocytomas. *Acta Neuropathol* 2020;139:603-608. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31996992>.

33. Jiang H, Cui Y, Wang J, Lin S. Impact of epidemiological characteristics of supratentorial gliomas in adults brought about by the 2016 world health organization classification of tumors of the central nervous system. *Oncotarget* 2017;8:20354-20361. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27888628>.

34. Wick W, Roth P, Hartmann C, et al. Long-term analysis of the NOA-04 randomized phase III trial of sequential radiochemotherapy of anaplastic glioma with PCV or temozolomide. *Neuro Oncol* 2016;18:1529-1537. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27370396>.

35. Cairncross JG, Wang M, Jenkins RB, et al. Benefit from procarbazine, lomustine, and vincristine in oligodendroglial tumors is associated with mutation of IDH. *J Clin Oncol* 2014;32:783-790. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24516018>.

36. Everhard S, Kaloshi G, Criniere E, et al. MGMT methylation: a marker of response to temozolomide in low-grade gliomas. *Ann Neurol* 2006;60:740-743. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17192931>.

37. Gorovets D, Kannan K, Shen R, et al. IDH mutation and neuroglial developmental features define clinically distinct subclasses of lower grade diffuse astrocytic glioma. *Clin Cancer Res* 2012;18:2490-2501. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22415316>.

38. Wahl M, Phillips JJ, Molinaro AM, et al. Chemotherapy for adult low-grade gliomas: clinical outcomes by molecular subtype in a phase II study of adjuvant temozolomide. *Neuro Oncol* 2017;19:242-251. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27571885>.

39. Arita H, Yamasaki K, Matsushita Y, et al. A combination of TERT promoter mutation and MGMT methylation status predicts clinically relevant subgroups of newly diagnosed glioblastomas. *Acta Neuropathol Commun* 2016;4:79. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27503138>.

40. Pekmezci M, Rice T, Molinaro AM, et al. Adult infiltrating gliomas with WHO 2016 integrated diagnosis: additional prognostic roles of ATRX and TERT. *Acta Neuropathol* 2017;133:1001-1016. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28255664>.

41. Mollemann M, Wolter M, Felsberg J, et al. Frequent promoter hypermethylation and low expression of the MGMT gene in oligodendroglial tumors. *Int J Cancer* 2005;113:379-385. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15455350>.

42. Hegi ME, Diserens AC, Gorlia T, et al. MGMT gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med* 2005;352:997-1003. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15758010>.

43. Hegi ME, Diserens AC, Godard S, et al. Clinical trial substantiates the predictive value of O-6-methylguanine-DNA methyltransferase promoter methylation in glioblastoma patients treated with temozolomide. *Clin Cancer Res* 2004;10:1871-1874. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15041700>.

44. Bell EH, Zhang P, Fisher BJ, et al. Association of MGMT promoter methylation status with survival outcomes in patients with high-risk glioma treated with radiotherapy and temozolomide: an analysis from the NRG Oncology/RTOG 0424 trial. *JAMA Oncol* 2018;4:1405-1409. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29955793>.

45. Malmstrom A, Gronberg BH, Marosi C, et al. Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised, phase 3 trial. *Lancet Oncol* 2012;13:916-926. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22877848>.

46. Wick W, Platten M, Meisner C, et al. Temozolomide chemotherapy alone versus radiotherapy alone for malignant astrocytoma in the elderly: the NOA-08 randomised, phase 3 trial. *Lancet Oncol* 2012;13:707-715. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22578793>.

47. Khuong-Quang DA, Buczkowicz P, Rakopoulos P, et al. K27M mutation in histone H3.3 defines clinically and biologically distinct



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

subgroups of pediatric diffuse intrinsic pontine gliomas. *Acta Neuropathol* 2012;124:439-447. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22661320>.

48. Sturm D, Witt H, Hovestadt V, et al. Hotspot mutations in H3F3A and IDH1 define distinct epigenetic and biological subgroups of glioblastoma. *Cancer Cell* 2012;22:425-437. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23079654>.

49. Korshunov A, Capper D, Reuss D, et al. Histologically distinct neuroepithelial tumors with histone 3 G34 mutation are molecularly similar and comprise a single nosologic entity. *Acta Neuropathol* 2016;131:137-146. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26482474>.

50. Horbinski C. To BRAF or not to BRAF: is that even a question anymore? *J Neuropathol Exp Neurol* 2013;72:2-7. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23242278>.

51. Hawkins C, Walker E, Mohamed N, et al. BRAF-KIAA1549 fusion predicts better clinical outcome in pediatric low-grade astrocytoma. *Clin Cancer Res* 2011;17:4790-4798. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21610142>.

52. Horbinski C, Nikiforova MN, Hagenkord JM, et al. Interplay among BRAF, p16, p53, and MIB1 in pediatric low-grade gliomas. *Neuro Oncol* 2012;14:777-789. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22492957>.

53. Penman CL, Faulkner C, Lowis SP, Kurian KM. Current understanding of BRAF alterations in diagnosis, prognosis, and therapeutic targeting in pediatric low-grade gliomas. *Front Oncol* 2015;5:54. Available at:

<http://na1.replicon.com/NCCN/main.aspx>.

54. Kleinschmidt-DeMasters BK, Aisner DL, Birks DK, Foreman NK. Epithelioid GBMs show a high percentage of BRAF V600E mutation. *Am J Surg Pathol* 2013;37:685-698. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23552385>.

55. Mistry M, Zhukova N, Merico D, et al. BRAF mutation and CDKN2A deletion define a clinically distinct subgroup of childhood secondary high-grade glioma. *J Clin Oncol* 2015;33:1015-1022. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25667294>.

56. Chapman PB, Hauschild A, Robert C, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med* 2011;364:2507-2516. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21639808>.

57. McArthur GA, Chapman PB, Robert C, et al. Safety and efficacy of vemurafenib in BRAF(V600E) and BRAF(V600K) mutation-positive melanoma (BRIM-3): extended follow-up of a phase 3, randomised, open-label study. *Lancet Oncol* 2014;15:323-332. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24508103>.

58. Kaley T, Touat M, Subbiah V, et al. BRAF inhibition in BRAF(V600)-mutant gliomas: results from the VE-BASKET Study. *J Clin Oncol* 2018;36:3477-3484. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30351999>.

59. Chang S, Zhang P, Cairncross JG, et al. Phase III randomized study of radiation and temozolomide versus radiation and nitrosourea therapy for anaplastic astrocytoma: results of NRG Oncology RTOG 9813. *Neuro Oncol* 2017;19:252-258. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27994066>.

60. Olar A, Wani KM, Alfaro-Munoz KD, et al. IDH mutation status and role of WHO grade and mitotic index in overall survival in grade II-III diffuse gliomas. *Acta Neuropathol* 2015;129:585-596. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25701198>.

61. Hegi ME, Genbrugge E, Gorlia T, et al. MGMT promoter methylation cutoff with safety margin for selecting glioblastoma patients into trials omitting temozolomide: a pooled analysis of four clinical trials. *Clin Cancer Res* 2019;25:1809-1816. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30514777>.





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

62. Ostrom QT, Gittleman H, Xu J, et al. CBTRUS Statistical Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2009-2013. *Neuro Oncol* 2016;18:v1-v75. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28475809>.

63. Chang EF, Potts MB, Keles GE, et al. Seizure characteristics and control following resection in 332 patients with low-grade gliomas. *J Neurosurg* 2008;108:227-235. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18240916>.

64. Schiff D, Van den Bent M, Vogelbaum MA, et al. Recent developments and future directions in adult lower-grade gliomas: Society for Neuro-Oncology (SNO) and European Association of Neuro-Oncology (EANO) consensus. *Neuro Oncol* 2019;21:837-853. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30753579>.

65. Piepmeier J, Christopher S, Spencer D, et al. Variations in the natural history and survival of patients with supratentorial low-grade astrocytomas. *Neurosurgery* 1996;38:872-878; discussion 878-879. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8727811>.

66. Afra D, Osztie E, Sipos L, Vitanovics D. Preoperative history and postoperative survival of supratentorial low-grade astrocytomas. *Br J Neurosurg* 1999;13:299-305. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10562842>.

67. Gallo P, Cecchi PC, Locatelli F, et al. Pleomorphic xanthoastrocytoma: long-term results of surgical treatment and analysis of prognostic factors. *Br J Neurosurg* 2013;27:759-764. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23514331>.

68. Giannini C, Scheithauer BW, Burger PC, et al. Pleomorphic xanthoastrocytoma: what do we really know about it? *Cancer* 1999;85:2033-2045. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10223246>.

69. Ida CM, Rodriguez FJ, Burger PC, et al. Pleomorphic xanthoastrocytoma: natural history and long-term follow-up. *Brain Pathol*

2015;25:575-586. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25318587>.

70. Varshneya K, Sarmiento JM, Nuno M, et al. A national perspective of adult gangliogliomas. *J Clin Neurosci* 2016;30:65-70. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27083133>.

71. Tahiri Elousrouti L, Lamchahab M, Bougtoub N, et al. Subependymal giant cell astrocytoma (SEGA): a case report and review of the literature. *J Med Case Rep* 2016;10:35. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26861567>.

72. Roth J, Roach ES, Bartels U, et al. Subependymal giant cell astrocytoma: diagnosis, screening, and treatment. Recommendations from the International Tuberous Sclerosis Complex Consensus Conference 2012. *Pediatr Neurol* 2013;49:439-444. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24138953>.

73. Adriaensen ME, Schaefer-Prokop CM, Stijnen T, et al. Prevalence of subependymal giant cell tumors in patients with tuberous sclerosis and a review of the literature. *Eur J Neurol* 2009;16:691-696. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19236458>.

74. Goh S, Butler W, Thiele EA. Subependymal giant cell tumors in tuberous sclerosis complex. *Neurology* 2004;63:1457-1461. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15505165>.

75. Skalicky AM, Rentz AM, Liu Z, et al. The burden of subependymal giant cell astrocytomas associated with tuberous sclerosis complex: results of a patient and caregiver survey. *J Child Neurol* 2015;30:563-569. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24667738>.

76. Franz DN, Bissler JJ, McCormack FX. Tuberous sclerosis complex: neurological, renal and pulmonary manifestations. *Neuropediatrics* 2010;41:199-208. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21210335>.

77. Sun P, Kohrman M, Liu J, et al. Outcomes of resecting subependymal giant cell astrocytoma (SEGA) among patients with SEGA-related





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

tuberous sclerosis complex: a national claims database analysis. *Curr Med Res Opin* 2012;28:657-663. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22375958>.

78. Berhouma M, Dubourg J, Messerer M. Neurology: Letter to the editor. Re: Sun P, Kohrman M, Liu J et al. Outcomes of resecting subependymal giant cell astrocytoma (SEGA) among patients with SEGA-related tuberous sclerosis complex: a national claims database analysis. *Curr Med Res Opin* 2012;28:657-63. *Curr Med Res Opin* 2012;28:1571-1572; author reply 1572-1573. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22876836>.

79. Wen P, Stein A, van den Bent M, et al. ACTR-30. Updated efficacy and safety of dabrafenib plus trametinib in patients with recurrent/refractory BRAF V600E-mutated high-grade glioma (HGG) and low-grade glioma (LGG). *Neuro-Oncology* 2019;21:vi19-vi20. Available at: <https://doi.org/10.1093/neuonc/noz175.073>.

80. Brown NF, Carter T, Kitchen N, Mulholland P. Dabrafenib and trametinib in BRAFV600E mutated glioma. *CNS Oncol* 2017;6:291-296. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28984141>.

81. Marks AM, Bindra RS, DiLuna ML, et al. Response to the BRAF/MEK inhibitors dabrafenib/trametinib in an adolescent with a BRAF V600E mutated anaplastic ganglioglioma intolerant to vemurafenib. *Pediatr Blood Cancer* 2018;65:e26969. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29380516>.

82. Franz DN, Belousova E, Sparagana S, et al. Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial. *Lancet* 2013;381:125-132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23158522>.

83. Franz DN, Belousova E, Sparagana S, et al. Everolimus for subependymal giant cell astrocytoma in patients with tuberous sclerosis complex: 2-year open-label extension of the randomised EXIST-1 study. *Lancet Oncol* 2014;15:1513-1520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25456370>.

84. Wahl M, Chang SM, Phillips JJ, et al. Probing the phosphatidylinositol 3-kinase/mammalian target of rapamycin pathway in gliomas: A phase 2 study of everolimus for recurrent adult low-grade gliomas. *Cancer* 2017;123:4631-4639. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28759109>.

85. Ebrahimi-Fakhari D, Franz DN. Pharmacological treatment strategies for subependymal giant cell astrocytoma (SEGA). *Expert Opin Pharmacother* 2020;21:1329-1336. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32338549>.

86. Ostrom QT, Cioffi G, Gittleman H, et al. CBTRUS Statistical Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2012-2016. *Neuro Oncol* 2019;21:v1-v100. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31675094>.

87. Pignatti F, van den Bent M, Curran D, et al. Prognostic factors for survival in adult patients with cerebral low-grade glioma. *J Clin Oncol* 2002;20:2076-2084. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11956268>.

88. Daniels TB, Brown PD, Felten SJ, et al. Validation of EORTC prognostic factors for adults with low-grade glioma: a report using intergroup 86-72-51. *Int J Radiat Oncol Biol Phys* 2011;81:218-224. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21549518>.

89. Lo SS, Cho KH, Hall WA, et al. Does the extent of surgery have an impact on the survival of patients who receive postoperative radiation therapy for supratentorial low-grade gliomas? *Int J Cancer* 2001;96 Suppl:71-78. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11992388>.

90. Shaw E, Arusell R, Scheithauer B, et al. Prospective randomized trial of low- versus high-dose radiation therapy in adults with supratentorial low-grade glioma: initial report of a North Central Cancer Treatment Group/Radiation Therapy Oncology Group/Eastern Cooperative Oncology Group study. *J Clin Oncol* 2002;20:2267-2276. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11980997>.



91. Lo SS, Hall WA, Cho KH, et al. Radiation dose response for supratentorial low-grade glioma--institutional experience and literature review. *J Neurol Sci* 2003;214:43-48. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12972387>.

92. Shaw EG, Berkey B, Coons SW, et al. Recurrence following neurosurgeon-determined gross-total resection of adult supratentorial low-grade glioma: results of a prospective clinical trial. *J Neurosurg* 2008;109:835-841. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18976072>.

93. Jackson RJ, Fuller GN, Abi-Said D, et al. Limitations of stereotactic biopsy in the initial management of gliomas. *Neuro Oncol* 2001;3:193-200. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11465400>.

94. Villena Martin M, Pena Pardo FJ, Jimenez Aragon F, et al. Metabolic targeting can improve the efficiency of brain tumor biopsies. *Semin Oncol* 2020;47:148-154. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32513423>.

95. Brown TJ, Bota DA, van Den Bent MJ, et al. Management of low-grade glioma: a systematic review and meta-analysis. *Neurooncol Pract* 2019;6:249-258. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31386075>.

96. Yang K, Nath S, Koziarz A, et al. Biopsy versus subtotal versus gross total resection in patients with low-grade glioma: a systematic review and meta-analysis. *World Neurosurg* 2018;120:e762-e775. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30172972>.

97. Tang S, Liao J, Long Y. Comparative assessment of the efficacy of gross total versus subtotal total resection in patients with glioma: a meta-analysis. *Int J Surg* 2019;63:90-97. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30742934>.

98. McGirt MJ, Chaichana KL, Attenello FJ, et al. Extent of surgical resection is independently associated with survival in patients with hemispheric infiltrating low-grade gliomas. *Neurosurgery* 2008;63:700-

707; author reply 707-708. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18981880>.

99. Jakola AS, Myrnes KS, Kloster R, et al. Comparison of a strategy favoring early surgical resection vs a strategy favoring watchful waiting in low-grade gliomas. *JAMA* 2012;308:1881-1888. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23099483>.

100. Berger MS, Deliganis AV, Dobbins J, Keles GE. The effect of extent of resection on recurrence in patients with low grade cerebral hemisphere gliomas. *Cancer* 1994;74:1784-1791. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8082081>.

101. Kilic T, Ozduman K, Elmaci I, et al. Effect of surgery on tumor progression and malignant degeneration in hemispheric diffuse low-grade astrocytomas. *J Clin Neurosci* 2002;9:549-552. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12383413>.

102. Roelz R, Strohmaier D, Jabbarli R, et al. Residual tumor volume as best outcome predictor in low grade glioma - a nine-years near-randomized survey of surgery vs. biopsy. *Sci Rep* 2016;6:32286. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27574036>.

103. Karim AB, Maat B, Hatlevoll R, et al. A randomized trial on dose-response in radiation therapy of low-grade cerebral glioma: European Organization for Research and Treatment of Cancer (EORTC) Study 22844. *Int J Radiat Oncol Biol Phys* 1996;36:549-556. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8948338>.

104. van den Bent MJ, Afra D, de Witte O, et al. Long-term efficacy of early versus delayed radiotherapy for low-grade astrocytoma and oligodendroglioma in adults: the EORTC 22845 randomised trial. *Lancet* 2005;366:985-990. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16168780>.

105. Gorlia T, Wu W, Wang M, et al. New validated prognostic models and prognostic calculators in patients with low-grade gliomas diagnosed by central pathology review: a pooled analysis of EORTC/RTOG/NCCTG



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

phase III clinical trials. *Neuro Oncol* 2013;15:1568-1579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24049111>.

106. Shaw EG, Wang M, Coons SW, et al. Randomized trial of radiation therapy plus procarbazine, lomustine, and vincristine chemotherapy for supratentorial adult low-grade glioma: initial results of RTOG 9802. *J Clin Oncol* 2012;30:3065-3070. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22851558>.

107. Buckner JC, Shaw EG, Pugh SL, et al. Radiation plus procarbazine, CCNU, and vincristine in low-grade glioma. *N Engl J Med* 2016;374:1344-1355. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27050206>.

108. Bell EH, Zhang P, Shaw EG, et al. Comprehensive genomic analysis in NRG Oncology/RTOG 9802: a phase III trial of radiation versus radiation plus procarbazine, lomustine (CCNU), and vincristine in high-risk low-grade glioma. *J Clin Oncol* 2020;JCO1902983. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32706640>.

109. Fisher BJ, Hu C, Macdonald DR, et al. Phase 2 study of temozolomide-based chemoradiation therapy for high-risk low-grade gliomas: preliminary results of Radiation Therapy Oncology Group 0424. *Int J Radiat Oncol Biol Phys* 2015;91:497-504. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25680596>.

110. Navarria P, Pessina F, Cozzi L, et al. Can advanced new radiation therapy technologies improve outcome of high grade glioma (HGG) patients? analysis of 3D-conformal radiotherapy (3DCRT) versus volumetric-modulated arc therapy (VMAT) in patients treated with surgery, concomitant and adjuvant chemo-radiotherapy. *BMC Cancer* 2016;16:362. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27287048>.

111. Ding M, Newman F, Chen C, et al. Dosimetric comparison between 3DCRT and IMRT using different multileaf collimators in the treatment of brain tumors. *Med Dosim* 2009;34:1-8. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19181248>.

112. Shaw EG, Daumas-Duport C, Scheithauer BW, et al. Radiation therapy in the management of low-grade supratentorial astrocytomas. *J*

*Neurosurg* 1989;70:853-861. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2715812>.

113. Indelicato DJ, Rotondo RL, Uezono H, et al. Outcomes following proton therapy for pediatric low-grade glioma. *Int J Radiat Oncol Biol Phys* 2019;104:149-156. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30684665>.

114. Nahed BV, Redjal N, Brat DJ, et al. Management of patients with recurrence of diffuse low grade glioma: A systematic review and evidence-based clinical practice guideline. *J Neurooncol* 2015;125:609-630. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26530264>.

115. Ramakrishna R, Hebb A, Barber J, et al. Outcomes in reoperated low-grade gliomas. *Neurosurgery* 2015;77:175-184; discussion 184. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25850601>.

116. Smith JS, Chang EF, Lamborn KR, et al. Role of extent of resection in the long-term outcome of low-grade hemispheric gliomas. *J Clin Oncol* 2008;26:1338-1345. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18323558>.

117. Uppstrom TJ, Singh R, Hadjigeorgiou GF, et al. Repeat surgery for recurrent low-grade gliomas should be standard of care. *Clin Neurol Neurosurg* 2016;151:18-23. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27736650>.

118. Kesari S, Schiff D, Drappatz J, et al. Phase II study of protracted daily temozolomide for low-grade gliomas in adults. *Clin Cancer Res* 2009;15:330-337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19118062>.

119. Nicholson HS, Kretschmar CS, Krailo M, et al. Phase 2 study of temozolomide in children and adolescents with recurrent central nervous system tumors: a report from the Children's Oncology Group. *Cancer* 2007;110:1542-1550. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17705175>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

120. Soffietti R, Ruda R, Bradac GB, Schiffer D. PCV chemotherapy for recurrent oligodendrogliomas and oligoastrocytomas. *Neurosurgery* 1998;43:1066-1073. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9802850>.

121. Moghrabi A, Friedman HS, Ashley DM, et al. Phase II study of carboplatin (CBDCA) in progressive low-grade gliomas. *Neurosurg Focus* 1998;4:e3. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17168503>.

122. Brandes AA, Basso U, Vastola F, et al. Carboplatin and teniposide as third-line chemotherapy in patients with recurrent oligodendroglioma or oligoastrocytoma: a phase II study. *Ann Oncol* 2003;14:1727-1731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14630676>.

123. Triebels VH, Taphoorn MJ, Brandes AA, et al. Salvage PCV chemotherapy for temozolomide-resistant oligodendrogliomas. *Neurology* 2004;63:904-906. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15365146>.

124. van den Bent MJ, Smits M, Kros JM, Chang SM. Diffuse infiltrating oligodendroglioma and astrocytoma. *J Clin Oncol* 2017;35:2394-2401. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28640702>.

125. van den Bent MJ, Baumert B, Erridge SC, et al. Interim results from the CATNON trial (EORTC study 26053-22054) of treatment with concurrent and adjuvant temozolomide for 1p/19q non-co-deleted anaplastic glioma: a phase 3, randomised, open-label intergroup study. *Lancet* 2017;390:1645-1653. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28801186>.

126. Perry JR, Laperriere N, O'Callaghan CJ, et al. Short-course radiation plus temozolomide in elderly patients with glioblastoma. *N Engl J Med* 2017;376:1027-1037. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28296618>.

127. Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 2005;352:987-996. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15758009>.

128. Acharya S, Robinson CG, Michalski JM, et al. Association of 1p/19q codeletion and radiation necrosis in adult cranial gliomas after proton or photon therapy. *Int J Radiat Oncol Biol Phys* 2018;101:334-343. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29534896>.

129. Wen PY, Weller M, Lee EQ, et al. Glioblastoma in adults: a Society for Neuro-Oncology (SNO) and European Society of Neuro-Oncology (EANO) consensus review on current management and future directions. *Neuro Oncol* 2020;22:1073-1113. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32328653>.

130. Arevalo OD, Soto C, Rabiei P, et al. Assessment of glioblastoma response in the era of bevacizumab: longstanding and emergent challenges in the imaging evaluation of pseudoresponse. *Front Neurol* 2019;10:460. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31133966>.

131. Li XZ, Li YB, Cao Y, et al. Prognostic implications of resection extent for patients with glioblastoma multiforme: a meta-analysis. *J Neurosurg Sci* 2017;61:631-639. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26824196>.

132. Beiko J, Suki D, Hess KR, et al. IDH1 mutant malignant astrocytomas are more amenable to surgical resection and have a survival benefit associated with maximal surgical resection. *Neuro Oncol* 2014;16:81-91. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24305719>.

133. Molinaro AM, Hervey-Jumper S, Morshed RA, et al. Association of maximal extent of resection of contrast-enhanced and non-contrast-enhanced tumor with survival within molecular subgroups of patients with newly diagnosed glioblastoma. *JAMA Oncol* 2020;6:495-503. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32027343>.

134. Barker FG, 2nd, Chang SM, Gutin PH, et al. Survival and functional status after resection of recurrent glioblastoma multiforme. *Neurosurgery* 1998;42:709-720; discussion 720-703. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9574634>.





135. Park JK, Hodges T, Arko L, et al. Scale to predict survival after surgery for recurrent glioblastoma multiforme. *J Clin Oncol* 2010;28:3838-3843. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20644085>.

136. Kristiansen K, Hagen S, Kollevold T, et al. Combined modality therapy of operated astrocytomas grade III and IV. Confirmation of the value of postoperative irradiation and lack of potentiation of bleomycin on survival time: a prospective multicenter trial of the Scandinavian Glioblastoma Study Group. *Cancer* 1981;47:649-652. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6164465>.

137. Walker MD, Alexander E, Jr., Hunt WE, et al. Evaluation of BCNU and/or radiotherapy in the treatment of anaplastic gliomas. A cooperative clinical trial. *J Neurosurg* 1978;49:333-343. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/355604>.

138. Cabrera AR, Kirkpatrick JP, Fiveash JB, et al. Radiation therapy for glioblastoma: Executive summary of an American Society for Radiation Oncology Evidence-Based Clinical Practice Guideline. *Pract Radiat Oncol* 2016;6:217-225. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27211230>.

139. Roa W, Brasher PM, Bauman G, et al. Abbreviated course of radiation therapy in older patients with glioblastoma multiforme: a prospective randomized clinical trial. *J Clin Oncol* 2004;22:1583-1588. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15051755>.

140. Roa W, Kepka L, Kumar N, et al. International Atomic Energy Agency randomized phase III study of radiation therapy in elderly and/or frail patients with newly diagnosed glioblastoma multiforme. *J Clin Oncol* 2015;33:4145-4150. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26392096>.

141. Finazzi T. One week of radiotherapy for glioblastoma: a noninferiority trial? *J Clin Oncol* 2016;34:2192. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27069081>.

142. Kim H, Leiby BE, Shi W. Too little, too soon: short-course radiotherapy in elderly patients with glioblastoma. *J Clin Oncol*

2016;34:2191-2192. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27069070>.

143. Stupp R, Hegi ME, Mason WP, et al. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol* 2009;10:459-466. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19269895>.

144. van den Bent MJ, Carpentier AF, Brandes AA, et al. Adjuvant procarbazine, lomustine, and vincristine improves progression-free survival but not overall survival in newly diagnosed anaplastic oligodendrogliomas and oligoastrocytomas: a randomized European Organisation for Research and Treatment of Cancer phase III trial. *J Clin Oncol* 2006;24:2715-2722. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16782911>.

145. van den Bent MJ, Brandes AA, Taphoorn MJ, et al. Adjuvant procarbazine, lomustine, and vincristine chemotherapy in newly diagnosed anaplastic oligodendroglioma: long-term follow-up of EORTC brain tumor group study 26951. *J Clin Oncol* 2013;31:344-350. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23071237>.

146. Cairncross G, Berkey B, Shaw E, et al. Phase III trial of chemotherapy plus radiotherapy compared with radiotherapy alone for pure and mixed anaplastic oligodendroglioma: Intergroup Radiation Therapy Oncology Group Trial 9402. *J Clin Oncol* 2006;24:2707-2714. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16782910>.

147. Cairncross G, Wang M, Shaw E, et al. Phase III trial of chemoradiotherapy for anaplastic oligodendroglioma: long-term results of RTOG 9402. *J Clin Oncol* 2013;31:337-343. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23071247>.

148. Jaeckle KA, Ballman KV, van den Bent M, et al. CODEL: phase III study of RT, RT + temozolomide (TMZ), or TMZ for newly-diagnosed 1p/19q codeleted oligodendroglioma. Analysis from the initial study design. *Neuro Oncol* 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32678879>.





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

149. van den Bent MJ, Erridge SC, Vogelbaum MA, et al. Second interim and first molecular analysis of the EORTC randomized phase III intergroup CATNON trial on concurrent and adjuvant temozolomide in anaplastic glioma without 1p/19q codeletion. ASCO 2019. Available at: <https://meetinglibrary.asco.org/record/173361/abstract>.

150. Clarke JL, Iwamoto FM, Sul J, et al. Randomized phase II trial of chemoradiotherapy followed by either dose-dense or metronomic temozolomide for newly diagnosed glioblastoma. J Clin Oncol 2009;27:3861-3867. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19506159>.

151. Gilbert MR, Wang M, Aldape KD, et al. Dose-dense temozolomide for newly diagnosed glioblastoma: a randomized phase III clinical trial. J Clin Oncol 2013;31:4085-4091. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24101040>.

152. Stupp R, Hegi ME, Gorlia T, et al. Cilengitide combined with standard treatment for patients with newly diagnosed glioblastoma with methylated MGMT promoter (CENTRIC EORTC 26071-22072 study): a multicentre, randomised, open-label, phase 3 trial. Lancet Oncol 2014;15:1100-1108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25163906>.

153. Nabors LB, Fink KL, Mikkelsen T, et al. Two cilengitide regimens in combination with standard treatment for patients with newly diagnosed glioblastoma and unmethylated MGMT gene promoter: results of the open-label, controlled, randomized phase II CORE study. Neuro Oncol 2015;17:708-717. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25762461>.

154. Blumenthal DT, Gorlia T, Gilbert MR, et al. Is more better? The impact of extended adjuvant temozolomide in newly diagnosed glioblastoma: a secondary analysis of EORTC and NRG Oncology/RTOG. Neuro Oncol 2017;19:1119-1126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28371907>.

155. Balana C, Vaz MA, Sepulveda JM, et al. A phase II randomized, multicenter, open-label trial of continuing adjuvant temozolomide beyond

six cycles in patients with glioblastoma (GEINO 14-01). Neuro Oncol 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32328662>.

156. Glas M, Happold C, Rieger J, et al. Long-term survival of patients with glioblastoma treated with radiotherapy and lomustine plus temozolomide. J Clin Oncol 2009;27:1257-1261. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19188676>.

157. Herrlinger U, Rieger J, Koch D, et al. Phase II trial of lomustine plus temozolomide chemotherapy in addition to radiotherapy in newly diagnosed glioblastoma: UKT-03. J Clin Oncol 2006;24:4412-4417. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16983109>.

158. Herrlinger U, Tzaridis T, Mack F, et al. Lomustine-temozolomide combination therapy versus standard temozolomide therapy in patients with newly diagnosed glioblastoma with methylated MGMT promoter (CeTeG/NOA-09): a randomised, open-label, phase 3 trial. Lancet 2019;393:678-688. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30782343>.

159. Weller J, Tzaridis T, Mack F, et al. Health-related quality of life and neurocognitive functioning with lomustine-temozolomide versus temozolomide in patients with newly diagnosed, MGMT-methylated glioblastoma (CeTeG/NOA-09): a randomised, multicentre, open-label, phase 3 trial. Lancet Oncol 2019;20:1444-1453. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31488360>.

160. Stupp R, Taillibert S, Kanner AA, et al. Maintenance therapy with tumor-treating fields plus temozolomide vs temozolomide alone for glioblastoma: a randomized clinical trial. JAMA 2015;314:2535-2543. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26670971>.

161. Stupp R, Taillibert S, Kanner A, et al. Effect of tumor-treating fields plus maintenance temozolomide vs maintenance temozolomide alone on survival in patients with glioblastoma: a randomized clinical trial. JAMA 2017;318:2306-2316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29260225>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

162. Taphoorn MJB, Dirven L, Kanner AA, et al. Influence of treatment with tumor-treating fields on health-related quality of life of patients with newly diagnosed glioblastoma: a secondary analysis of a randomized clinical trial. *JAMA Oncol* 2018;4:495-504. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29392280>.

163. Chamberlain MC. Treatment for patients with newly diagnosed glioblastoma. *JAMA* 2016;315:2348. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27272590>.

164. Sampson JH. Alternating electric fields for the treatment of glioblastoma. *JAMA* 2015;314:2511-2513. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26670969>.

165. Perry JR, Rizek P, Cashman R, et al. Temozolomide rechallenge in recurrent malignant glioma by using a continuous temozolomide schedule: the "rescue" approach. *Cancer* 2008;113:2152-2157. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18756530>.

166. Brandes AA, Tosoni A, Amista P, et al. How effective is BCNU in recurrent glioblastoma in the modern era? A phase II trial. *Neurology* 2004;63:1281-1284. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15477552>.

167. Reithmeier T, Graf E, Piroth T, et al. BCNU for recurrent glioblastoma multiforme: efficacy, toxicity and prognostic factors. *BMC Cancer* 2010;10:30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20122270>.

168. Wick W, Puduvalli VK, Chamberlain MC, et al. Phase III study of enzastaurin compared with lomustine in the treatment of recurrent intracranial glioblastoma. *J Clin Oncol* 2010;28:1168-1174. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20124186>.

169. Taal W, Oosterkamp HM, Walenkamp AM, et al. Single-agent bevacizumab or lomustine versus a combination of bevacizumab plus lomustine in patients with recurrent glioblastoma (BELOB trial): a randomised controlled phase 2 trial. *Lancet Oncol* 2014;15:943-953. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25035291>.

170. Friedman HS, Prados MD, Wen PY, et al. Bevacizumab alone and in combination with irinotecan in recurrent glioblastoma. *J Clin Oncol* 2009;27:4733-4740. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19720927>.

171. Wick W, Gorlia T, Bendszus M, et al. Lomustine and bevacizumab in progressive glioblastoma. *N Engl J Med* 2017;377:1954-1963. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29141164>.

172. Ameratunga M, Pavlakakis N, Wheeler H, et al. Anti-angiogenic therapy for high-grade glioma. *Cochrane Database Syst Rev* 2018;11:CD008218. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30480778>.

173. Wick W, Weller M, van den Bent M, Stupp R. Bevacizumab and recurrent malignant gliomas: a European perspective. *J Clin Oncol* 2010;28:e188-189; author reply e190-182. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20159801>.

174. Kaley T, Nolan C, Carver A, Omuro A. Bevacizumab for acute neurologic deterioration in patients with glioblastoma. *CNS Oncol* 2013;2:413-418. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25054664>.

175. Lombardi G, De Salvo GL, Brandes AA, et al. Regorafenib compared with lomustine in patients with relapsed glioblastoma (REGOMA): a multicentre, open-label, randomised, controlled, phase 2 trial. *Lancet Oncol* 2019;20:110-119. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30522967>.

176. Brem H, Piantadosi S, Burger PC, et al. Placebo-controlled trial of safety and efficacy of intraoperative controlled delivery by biodegradable polymers of chemotherapy for recurrent gliomas. The Polymer-brain Tumor Treatment Group. *Lancet* 1995;345:1008-1012. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7723496>.

177. McGirt MJ, Brem H. Carmustine wafers (Gliadel) plus concomitant temozolomide therapy after resection of malignant astrocytoma: growing evidence for safety and efficacy. *Ann Surg Oncol* 2010;17:1729-1731. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20422453>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

178. Stupp R, Wong ET, Kanner AA, et al. NovoTTF-100A versus physician's choice chemotherapy in recurrent glioblastoma: a randomised phase III trial of a novel treatment modality. *Eur J Cancer* 2012;48:2192-2202. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22608262>.

179. Kazmi F, Soon YY, Leong YH, et al. Re-irradiation for recurrent glioblastoma (GBM): a systematic review and meta-analysis. *J Neurooncol* 2019;142:79-90. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30523605>.

180. Mohammadi AM, Sullivan TB, Barnett GH, et al. Use of high-field intraoperative magnetic resonance imaging to enhance the extent of resection of enhancing and nonenhancing gliomas. *Neurosurgery* 2014;74:339-348; discussion 349; quiz 349-350. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24368543>.

181. Coburger J, Wirtz CR. Fluorescence guided surgery by 5-ALA and intraoperative MRI in high grade glioma: a systematic review. *J Neurooncol* 2019;141:533-546. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30488293>.

182. Panageas KS, Iwamoto FM, Cloughesy TF, et al. Initial treatment patterns over time for anaplastic oligodendroglial tumors. *Neuro Oncol* 2012;14:761-767. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22661585>.

183. Brandes AA, Nicolardi L, Tosoni A, et al. Survival following adjuvant PCV or temozolomide for anaplastic astrocytoma. *Neuro Oncol* 2006;8:253-260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16723632>.

184. Levin VA, Silver P, Hannigan J, et al. Superiority of post-radiotherapy adjuvant chemotherapy with CCNU, procarbazine, and vincristine (PCV) over BCNU for anaplastic gliomas: NCOG 6G61 final report. *Int J Radiat Oncol Biol Phys* 1990;18:321-324. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2154418>.

185. Minniti G, Scaringi C, Lanzetta G, et al. Standard (60 Gy) or short-course (40 Gy) irradiation plus concomitant and adjuvant temozolomide for

elderly patients with glioblastoma: a propensity-matched analysis. *Int J Radiat Oncol Biol Phys* 2015;91:109-115. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25442339>.

186. Brandes AA, Franceschi E, Tosoni A, et al. MGMT promoter methylation status can predict the incidence and outcome of pseudoprogression after concomitant radiochemotherapy in newly diagnosed glioblastoma patients. *J Clin Oncol* 2008;26:2192-2197. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18445844>.

187. Ellingson BM, Chung C, Pope WB, et al. Pseudoprogression, radionecrosis, inflammation or true tumor progression? challenges associated with glioblastoma response assessment in an evolving therapeutic landscape. *J Neurooncol* 2017;134:495-504. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28382534>.

188. Tsien C, Galban CJ, Chenevert TL, et al. Parametric response map as an imaging biomarker to distinguish progression from pseudoprogression in high-grade glioma. *J Clin Oncol* 2010;28:2293-2299. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20368564>.

189. Fink J, Born D, Chamberlain MC. Pseudoprogression: relevance with respect to treatment of high-grade gliomas. *Curr Treat Options Oncol* 2011;12:240-252. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21594589>.

190. Perry JR, Belanger K, Mason WP, et al. Phase II trial of continuous dose-intense temozolomide in recurrent malignant glioma: RESCUE study. *J Clin Oncol* 2010;28:2051-2057. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20308655>.

191. Weller M, Tabatabai G, Kastner B, et al. MGMT promoter methylation is a strong prognostic biomarker for benefit from dose-intensified temozolomide rechallenge in progressive glioblastoma: the DIRECTOR trial. *Clin Cancer Res* 2015;21:2057-2064. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25655102>.

192. Yung WK, Albright RE, Olson J, et al. A phase II study of temozolomide vs. procarbazine in patients with glioblastoma multiforme at



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

first relapse. Br J Cancer 2000;83:588-593. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/10944597>.

193. Prados M, Rodriguez L, Chamberlain M, et al. Treatment of recurrent gliomas with 1,3-bis(2-chloroethyl)-1-nitrosourea and alpha-difluoromethylornithine. Neurosurgery 1989;24:806-809. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/2546093>.

194. Chamberlain MC, Johnston S. Bevacizumab for recurrent alkylator-refractory anaplastic oligodendroglioma. Cancer 2009;115:1734-1743. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19197992>.

195. Chamberlain MC, Johnston S. Salvage chemotherapy with bevacizumab for recurrent alkylator-refractory anaplastic astrocytoma. J Neurooncol 2009;91:359-367. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18953491>.

196. Norden AD, Young GS, Setayesh K, et al. Bevacizumab for recurrent malignant gliomas: efficacy, toxicity, and patterns of recurrence. Neurology 2008;70:779-787. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18316689>.

197. Cloughesy T, Prados MD, Mikkelsen T. A phase 2 randomized non-comparative clinical trial of the effect of bevacizumab alone or in combination with irinotecan on 6-month progression free survival in recurrent refractory glioblastoma [abstract]. J Clin Oncol 2008;26(Suppl 15):2010b. Available at:  
[http://meeting.ascopubs.org/cgi/content/abstract/26/15\\_suppl/2010b](http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/2010b).

198. Kreisl TN, Kim L, Moore K, et al. Phase II trial of single-agent bevacizumab followed by bevacizumab plus irinotecan at tumor progression in recurrent glioblastoma. J Clin Oncol 2009;27:740-745. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19114704>.

199. Vredenburgh JJ, Desjardins A, Herndon JE, 2nd, et al. Phase II trial of bevacizumab and irinotecan in recurrent malignant glioma. Clin Cancer Res 2007;13:1253-1259. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/17317837>.

200. Carvalho BF, Fernandes AC, Almeida DS, et al. Second-line chemotherapy in recurrent glioblastoma: a 2-cohort study. Oncol Res Treat 2015;38:348-354. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/26278578>.

201. Schmidt F, Fischer J, Herrlinger U, et al. PCV chemotherapy for recurrent glioblastoma. Neurology 2006;66:587-589. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/16505319>.

202. DeVita VT, Lawrence TS, Rosenberg SA. DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology (ed 8). Philadelphia: Lippincott Williams & Wilkins; 2008.

203. Packer RJ, Hoffman HJ, Friedman HS. Tumors of the fourth ventricle. In: Levin VA, ed. Cancer in the Nervous System. New York: Churchill Livingstone; 1996:153-156.

204. Pajtler KW, Witt H, Sill M, et al. Molecular classification of ependymal tumors across all CNS compartments, histopathological grades, and age groups. Cancer Cell 2015;27:728-743. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/25965575>.

205. Hubner JM, Kool M, Pfister SM, Pajtler KW. Epidemiology, molecular classification and WHO grading of ependymoma. J Neurosurg Sci 2018;62:46-50. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/28895660>.

206. Kawabata Y, Takahashi JA, Arakawa Y, Hashimoto N. Long-term outcome in patients harboring intracranial ependymoma. J Neurosurg 2005;103:31-37. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/16121970>.

207. Metellus P, Figarella-Branger D, Guyotat J, et al. Supratentorial ependymomas: prognostic factors and outcome analysis in a retrospective series of 46 adult patients. Cancer 2008;113:175-185. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18470910>.

208. Metellus P, Barrie M, Figarella-Branger D, et al. Multicentric French study on adult intracranial ependymomas: prognostic factors analysis and





therapeutic considerations from a cohort of 152 patients. Brain 2007;130:1338-1349. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17449478>.

209. Paulino AC, Wen BC, Buatti JM, et al. Intracranial ependymomas: an analysis of prognostic factors and patterns of failure. Am J Clin Oncol 2002;25:117-122. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11943886>.

210. Schwartz TH, Kim S, Glick RS, et al. Supratentorial ependymomas in adult patients. Neurosurgery 1999;44:721-731. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10201296>.

211. Abdulaziz M, Mallory GW, Bydon M, et al. Outcomes following myxopapillary ependymoma resection: the importance of capsule integrity. Neurosurg Focus 2015;39:E8. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26235025>.

212. Rodriguez D, Cheung MC, Housri N, et al. Outcomes of malignant CNS ependymomas: an examination of 2408 cases through the Surveillance, Epidemiology, and End Results (SEER) database (1973-2005). J Surg Res 2009;156:340-351. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19577759>.

213. Mansur DB, Perry A, Rajaram V, et al. Postoperative radiation therapy for grade II and III intracranial ependymoma. Int J Radiat Oncol Biol Phys 2005;61:387-391. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15667957>.

214. Merchant TE, Fouladi M. Ependymoma: new therapeutic approaches including radiation and chemotherapy. J Neurooncol 2005;75:287-299. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16195801>.

215. Taylor RE. Review of radiotherapy dose and volume for intracranial ependymoma. Pediatr Blood Cancer 2004;42:457-460. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15049020>.

216. Reni M, Brandes AA, Vavassori V, et al. A multicenter study of the prognosis and treatment of adult brain ependymal tumors. Cancer

2004;100:1221-1229. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15022290>.

217. Snider CA, Yang K, Mack SC, et al. Impact of radiation therapy and extent of resection for ependymoma in young children: A population-based study. Pediatr Blood Cancer 2018;65. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29115718>.

218. Ghia AJ, Mahajan A, Allen PK, et al. Supratentorial gross-totally resected non-anaplastic ependymoma: population based patterns of care and outcomes analysis. J Neurooncol 2013;115:513-520. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24085643>.

219. Metellus P, Guyotat J, Chinot O, et al. Adult intracranial WHO grade II ependymomas: long-term outcome and prognostic factor analysis in a series of 114 patients. Neuro Oncol 2010;12:976-984. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/20484442>.

220. Nuno M, Yu JJ, Varshneya K, et al. Treatment and survival of supratentorial and posterior fossa ependymomas in adults. J Clin Neurosci 2016;28:24-30. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26810473>.

221. Goldwein JW, Corn BW, Finlay JL, et al. Is craniospinal irradiation required to cure children with malignant (anaplastic) intracranial ependymomas? Cancer 1991;67:2766-2771. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/2025840>.

222. Vanuytsel L, Brada M. The role of prophylactic spinal irradiation in localized intracranial ependymoma. Int J Radiat Oncol Biol Phys 1991;21:825-830. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/1831193>.

223. Vanuytsel LJ, Bessell EM, Ashley SE, et al. Intracranial ependymoma: long-term results of a policy of surgery and radiotherapy. Int J Radiat Oncol Biol Phys 1992;23:313-319. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/1587752>.





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

224. Merchant TE, Li C, Xiong X, et al. Conformal radiotherapy after surgery for paediatric ependymoma: a prospective study. *Lancet Oncol* 2009;10:258-266. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19274783>.

225. Pieters RS, Niemierko A, Fullerton BC, Munzenrider JE. Cauda equina tolerance to high-dose fractionated irradiation. *Int J Radiat Oncol Biol Phys* 2006;64:251-257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15993548>.

226. Emami B, Lyman J, Brown A, et al. Tolerance of normal tissue to therapeutic irradiation. *Int J Radiat Oncol Biol Phys* 1991;21:109-122. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2032882>.

227. Lee SH, Chung CK, Kim CH, et al. Long-term outcomes of surgical resection with or without adjuvant radiation therapy for treatment of spinal ependymoma: a retrospective multicenter study by the Korea Spinal Oncology Research Group. *Neuro Oncol* 2013;15:921-929. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23576600>.

228. Abdel-Wahab M, Etuk B, Palermo J, et al. Spinal cord gliomas: A multi-institutional retrospective analysis. *Int J Radiat Oncol Biol Phys* 2006;64:1060-1071. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16373081>.

229. Kano H, Niranjan A, Kondziolka D, et al. Outcome predictors for intracranial ependymoma radiosurgery. *Neurosurgery* 2009;64:279-287; discussion 287-278. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19190457>.

230. Lo SS, Abdulrahman R, Desrosiers PM, et al. The role of Gamma Knife Radiosurgery in the management of unresectable gross disease or gross residual disease after surgery in ependymoma. *J Neurooncol* 2006;79:51-56. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16557349>.

231. Mansur DB, Drzymala RE, Rich KM, et al. The efficacy of stereotactic radiosurgery in the management of intracranial ependymoma. *J*

*Neurooncol* 2004;66:187-190. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15015785>.

232. Brandes AA, Cavallo G, Reni M, et al. A multicenter retrospective study of chemotherapy for recurrent intracranial ependymal tumors in adults by the Gruppo Italiano Cooperativo di Neuro-Oncologia. *Cancer* 2005;104:143-148. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15912507>.

233. Gornet MK, Buckner JC, Marks RS, et al. Chemotherapy for advanced CNS ependymoma. *J Neurooncol* 1999;45:61-67. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10728911>.

234. Chamberlain MC. Recurrent intracranial ependymoma in children: salvage therapy with oral etoposide. *Pediatr Neurol* 2001;24:117-121. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11275460>.

235. Sandri A, Massimino M, Mastrodicasa L, et al. Treatment with oral etoposide for childhood recurrent ependymomas. *J Pediatr Hematol Oncol* 2005;27:486-490. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16189442>.

236. Green RM, Cloughesy TF, Stupp R, et al. Bevacizumab for recurrent ependymoma. *Neurology* 2009;73:1677-1680. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19917990>.

237. Ruda R, Bosa C, Magistrello M, et al. Temozolomide as salvage treatment for recurrent intracranial ependymomas of the adult: a retrospective study. *Neuro Oncol* 2016;18:261-268. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26323606>.

238. Gilbert M, Yuan Y, Wani K, et al. AT-23 a phase II study of lapatinib and dose-dense temozolomide (TMZ) for adults with recurrent ependymoma: a CERN clinical trial. *Neuro Oncol* 2014;16:v13-v13. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4217801/>.

239. Kool M, Korshunov A, Remke M, et al. Molecular subgroups of medulloblastoma: an international meta-analysis of transcriptome, genetic aberrations, and clinical data of WNT, SHH, Group 3, and Group 4



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

medulloblastomas. *Acta Neuropathol* 2012;123:473-484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22358457>.

240. Sant M, Minicozzi P, Lagorio S, et al. Survival of European patients with central nervous system tumors. *Int J Cancer* 2012;131:173-185. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21805473>.

241. Brandes AA, Franceschi E. Shedding light on adult medulloblastoma: current management and opportunities for advances. *Am Soc Clin Oncol Educ Book* 2014:e82-87. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24857151>.

242. Northcott PA, Buchhalter I, Morrissy AS, et al. The whole-genome landscape of medulloblastoma subtypes. *Nature* 2017;547:311-317. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28726821>.

243. Sharma T, Schwalbe EC, Williamson D, et al. Second-generation molecular subgrouping of medulloblastoma: an international meta-analysis of Group 3 and Group 4 subtypes. *Acta Neuropathol* 2019;138:309-326. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31076851>.

244. Korshunov A, Remke M, Werft W, et al. Adult and pediatric medulloblastomas are genetically distinct and require different algorithms for molecular risk stratification. *J Clin Oncol* 2010;28:3054-3060. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20479417>.

245. Remke M, Hielscher T, Northcott PA, et al. Adult medulloblastoma comprises three major molecular variants. *J Clin Oncol* 2011;29:2717-2723. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21632505>.

246. Taylor MD, Northcott PA, Korshunov A, et al. Molecular subgroups of medulloblastoma: the current consensus. *Acta Neuropathol* 2012;123:465-472. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22134537>.

247. Prados MD, Warnick RE, Wara WM, et al. Medulloblastoma in adults. *Int J Radiat Oncol Biol Phys* 1995;32:1145-1152. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7607936>.

248. Waszak SM, Northcott PA, Buchhalter I, et al. Spectrum and prevalence of genetic predisposition in medulloblastoma: a retrospective genetic study and prospective validation in a clinical trial cohort. *Lancet Oncol* 2018;19:785-798. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29753700>.

249. Zhukova N, Ramaswamy V, Remke M, et al. Subgroup-specific prognostic implications of TP53 mutation in medulloblastoma. *J Clin Oncol* 2013;31:2927-2935. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23835706>.

250. Brandes AA, Bartolotti M, Marucci G, et al. New perspectives in the treatment of adult medulloblastoma in the era of molecular oncology. *Crit Rev Oncol Hematol* 2015;94:348-359. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25600839>.

251. Chan AW, Tarbell NJ, Black PM, et al. Adult medulloblastoma: prognostic factors and patterns of relapse. *Neurosurgery* 2000;47:623-631; discussion 631-622. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10981749>.

252. Frost PJ, Laperriere NJ, Wong CS, et al. Medulloblastoma in adults. *Int J Radiat Oncol Biol Phys* 1995;32:951-957. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7607969>.

253. Kunschner LJ, Kuttlesch J, Hess K, Yung WK. Survival and recurrence factors in adult medulloblastoma: the M.D. Anderson Cancer Center experience from 1978 to 1998. *Neuro Oncol* 2001;3:167-173. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11465397>.

254. Chargari C, Feuvret L, Levy A, et al. Reappraisal of clinical outcome in adult medulloblastomas with emphasis on patterns of relapse. *Br J Neurosurg* 2010;24:460-467. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20726753>.

255. Thompson EM, Hielscher T, Bouffet E, et al. Prognostic value of medulloblastoma extent of resection after accounting for molecular subgroup: a retrospective integrated clinical and molecular analysis.



Lancet Oncol 2016;17:484-495. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/26976201>.

256. Douglas JG, Barker JL, Ellenbogen RG, Geyer JR. Concurrent chemotherapy and reduced-dose cranial spinal irradiation followed by conformal posterior fossa tumor bed boost for average-risk medulloblastoma: efficacy and patterns of failure. Int J Radiat Oncol Biol Phys 2004;58:1161-1164. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/15001259>.

257. Merchant TE, Kun LE, Krasin MJ, et al. Multi-institution prospective trial of reduced-dose craniospinal irradiation (23.4 Gy) followed by conformal posterior fossa (36 Gy) and primary site irradiation (55.8 Gy) and dose-intensive chemotherapy for average-risk medulloblastoma. Int J Radiat Oncol Biol Phys 2008;70:782-787. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/17892918>.

258. Packer RJ, Gajjar A, Vezina G, et al. Phase III study of craniospinal radiation therapy followed by adjuvant chemotherapy for newly diagnosed average-risk medulloblastoma. J Clin Oncol 2006;24:4202-4208. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16943538>.

259. Deutsch M, Thomas PR, Krischer J, et al. Results of a prospective randomized trial comparing standard dose neuraxis irradiation (3,600 cGy/20) with reduced neuraxis irradiation (2,340 cGy/13) in patients with low-stage medulloblastoma. A Combined Children's Cancer Group-Pediatric Oncology Group Study. Pediatr Neurosurg 1996;24:167-176; discussion 176-167. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/8873158>.

260. Friedrich C, von Bueren AO, von Hoff K, et al. Treatment of adult nonmetastatic medulloblastoma patients according to the paediatric HIT 2000 protocol: a prospective observational multicentre study. Eur J Cancer 2013;49:893-903. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/23182688>.

261. Brown AP, Barney CL, Grosshans DR, et al. Proton beam craniospinal irradiation reduces acute toxicity for adults with

medulloblastoma. Int J Radiat Oncol Biol Phys 2013;86:277-284. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23433794>.

262. Germanwala AV, Mai JC, Tomycz ND, et al. Boost Gamma Knife surgery during multimodality management of adult medulloblastoma. J Neurosurg 2008;108:204-209. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18240913>.

263. Carrie C, Lasset C, Alapetite C, et al. Multivariate analysis of prognostic factors in adult patients with medulloblastoma. Retrospective study of 156 patients. Cancer 1994;74:2352-2360. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/7922986>.

264. Riffaud L, Saikali S, Leray E, et al. Survival and prognostic factors in a series of adults with medulloblastomas. J Neurosurg 2009;111:478-487. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19231932>.

265. Padovani L, Sunyach MP, Perol D, et al. Common strategy for adult and pediatric medulloblastoma: a multicenter series of 253 adults. Int J Radiat Oncol Biol Phys 2007;68:433-440. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/17498567>.

266. Herrlinger U, Steinbrecher A, Rieger J, et al. Adult medulloblastoma: prognostic factors and response to therapy at diagnosis and at relapse. J Neurol 2005;252:291-299. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/16189725>.

267. Moots PL, O'Neill A, Londer H, et al. Preradiation Chemotherapy for Adult High-risk Medulloblastoma: A Trial of the ECOG-ACRIN Cancer Research Group (E4397). Am J Clin Oncol 2018;41:588-594. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/27635620>.

268. Tarbell NJ, Friedman H, Polkinghorn WR, et al. High-risk medulloblastoma: a pediatric oncology group randomized trial of chemotherapy before or after radiation therapy (POG 9031). J Clin Oncol 2013;31:2936-2941. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/23857975>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

269. Zeltzer PM, Boyett JM, Finlay JL, et al. Metastasis stage, adjuvant treatment, and residual tumor are prognostic factors for medulloblastoma in children: conclusions from the Children's Cancer Group 921 randomized phase III study. *J Clin Oncol* 1999;17:832-845. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10071274>.

270. Ashley DM, Meier L, Kerby T, et al. Response of recurrent medulloblastoma to low-dose oral etoposide. *J Clin Oncol* 1996;14:1922-1927. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8656261>.

271. Chamberlain MC, Kormanik PA. Chronic oral VP-16 for recurrent medulloblastoma. *Pediatr Neurol* 1997;17:230-234. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9390699>.

272. Dunkel IJ, Gardner SL, Garvin JH, Jr., et al. High-dose carboplatin, thiopeta, and etoposide with autologous stem cell rescue for patients with previously irradiated recurrent medulloblastoma. *Neuro Oncol* 2010;12:297-303. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20167818>.

273. Wang CH, Hsu TR, Wong TT, Chang KP. Efficacy of temozolomide for recurrent embryonal brain tumors in children. *Childs Nerv Syst* 2009;25:535-541. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19107490>.

274. Gill P, Litzow M, Buckner J, et al. High-dose chemotherapy with autologous stem cell transplantation in adults with recurrent embryonal tumors of the central nervous system. *Cancer* 2008;112:1805-1811. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18300237>.

275. Robinson GW, Orr BA, Wu G, et al. Vismodegib exerts targeted efficacy against recurrent sonic hedgehog-subgroup medulloblastoma: results from phase II Pediatric Brain Tumor Consortium Studies PBTC-025B and PBTC-032. *J Clin Oncol* 2015;33:2646-2654. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26169613>.

276. Kieran MW, Chisholm J, Casanova M, et al. Phase I study of oral sonidegib (LDE225) in pediatric brain and solid tumors and a phase II study in children and adults with relapsed medulloblastoma. *Neuro Oncol*

2017;19:1542-1552. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28605510>.

277. Cohen ME, Duffner P, eds. Brain tumors in children (ed 2). New York: McGraw-Hill; 1994.

278. Chang CH, Housepian EM, Herbert C, Jr. An operative staging system and a megavoltage radiotherapeutic technic for cerebellar medulloblastomas. *Radiology* 1969;93:1351-1359. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/4983156>.

279. Brandes AA, Franceschi E, Tosoni A, et al. Adult neuroectodermal tumors of posterior fossa (medulloblastoma) and of supratentorial sites (stPNET). *Crit Rev Oncol Hematol* 2009;71:165-179. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19303318>.

280. Villano JL, Koshy M, Shaikh H, et al. Age, gender, and racial differences in incidence and survival in primary CNS lymphoma. *Br J Cancer* 2011;105:1414-1418. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21915121>.

281. O'Neill BP, Decker PA, Tieu C, Cerhan JR. The changing incidence of primary central nervous system lymphoma is driven primarily by the changing incidence in young and middle-aged men and differs from time trends in systemic diffuse large B-cell non-Hodgkin's lymphoma. *Am J Hematol* 2013;88:997-1000. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23873804>.

282. Fine HA, Mayer RJ. Primary central nervous system lymphoma. *Ann Intern Med* 1993;119:1093-1104. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8239229>.

283. Norden AD, Drappatz J, Wen PY, Claus EB. Survival among patients with primary central nervous system lymphoma, 1973-2004. *J Neurooncol* 2010. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20556477>.

284. Uldrick TS, Pipkin S, Scheer S, Hessel NA. Factors associated with survival among patients with AIDS-related primary central nervous system





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

lymphoma. *AIDS* 2014;28:397-405. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24076659>.

285. Gerstner ER, Batchelor TT. Primary central nervous system lymphoma. *Arch Neurol* 2010;67:291-297. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20212226>.

286. Batchelor TT. Primary central nervous system lymphoma: A curable disease. *Hematol Oncol* 2019;37 Suppl 1:15-18. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31187523>.

287. Bataille B, Delwail V, Menet E, et al. Primary intracerebral malignant lymphoma: report of 248 cases. *J Neurosurg* 2000;92:261-266. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10659013>.

288. Buell JF, Gross TG, Hanaway MJ, et al. Posttransplant lymphoproliferative disorder: significance of central nervous system involvement. *Transplant Proc* 2005;37:954-955. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15848587>.

289. Cavaliere R, Petroni G, Lopes MB, et al. Primary central nervous system post-transplantation lymphoproliferative disorder: an International Primary Central Nervous System Lymphoma Collaborative Group Report. *Cancer* 2010;116:863-870. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20052713>.

290. Evens AM, David KA, Helenowski I, et al. Multicenter analysis of 80 solid organ transplantation recipients with post-transplantation lymphoproliferative disease: outcomes and prognostic factors in the modern era. *J Clin Oncol* 2010;28:1038-1046. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20085936>.

291. Penn I, Porat G. Central nervous system lymphomas in organ allograft recipients. *Transplantation* 1995;59:240-244. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7839447>.

292. Leblond V, Dhedin N, Mamzer Bruneel MF, et al. Identification of prognostic factors in 61 patients with posttransplantation

lymphoproliferative disorders. *J Clin Oncol* 2001;19:772-778. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11157030>.

293. Nalesnik MA. Clinicopathologic characteristics of post-transplant lymphoproliferative disorders. *Recent Results Cancer Res* 2002;159:9-18. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11785849>.

294. Curtis RE, Travis LB, Rowlings PA, et al. Risk of lymphoproliferative disorders after bone marrow transplantation: a multi-institutional study. *Blood* 1999;94:2208-2216. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10498590>.

295. Jacobson CA, LaCasce AS. Lymphoma: risk and response after solid organ transplant. *Oncology (Williston Park)* 2010;24:936-944. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21138175>.

296. Wagner HJ, Rooney CM, Heslop HE. Diagnosis and treatment of posttransplantation lymphoproliferative disease after hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant* 2002;8:1-8. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11846351>.

297. Abrey LE, Batchelor TT, Ferreri AJM, et al. Report of an international workshop to standardize baseline evaluation and response criteria for primary CNS lymphoma. *J Clin Oncol* 2005;23:5034-5043. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15955902>.

298. Batchelor T, Carson K, O'Neill A, et al. Treatment of primary CNS lymphoma with methotrexate and deferred radiotherapy: a report of NABTT 96-07. *J Clin Oncol* 2003;21:1044-1049. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12637469>.

299. Chamberlain MC, Johnston SK. High-dose methotrexate and rituximab with deferred radiotherapy for newly diagnosed primary B-cell CNS lymphoma. *Neuro Oncol* 2010;12:736-744. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20511181>.

300. DeAngelis LM, Seiferheld W, Schold SC, et al. Combination chemotherapy and radiotherapy for primary central nervous system lymphoma: Radiation Therapy Oncology Group Study 93-10. *J Clin Oncol*





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

2002;20:4643-4648. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12488408>.

301. Gavrilovic IT, Hormigo A, Yahalom J, et al. Long-term follow-up of high-dose methotrexate-based therapy with and without whole brain irradiation for newly diagnosed primary CNS lymphoma. *J Clin Oncol* 2006;24:4570-4574. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17008697>.

302. Shah GD, Yahalom J, Correa DD, et al. Combined immunotherapy with reduced whole-brain radiotherapy for newly diagnosed primary CNS lymphoma. *J Clin Oncol* 2007;25:4730-4735. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17947720>.

303. Wieduwilt MJ, Valles F, Issa S, et al. Immunotherapy with intensive consolidation for primary CNS lymphoma: a pilot study and prognostic assessment by diffusion-weighted MRI. *Clin Cancer Res* 2012;18:1146-1155. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22228634>.

304. Gregory G, Arumugaswamy A, Leung T, et al. Rituximab is associated with improved survival for aggressive B cell CNS lymphoma. *Neuro Oncol* 2013;15:1068-1073. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23502429>.

305. Holdhoff M, Ambady P, Abdelaziz A, et al. High-dose methotrexate with or without rituximab in newly diagnosed primary CNS lymphoma. *Neurology* 2014;83:235-239. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24928128>.

306. Ly KI, Crew LL, Graham CA, Mrugala MM. Primary central nervous system lymphoma treated with high-dose methotrexate and rituximab: A single-institution experience. *Oncol Lett* 2016;11:3471-3476. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27123138>.

307. Omuro A, Correa DD, DeAngelis LM, et al. R-MPV followed by high-dose chemotherapy with TBC and autologous stem-cell transplant for newly diagnosed primary CNS lymphoma. *Blood* 2015;125:1403-1410. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25568347>.

308. Glass J, Won M, Schultz CJ, et al. Phase I and II study of induction chemotherapy with methotrexate, rituximab, and temozolomide, followed by whole-brain radiotherapy and postirradiation temozolomide for primary CNS lymphoma: NRG Oncology RTOG 0227. *J Clin Oncol* 2016;34:1620-1625. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27022122>.

309. Song Y, Wen Y, Xue W, et al. Effect of rituximab on primary central nervous system lymphoma: a meta-analysis. *Int J Hematol* 2017;106:612-621. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28900847>.

310. Ferreri AJ, Reni M, Foppoli M, et al. High-dose cytarabine plus high-dose methotrexate versus high-dose methotrexate alone in patients with primary CNS lymphoma: a randomised phase 2 trial. *Lancet* 2009;374:1512-1520. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19767089>.

311. Bromberg JEC, Issa S, Bakunina K, et al. Rituximab in patients with primary CNS lymphoma (HOVON 105/ALLG NHL 24): a randomised, open-label, phase 3 intergroup study. *Lancet Oncol* 2019;20:216-228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30630772>.

312. Ferreri AJ, Reni M, Pasini F, et al. A multicenter study of treatment of primary CNS lymphoma. *Neurology* 2002;58:1513-1520. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/12034789>.

313. Schulz H, Pels H, Schmidt-Wolf I, et al. Intraventricular treatment of relapsed central nervous system lymphoma with the anti-CD20 antibody rituximab. *Haematologica* 2004;89:753-754. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15194546>.

314. Abrey LE, Moskowitz CH, Mason WP, et al. Intensive methotrexate and cytarabine followed by high-dose chemotherapy with autologous stem-cell rescue in patients with newly diagnosed primary CNS lymphoma: an intent-to-treat analysis. *J Clin Oncol* 2003;21:4151-4156. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14615443>.

315. Montemurro M, Kiefer T, Schuler F, et al. Primary central nervous system lymphoma treated with high-dose methotrexate, high-dose busulfan/thiotepa, autologous stem-cell transplantation and response-



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

adapted whole-brain radiotherapy: results of the multicenter Ostdeutsche Studiengruppe Hamato-Onkologie OSHO-53 phase II study. *Ann Oncol* 2007;18:665-671. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17185743>.

316. Colombat P, Lemevel A, Bertrand P, et al. High-dose chemotherapy with autologous stem cell transplantation as first-line therapy for primary CNS lymphoma in patients younger than 60 years: a multicenter phase II study of the GOELAMS group. *Bone Marrow Transplant* 2006;38:417-420. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16951691>.

317. Illerhaus G, Muller F, Feuerhake F, et al. High-dose chemotherapy and autologous stem-cell transplantation without consolidating radiotherapy as first-line treatment for primary lymphoma of the central nervous system. *Haematologica* 2008;93:147-148. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18166803>.

318. Cote GM, Hochberg EP, Muzikansky A, et al. Autologous stem cell transplantation with thiotepa, busulfan, and cyclophosphamide (TBC) conditioning in patients with CNS involvement by non-Hodgkin lymphoma. *Biol Blood Marrow Transplant* 2012;18:76-83. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21749848>.

319. Kasenda B, Schorb E, Fritsch K, et al. Prognosis after high-dose chemotherapy followed by autologous stem-cell transplantation as first-line treatment in primary CNS lymphoma--a long-term follow-up study. *Ann Oncol* 2012;23:2670-2675. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22473593>.

320. DeFilipp Z, Li S, El-Jawahri A, et al. High-dose chemotherapy with thiotepa, busulfan, and cyclophosphamide and autologous stem cell transplantation for patients with primary central nervous system lymphoma in first complete remission. *Cancer* 2017;123:3073-3079. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28369839>.

321. Ferreri AJM, Cwynarski K, Pulczynski E, et al. Whole-brain radiotherapy or autologous stem-cell transplantation as consolidation strategies after high-dose methotrexate-based chemoimmunotherapy in patients with primary CNS lymphoma: results of the second randomisation

of the International Extranodal Lymphoma Study Group-32 phase 2 trial. *Lancet Haematol* 2017;4:e510-e523. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29054815>.

322. Widemann BC, Balis FM, Kim A, et al. Glucarpidase, leucovorin, and thymidine for high-dose methotrexate-induced renal dysfunction: clinical and pharmacologic factors affecting outcome. *J Clin Oncol* 2010;28:3979-3986. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20679598>.

323. Widemann BC, Balis FM, Kempf-Bielack B, et al. High-dose methotrexate-induced nephrotoxicity in patients with osteosarcoma. *Cancer* 2004;100:2222-2232. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15139068>.

324. Abrey LE, Yahalom J, DeAngelis LM. Treatment for primary CNS lymphoma: the next step. *J Clin Oncol* 2000;18:3144-3150. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10963643>.

325. Bessell EM, Lopez-Guillermo A, Villa S, et al. Importance of radiotherapy in the outcome of patients with primary CNS lymphoma: an analysis of the CHOD/BVAM regimen followed by two different radiotherapy treatments. *J Clin Oncol* 2002;20:231-236. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11773174>.

326. Gerstner ER, Carson KA, Grossman SA, Batchelor TT. Long-term outcome in PCNSL patients treated with high-dose methotrexate and deferred radiation. *Neurology* 2008;70:401-402. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18227422>.

327. Hoang-Xuan K, Taillandier L, Chinot O, et al. Chemotherapy alone as initial treatment for primary CNS lymphoma in patients older than 60 years: a multicenter phase II study (26952) of the European Organization for Research and Treatment of Cancer Brain Tumor Group. *J Clin Oncol* 2003;21:2726-2731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12860951>.

328. Pels H, Schmidt-Wolf IG, Glasmacher A, et al. Primary central nervous system lymphoma: results of a pilot and phase II study of systemic and intraventricular chemotherapy with deferred radiotherapy. *J*



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Clin Oncol 2003;21:4489-4495. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/14597744>.

329. Illerhaus G, Marks R, Muller F, et al. High-dose methotrexate combined with procarbazine and CCNU for primary CNS lymphoma in the elderly: results of a prospective pilot and phase II study. Ann Oncol 2009;20:319-325. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18953065>.

330. Juergens A, Pels H, Rogowski S, et al. Long-term survival with favorable cognitive outcome after chemotherapy in primary central nervous system lymphoma. Ann Neurol 2010;67:182-189. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/20225195>.

331. Rubenstein JL, Hsi ED, Johnson JL, et al. Intensive chemotherapy and immunotherapy in patients with newly diagnosed primary CNS lymphoma: CALGB 50202 (Alliance 50202). J Clin Oncol 2013;31:3061-3068. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23569323>.

332. Soussain C, Hoang-Xuan K, Taillandier L, et al. Intensive chemotherapy followed by hematopoietic stem-cell rescue for refractory and recurrent primary CNS and intraocular lymphoma: Societe Francaise de Greffe de Moelle Osseuse-Therapie Cellulaire. J Clin Oncol 2008;26:2512-2518. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18413641>.

333. Kasenda B, Ihorst G, Schroers R, et al. High-dose chemotherapy with autologous haematopoietic stem cell support for relapsed or refractory primary CNS lymphoma: a prospective multicentre trial by the German Cooperative PCNSL study group. Leukemia 2017;31:2623-2629. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28559537>.

334. Illerhaus G, Kasenda B, Ihorst G, et al. High-dose chemotherapy with autologous haemopoietic stem cell transplantation for newly diagnosed primary CNS lymphoma: a prospective, single-arm, phase 2 trial. Lancet Haematol 2016;3:e388-397. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/27476790>.

335. Schorb E, Finke J, Ferreri AJ, et al. High-dose chemotherapy and autologous stem cell transplant compared with conventional chemotherapy for consolidation in newly diagnosed primary CNS lymphoma--a randomized phase III trial (MATRix). BMC Cancer 2016;16:282. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/27098429>.

336. Houillier C, Taillandier L, Dureau S, et al. Radiotherapy or autologous stem-cell transplantation for primary CNS lymphoma in patients 60 years of age and younger: results of the Intergroup ANOCEF-GOELAMS randomized phase II PRECIS study. J Clin Oncol 2019;37:823-833. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30785830>.

337. Plotkin SR, Betensky RA, Hochberg FH, et al. Treatment of relapsed central nervous system lymphoma with high-dose methotrexate. Clin Cancer Res 2004;10:5643-5646. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/15355887>.

338. Grommes C, Tang SS, Wolfe J, et al. Phase 1b trial of an ibrutinib-based combination therapy in recurrent/refractory CNS lymphoma. Blood 2019;133:436-445. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/30567753>.

339. Grommes C, Pastore A, Palaskas N, et al. Ibrutinib unmasks critical role of bruton tyrosine kinase in primary CNS lymphoma. Cancer Discov 2017;7:1018-1029. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/28619981>.

340. Soussain C, Choquet S, Blonski M, et al. Ibrutinib monotherapy for relapse or refractory primary CNS lymphoma and primary vitreoretinal lymphoma: final analysis of the phase II 'proof-of-concept' iLOC study by the Lymphoma study association (LYSA) and the French oculo-cerebral lymphoma (LOC) network. Eur J Cancer 2019;117:121-130. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/31279304>.

341. Batchelor TT, Grossman SA, Mikkelsen T, et al. Rituximab monotherapy for patients with recurrent primary CNS lymphoma. Neurology 2011;76:929-930. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/21383331>.





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

342. Nayak L, Abrey LE, Drappatz J, et al. Multicenter phase II study of rituximab and temozolomide in recurrent primary central nervous system lymphoma. *Leuk Lymphoma* 2013;54:58-61. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22656234>.

343. Reni M, Zaja F, Mason W, et al. Temozolomide as salvage treatment in primary brain lymphomas. *Br J Cancer* 2007;96:864-867. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17325700>.

344. Makino K, Nakamura H, Hide T, Kuratsu J. Salvage treatment with temozolomide in refractory or relapsed primary central nervous system lymphoma and assessment of the MGMT status. *J Neurooncol* 2012;106:155-160. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21720808>.

345. Enting RH, Demopoulos A, DeAngelis LM, Abrey LE. Salvage therapy for primary CNS lymphoma with a combination of rituximab and temozolomide. *Neurology* 2004;63:901-903. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15365145>.

346. Rubenstein JL, Geng H, Fraser EJ, et al. Phase 1 investigation of lenalidomide/rituximab plus outcomes of lenalidomide maintenance in relapsed CNS lymphoma. *Blood Adv* 2018;2:1595-1607. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29986852>.

347. DeAngelis LM, Kreis W, Chan K, et al. Pharmacokinetics of ara-C and ara-U in plasma and CSF after high-dose administration of cytosine arabinoside. *Cancer Chemother Pharmacol* 1992;29:173-177. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1733548>.

348. Tun HW, Johnston PB, DeAngelis LM, et al. Phase 1 study of pomalidomide and dexamethasone for relapsed/refractory primary CNS or vitreoretinal lymphoma. *Blood* 2018;132:2240-2248. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30262659>.

349. Raizer JJ, Rademaker A, Evens AM, et al. Pemetrexed in the treatment of relapsed/refractory primary central nervous system lymphoma. *Cancer* 2012;118:3743-3748. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22179954>.

350. DeAngelis LM, Yahalom J, Thaler HT, Kher U. Combined modality therapy for primary CNS lymphoma. *J Clin Oncol* 1992;10:635-643. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1548527>.

351. Fisher B, Seiferheld W, Schultz C, et al. Secondary analysis of Radiation Therapy Oncology Group study (RTOG) 9310: an intergroup phase II combined modality treatment of primary central nervous system lymphoma. *J Neurooncol* 2005;74:201-205. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16193393>.

352. Nelson DF, Martz KL, Bonner H, et al. Non-Hodgkin's lymphoma of the brain: can high dose, large volume radiation therapy improve survival? Report on a prospective trial by the Radiation Therapy Oncology Group (RTOG): RTOG 8315. *Int J Radiat Oncol Biol Phys* 1992;23:9-17. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1572835>.

353. Schultz C, Scott C, Sherman W, et al. Preirradiation chemotherapy with cyclophosphamide, doxorubicin, vincristine, and dexamethasone for primary CNS lymphomas: initial report of radiation therapy oncology group protocol 88-06. *J Clin Oncol* 1996;14:556-564. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8636771>.

354. Poortmans PM, Kluin-Nelemans HC, Haaxma-Reiche H, et al. High-dose methotrexate-based chemotherapy followed by consolidating radiotherapy in non-AIDS-related primary central nervous system lymphoma: European Organization for Research and Treatment of Cancer Lymphoma Group Phase II Trial 20962. *J Clin Oncol* 2003;21:4483-4488. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14597741>.

355. Ghesquieres H, Ferlay C, Sebban C, et al. Long-term follow-up of an age-adapted C5R protocol followed by radiotherapy in 99 newly diagnosed primary CNS lymphomas: a prospective multicentric phase II study of the Groupe d'Etude des Lymphomes de l'Adulte (GELA). *Ann Oncol* 2010;21:842-850. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19914958>.

356. Thiel E, Korfel A, Martus P, et al. High-dose methotrexate with or without whole brain radiotherapy for primary CNS lymphoma (G-PCNSL-SG-1): a phase 3, randomised, non-inferiority trial. *Lancet Oncol*



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

2010;11:1036-1047. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20970380>.

357. Morris PG, Correa DD, Yahalom J, et al. Rituximab, methotrexate, procarbazine, and vincristine followed by consolidation reduced-dose whole-brain radiotherapy and cytarabine in newly diagnosed primary CNS lymphoma: final results and long-term outcome. *J Clin Oncol* 2013;31:3971-3979. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24101038>.

358. Nguyen PL, Chakravarti A, Finkelstein DM, et al. Results of whole-brain radiation as salvage of methotrexate failure for immunocompetent patients with primary CNS lymphoma. *J Clin Oncol* 2005;23:1507-1513. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15735126>.

359. Coulon A, Lafitte F, Hoang-Xuan K, et al. Radiographic findings in 37 cases of primary CNS lymphoma in immunocompetent patients. *Eur Radiol* 2002;12:329-340. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11870430>.

360. Mohile NA, Deangelis LM, Abrey LE. The utility of body FDG PET in staging primary central nervous system lymphoma. *Neuro Oncol* 2008;10:223-228. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18287338>.

361. Ferreri AJ, Blay JY, Reni M, et al. Prognostic scoring system for primary CNS lymphomas: the International Extranodal Lymphoma Study Group experience. *J Clin Oncol* 2003;21:266-272. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/12525518>.

362. Jahr G, Broi MD, Holte H, Jr., et al. Evaluation of Memorial Sloan-Kettering Cancer Center and International Extranodal Lymphoma Study Group prognostic scoring systems to predict overall survival in intracranial primary CNS lymphoma. *Brain Behav* 2018;8:e00928. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29541540>.

363. Doolittle ND, Korfel A, Lubow MA, et al. Long-term cognitive function, neuroimaging, and quality of life in primary CNS lymphoma. *Neurology*

2013;81:84-92. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23685932>.

364. Schellinger KA, Propp JM, Villano JL, McCarthy BJ. Descriptive epidemiology of primary spinal cord tumors. *J Neurooncol* 2008;87:173-179. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18084720>.

365. Grimm S, Chamberlain MC. Adult primary spinal cord tumors. *Expert Rev Neurother* 2009;9:1487-1495. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19831838>.

366. Gezen F, Kahraman S, Canakci Z, Beduk A. Review of 36 cases of spinal cord meningioma. *Spine (Phila Pa 1976)* 2000;25:727-731. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10752106>.

367. Solero CL, Fornari M, Giombini S, et al. Spinal meningiomas: review of 174 operated cases. *Neurosurgery* 1989;25:153-160. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/2671779>.

368. Volpp PB, Han K, Kagan AR, Tome M. Outcomes in treatment for intradural spinal cord ependymomas. *Int J Radiat Oncol Biol Phys* 2007;69:1199-1204. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17689025>.

369. Yang S, Yang X, Hong G. Surgical treatment of one hundred seventy-four intramedullary spinal cord tumors. *Spine (Phila Pa 1976)* 2009;34:2705-2710. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19910775>.

370. Raco A, Esposito V, Lenzi J, et al. Long-term follow-up of intramedullary spinal cord tumors: a series of 202 cases. *Neurosurgery* 2005;56:972-981; discussion 972-981. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15854245>.

371. Benes V, 3rd, Barsa P, Benes V, Jr., Suchomel P. Prognostic factors in intramedullary astrocytomas: a literature review. *Eur Spine J* 2009;18:1397-1422. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19562388>.





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

372. Milano MT, Johnson MD, Sul J, et al. Primary spinal cord glioma: a Surveillance, Epidemiology, and End Results database study. *J Neurooncol* 2010;98:83-92. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19898743>.

373. Sgouros S, Malluci CL, Jackowski A. Spinal ependymomas--the value of postoperative radiotherapy for residual disease control. *Br J Neurosurg* 1996;10:559-566. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9115651>.

374. Lee TT, Gromelski EB, Green BA. Surgical treatment of spinal ependymoma and post-operative radiotherapy. *Acta Neurochir (Wien)* 1998;140:309-313. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9689321>.

375. Akyurek S, Chang EL, Yu TK, et al. Spinal myxopapillary ependymoma outcomes in patients treated with surgery and radiotherapy at M.D. Anderson Cancer Center. *J Neurooncol* 2006;80:177-183. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16648988>.

376. Pica A, Miller R, Villa S, et al. The results of surgery, with or without radiotherapy, for primary spinal myxopapillary ependymoma: a retrospective study from the rare cancer network. *Int J Radiat Oncol Biol Phys* 2009;74:1114-1120. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19250760>.

377. Ryu SI, Kim DH, Chang SD. Stereotactic radiosurgery for hemangiomas and ependymomas of the spinal cord. *Neurosurg Focus* 2003;15:E10. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15323467>.

378. Gerszten PC, Burton SA, Ozhasoglu C, et al. Radiosurgery for benign intradural spinal tumors. *Neurosurgery* 2008;62:887-895; discussion 895-886. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18496194>.

379. Dodd RL, Ryu MR, Kamnerdsupaphon P, et al. CyberKnife radiosurgery for benign intradural extramedullary spinal tumors. *Neurosurgery* 2006;58:674-685; discussion 674-685. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16575331>.

380. Garces-Ambrossi GL, McGirt MJ, Mehta VA, et al. Factors associated with progression-free survival and long-term neurological outcome after resection of intramedullary spinal cord tumors: analysis of 101 consecutive cases. *J Neurosurg Spine* 2009;11:591-599. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19929363>.

381. Alliance TV. The VHL Handbook: What You Need to Know About VHL (ed 5th ed.). Boston, MA: VHL Alliance; 2015.

382. Surawicz TS, McCarthy BJ, Kupelian V, et al. Descriptive epidemiology of primary brain and CNS tumors: results from the Central Brain Tumor Registry of the United States, 1990-1994. *Neuro Oncol* 1999;1:14-25. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11554386>.

383. Mahmood A, Caccamo DV, Tomecek FJ, Malik GM. Atypical and malignant meningiomas: a clinicopathological review. *Neurosurgery* 1993;33:955-963. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8134008>.

384. Vernooij MW, Ikram MA, Tanghe HL, et al. Incidental findings on brain MRI in the general population. *N Engl J Med* 2007;357:1821-1828. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17978290>.

385. Lieu AS, Howng SL. Intracranial meningiomas and epilepsy: incidence, prognosis and influencing factors. *Epilepsy Res* 2000;38:45-52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10604605>.

386. Campbell BA, Jhamb A, Maguire JA, et al. Meningiomas in 2009: controversies and future challenges. *Am J Clin Oncol* 2009;32:73-85. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19194129>.

387. Nathoo N, Ugokwe K, Chang AS, et al. The role of 111indium-octreotide brain scintigraphy in the diagnosis of cranial, dural-based meningiomas. *J Neurooncol* 2007;81:167-174. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16850106>.

388. Nyuyki F, Plotkin M, Graf R, et al. Potential impact of (68)Ga-DOTATOC PET/CT on stereotactic radiotherapy planning of meningiomas.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Eur J Nucl Med Mol Imaging 2010;37:310-318. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/19763565>.

389. Gehler B, Paulsen F, Oksuz MO, et al. [68Ga]-DOTATOC-PET/CT for meningioma IMRT treatment planning. Radiat Oncol 2009;4:56. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19922642>.

390. Nakamura M, Roser F, Michel J, et al. The natural history of incidental meningiomas. Neurosurgery 2003;53:62-70; discussion 70-61. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12823874>.

391. Olivero WC, Lister JR, Elwood PW. The natural history and growth rate of asymptomatic meningiomas: a review of 60 patients. J Neurosurg 1995;83:222-224. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7616265>.

392. Stafford SL, Perry A, Suman VJ, et al. Primarily resected meningiomas: outcome and prognostic factors in 581 Mayo Clinic patients, 1978 through 1988. Mayo Clin Proc 1998;73:936-942. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9787740>.

393. Mahmood A, Qureshi NH, Malik GM. Intracranial meningiomas: analysis of recurrence after surgical treatment. Acta Neurochir (Wien) 1994;126:53-58. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8042555>.

394. Mathiesen T, Lindquist C, Kihlstrom L, Karlsson B. Recurrence of cranial base meningiomas. Neurosurgery 1996;39:2-7; discussion 8-9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8805134>.

395. Perry A, Stafford SL, Scheithauer BW, et al. Meningioma grading: an analysis of histologic parameters. Am J Surg Pathol 1997;21:1455-1465. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9414189>.

396. Simpson D. The recurrence of intracranial meningiomas after surgical treatment. J Neurol Neurosurg Psychiatry 1957;20:22-39. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/13406590>.

397. Condra KS, Buatti JM, Mendenhall WM, et al. Benign meningiomas: primary treatment selection affects survival. Int J Radiat Oncol Biol Phys 1997;39:427-436. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9308947>.

398. Soyuer S, Chang EL, Selek U, et al. Radiotherapy after surgery for benign cerebral meningioma. Radiother Oncol 2004;71:85-90. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15066300>.

399. Pirzkall A, Debus J, Haering P, et al. Intensity modulated radiotherapy (IMRT) for recurrent, residual, or untreated skull-base meningiomas: preliminary clinical experience. Int J Radiat Oncol Biol Phys 2003;55:362-372. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12527049>.

400. Aghi MK, Carter BS, Cosgrove GR, et al. Long-term recurrence rates of atypical meningiomas after gross total resection with or without postoperative adjuvant radiation. Neurosurgery 2009;64:56-60; discussion 60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19145156>.

401. Hug EB, Devries A, Thornton AF, et al. Management of atypical and malignant meningiomas: role of high-dose, 3D-conformal radiation therapy. J Neurooncol 2000;48:151-160. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11083080>.

402. Rogers CL, Won M, Vogelbaum MA, et al. High-risk meningioma: initial outcomes from NRG Oncology/RTOG 0539. Int J Radiat Oncol Biol Phys 2020;106:790-799. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31786276>.

403. Selch MT, Ahn E, Laskari A, et al. Stereotactic radiotherapy for treatment of cavernous sinus meningiomas. Int J Radiat Oncol Biol Phys 2004;59:101-111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15093905>.

404. Pollock BE, Stafford SL, Utter A, et al. Stereotactic radiosurgery provides equivalent tumor control to Simpson Grade 1 resection for patients with small- to medium-size meningiomas. Int J Radiat Oncol Biol



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

Phys 2003;55:1000-1005. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12605979>.

405. Kondziolka D, Mathieu D, Lunsford LD, et al. Radiosurgery as definitive management of intracranial meningiomas. Neurosurgery 2008;62:53-58; discussion 58-60. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18300891>.

406. Compter I, Zaugg K, Houben RM, et al. High symptom improvement and local tumor control using stereotactic radiotherapy when given early after diagnosis of meningioma. A multicentre study. Strahlenther Onkol 2012;188:887-893. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22961046>.

407. Debus J, Wuendrich M, Pirzkall A, et al. High efficacy of fractionated stereotactic radiotherapy of large base-of-skull meningiomas: long-term results. J Clin Oncol 2001;19:3547-3553. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11481362>.

408. Kaley T, Barani I, Chamberlain M, et al. Historical benchmarks for medical therapy trials in surgery- and radiation-refractory meningioma: a RANO review. Neuro Oncol 2014;16:829-840. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24500419>.

409. Chamberlain MC, Glantz MJ, Fadul CE. Recurrent meningioma: salvage therapy with long-acting somatostatin analogue. Neurology 2007;69:969-973. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17785665>.

410. Johnson DR, Kimmel DW, Burch PA, et al. Phase II study of subcutaneous octreotide in adults with recurrent or progressive meningioma and meningeal hemangiopericytoma. Neuro Oncol 2011;13:530-535. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21558077>.

411. Kaley TJ, Wen P, Schiff D, et al. Phase II trial of sunitinib for recurrent and progressive atypical and anaplastic meningioma. Neuro Oncol 2015;17:116-121. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25100872>.

412. Lou E, Sumrall AL, Turner S, et al. Bevacizumab therapy for adults with recurrent/progressive meningioma: a retrospective series. J Neurooncol 2012;109:63-70. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22535433>.

413. Nayak L, Iwamoto FM, Rudnick JD, et al. Atypical and anaplastic meningiomas treated with bevacizumab. J Neurooncol 2012;109:187-193. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22544653>.

414. Shih KC, Chowdhary S, Rosenblatt P, et al. A phase II trial of bevacizumab and everolimus as treatment for patients with refractory, progressive intracranial meningioma. J Neurooncol 2016;129:281-288. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27311730>.

415. Barnholtz-Sloan JS, Sloan AE, Davis FG, et al. Incidence proportions of brain metastases in patients diagnosed (1973 to 2001) in the Metropolitan Detroit Cancer Surveillance System. J Clin Oncol 2004;22:2865-2872. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15254054>.

416. Schouten LJ, Rutten J, Huveneers HA, Twijnstra A. Incidence of brain metastases in a cohort of patients with carcinoma of the breast, colon, kidney, and lung and melanoma. Cancer 2002;94:2698-2705. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12173339>.

417. Fox BD, Cheung VJ, Patel AJ, et al. Epidemiology of metastatic brain tumors. Neurosurg Clin N Am 2011;22:1-6, v. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21109143>.

418. Gould J. Breaking down the epidemiology of brain cancer. Nature 2018;561:S40-S41. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30258156>.

419. Maher EA, Mietz J, Arteaga CL, et al. Brain metastasis: opportunities in basic and translational research. Cancer Res 2009;69:6015-6020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19638593>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

420. Lin NU, Bellon JR, Winer EP. CNS metastases in breast cancer. *J Clin Oncol* 2004;22:3608-3617. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15337811>.

421. Eichler AF, Loeffler JS. Multidisciplinary management of brain metastases. *Oncologist* 2007;12:884-898. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17673619>.

422. Mahajan A, Ahmed S, McAleer MF, et al. Post-operative stereotactic radiosurgery versus observation for completely resected brain metastases: a single-centre, randomised, controlled, phase 3 trial. *Lancet Oncol* 2017;18:1040-1048. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28687375>.

423. Churilla TM, Chowdhury IH, Handorf E, et al. Comparison of local control of brain metastases with stereotactic radiosurgery vs surgical resection: a secondary analysis of a randomized clinical trial. *JAMA Oncol* 2019;5:243-247. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30419088>.

424. Ewend MG, Morris DE, Carey LA, et al. Guidelines for the initial management of metastatic brain tumors: role of surgery, radiosurgery, and radiation therapy. *J Natl Compr Canc Netw* 2008;6:505-513; quiz 514. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18492462>.

425. Patchell RA, Tibbs PA, Walsh JW, et al. A randomized trial of surgery in the treatment of single metastases to the brain. *N Engl J Med* 1990;322:494-500. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2405271>.

426. Vecht CJ, Haaxma-Reiche H, Noordijk EM, et al. Treatment of single brain metastasis: radiotherapy alone or combined with neurosurgery? *Ann Neurol* 1993;33:583-590. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8498838>.

427. Mintz AH, Kestle J, Rathbone MP, et al. A randomized trial to assess the efficacy of surgery in addition to radiotherapy in patients with a single cerebral metastasis. *Cancer* 1996;78:1470-1476. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8839553>.

428. Suh JH. Stereotactic radiosurgery for the management of brain metastases. *N Engl J Med* 2010;362:1119-1127. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20335588>.

429. Aoyama H, Shirato H, Tago M, et al. Stereotactic radiosurgery plus whole-brain radiation therapy vs stereotactic radiosurgery alone for treatment of brain metastases: a randomized controlled trial. *JAMA* 2006;295:2483-2491. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16757720>.

430. Brown PD, Jaeckle K, Ballman KV, et al. Effect of radiosurgery alone vs radiosurgery with whole brain radiation therapy on cognitive function in patients with 1 to 3 brain metastases: a randomized clinical trial. *JAMA* 2016;316:401-409. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27458945>.

431. Chang EL, Wefel JS, Hess KR, et al. Neurocognition in patients with brain metastases treated with radiosurgery or radiosurgery plus whole-brain irradiation: a randomised controlled trial. *Lancet Oncol* 2009;10:1037-1044. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19801201>.

432. Kocher M, Soffiotti R, Abacioglu U, et al. Adjuvant whole-brain radiotherapy versus observation after radiosurgery or surgical resection of one to three cerebral metastases: results of the EORTC 22952-26001 study. *J Clin Oncol* 2011;29:134-141. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21041710>.

433. Yamamoto M, Serizawa T, Shuto T, et al. Stereotactic radiosurgery for patients with multiple brain metastases (JLKG0901): a multi-institutional prospective observational study. *Lancet Oncol* 2014;15:387-395. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24621620>.

434. Bhatnagar AK, Flickinger JC, Kondziolka D, Lunsford LD. Stereotactic radiosurgery for four or more intracranial metastases. *Int J Radiat Oncol Biol Phys* 2006;64:898-903. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16338097>.





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

435. Banfill KE, Bownes PJ, St Clair SE, et al. Stereotactic radiosurgery for the treatment of brain metastases: impact of cerebral disease burden on survival. *Br J Neurosurg* 2012;26:674-678. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22747250>.

436. Chang WS, Kim HY, Chang JW, et al. Analysis of radiosurgical results in patients with brain metastases according to the number of brain lesions: is stereotactic radiosurgery effective for multiple brain metastases? *J Neurosurg* 2010;113 Suppl:73-78. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21121789>.

437. Farris M, McTyre ER, Cramer CK, et al. Brain metastasis velocity: a novel prognostic metric predictive of overall survival and freedom from whole-brain radiation therapy after distant brain failure following upfront radiosurgery alone. *Int J Radiat Oncol Biol Phys* 2017;98:131-141. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28586952>.

438. Karlsson B, Hanssens P, Wolff R, et al. Thirty years' experience with Gamma Knife surgery for metastases to the brain. *J Neurosurg* 2009;111:449-457. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19199505>.

439. Kased N, Binder DK, McDermott MW, et al. Gamma Knife radiosurgery for brain metastases from primary breast cancer. *Int J Radiat Oncol Biol Phys* 2009;75:1132-1140. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19345514>.

440. Rusthoven CG, Yamamoto M, Bernhardt D, et al. Evaluation of first-line radiosurgery vs whole-brain radiotherapy for small cell lung cancer brain metastases: the FIRE-SCLC Cohort Study. *JAMA Oncol* 2020;6:1028-1037. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32496550>.

441. Hanson PW, Elaimy AL, Lamoreaux WT, et al. A concise review of the efficacy of stereotactic radiosurgery in the management of melanoma and renal cell carcinoma brain metastases. *World J Surg Oncol* 2012;10:176. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22931379>.

442. Hunter GK, Suh JH, Reuther AM, et al. Treatment of five or more brain metastases with stereotactic radiosurgery. *Int J Radiat Oncol Biol Phys* 2012;83:1394-1398. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22209150>.

443. Minniti G, Esposito V, Clarke E, et al. Stereotactic radiosurgery in elderly patients with brain metastases. *J Neurooncol* 2013;111:319-325. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23187817>.

444. Higuchi Y, Yamamoto M, Serizawa T, et al. Stereotactic radiosurgery in elderly patients with brain metastases: comparison with non-elderly patients using database of a multi-institutional prospective observational study (JLGK0901-Elderly). *J Neurooncol* 2019;144:393-402. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31338786>.

445. Raman S, Mou B, Hsu F, et al. Whole brain radiotherapy versus stereotactic radiosurgery in poor-prognosis patients with one to 10 brain metastases: a randomised feasibility study. *Clin Oncol (R Coll Radiol)* 2020;32:442-451. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32085923>.

446. Chen L, Shen C, Redmond KJ, et al. Use of stereotactic radiosurgery in elderly and very elderly patients with brain metastases to limit toxicity associated with whole brain radiation therapy. *Int J Radiat Oncol Biol Phys* 2017;98:939-947. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28602418>.

447. Shaw E, Scott C, Souhami L, et al. Single dose radiosurgical treatment of recurrent previously irradiated primary brain tumors and brain metastases: final report of RTOG protocol 90-05. *Int J Radiat Oncol Biol Phys* 2000;47:291-298. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10802351>.

448. Minniti G, Scaringi C, Paolini S, et al. Single-fraction versus multifraction (3 x 9 Gy) stereotactic radiosurgery for large (>2 cm) brain metastases: a comparative analysis of local control and risk of radiation-induced brain necrosis. *Int J Radiat Oncol Biol Phys* 2016;95:1142-1148. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27209508>.





449. Al-Omair A, Soliman H, Xu W, et al. Hypofractionated stereotactic radiotherapy in five daily fractions for post-operative surgical cavities in brain metastases patients with and without prior whole brain radiation. *Technol Cancer Res Treat* 2013;12:493-499. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23617283>.

450. Soltys SG, Seiger K, Modlin LA, et al. A phase I/II dose-escalation trial of 3-fraction stereotactic radiosurgery (SRS) for large resection cavities of brain metastases. *Int J Radiat Oncol Biol Phys* 2015;93:S38. Available at: <https://doi.org/10.1016/j.ijrobp.2015.07.093>.

451. Soliman H, Ruschin M, Angelov L, et al. Consensus contouring guidelines for postoperative completely resected cavity stereotactic radiosurgery for brain metastases. *Int J Radiat Oncol Biol Phys* 2018;100:436-442. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29157748>.

452. Akyurek S, Chang EL, Mahajan A, et al. Stereotactic radiosurgical treatment of cerebral metastases arising from breast cancer. *Am J Clin Oncol* 2007;30:310-314. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17551311>.

453. Noel G, Proudhon MA, Valery CA, et al. Radiosurgery for re-irradiation of brain metastasis: results in 54 patients. *Radiother Oncol* 2001;60:61-67. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11410305>.

454. Sheehan J, Kondziolka D, Flickinger J, Lunsford LD. Radiosurgery for patients with recurrent small cell lung carcinoma metastatic to the brain: outcomes and prognostic factors. *J Neurosurg* 2005;102 Suppl:247-254. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15662819>.

455. Caballero JA, Sneed PK, Lamborn KR, et al. Prognostic factors for survival in patients treated with stereotactic radiosurgery for recurrent brain metastases after prior whole brain radiotherapy. *Int J Radiat Oncol Biol Phys* 2012;83:303-309. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22079723>.

456. Brown PD, Ballman KV, Cerhan JH, et al. Postoperative stereotactic radiosurgery compared with whole brain radiotherapy for resected metastatic brain disease (NCCTG N107C/CEC.3): a multicentre, randomised, controlled, phase 3 trial. *Lancet Oncol* 2017;18:1049-1060. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28687377>.

457. Andrews DW, Scott CB, Sperduto PW, et al. Whole brain radiation therapy with or without stereotactic radiosurgery boost for patients with one to three brain metastases: phase III results of the RTOG 9508 randomised trial. *Lancet* 2004;363:1665-1672. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15158627>.

458. Tsao MN, Xu W, Wong RK, et al. Whole brain radiotherapy for the treatment of newly diagnosed multiple brain metastases. *Cochrane Database Syst Rev* 2018;1:CD003869. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29365347>.

459. Churilla TM, Handorf E, Collette S, et al. Whole brain radiotherapy after stereotactic radiosurgery or surgical resection among patients with one to three brain metastases and favorable prognoses: a secondary analysis of EORTC 22952-26001. *Ann Oncol* 2017;28:2588-2594. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28961826>.

460. Mulvenna P, Nankivell M, Barton R, et al. Dexamethasone and supportive care with or without whole brain radiotherapy in treating patients with non-small cell lung cancer with brain metastases unsuitable for resection or stereotactic radiotherapy (QUARTZ): results from a phase 3, non-inferiority, randomised trial. *Lancet* 2016;388:2004-2014. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27604504>.

461. Yomo S, Hayashi M. Is upfront stereotactic radiosurgery a rational treatment option for very elderly patients with brain metastases? A retrospective analysis of 106 consecutive patients age 80 years and older. *BMC Cancer* 2016;16:948. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27978813>.

462. Brown PD, Pugh S, Laack NN, et al. Memantine for the prevention of cognitive dysfunction in patients receiving whole-brain radiotherapy: a randomized, double-blind, placebo-controlled trial. *Neuro Oncol*



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

2013;15:1429-1437. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23956241>.

463. Gondi V, Pugh SL, Tome WA, et al. Preservation of memory with conformal avoidance of the hippocampal neural stem-cell compartment during whole-brain radiotherapy for brain metastases (RTOG 0933): a phase II multi-institutional trial. *J Clin Oncol* 2014;32:3810-3816. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25349290>.

464. Brown PD, Gondi V, Pugh S, et al. Hippocampal avoidance during whole-brain radiotherapy plus memantine for patients with brain metastases: phase III trial NRG Oncology CC001. *J Clin Oncol* 2020;38:1019-1029. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32058845>.

465. Patchell RA, Tibbs PA, Regine WF, et al. Postoperative radiotherapy in the treatment of single metastases to the brain: a randomized trial. *JAMA* 1998;280:1485-1489. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9809728>.

466. Soffietti R, Kocher M, Abacioglu UM, et al. A European Organisation for Research and Treatment of Cancer phase III trial of adjuvant whole-brain radiotherapy versus observation in patients with one to three brain metastases from solid tumors after surgical resection or radiosurgery: quality-of-life results. *J Clin Oncol* 2013;31:65-72. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23213105>.

467. Hong AM, Fogarty GB, Dolven-Jacobsen K, et al. Adjuvant whole-brain radiation therapy compared with observation after local treatment of melanoma brain metastases: a multicenter, randomized phase III trial. *J Clin Oncol* 2019;37:3132-3141. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31553661>.

468. Doebele RC, Drilon A, Paz-Ares L, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1-2 trials. *Lancet Oncol* 2020;21:271-282. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31838007>.

469. Hong DS, DuBois SG, Kummar S, et al. Larotrectinib in patients with TRK fusion-positive solid tumours: a pooled analysis of three phase 1/2 clinical trials. *Lancet Oncol* 2020;21:531-540. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32105622>.

470. Glitza Oliva IC, Schvartsman G, Tawbi H. Advances in the systemic treatment of melanoma brain metastases. *Ann Oncol* 2018;29:1509-1520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29790899>.

471. Turajlic S, Larkin J. Immunotherapy for melanoma metastatic to the brain. *N Engl J Med* 2018;379:789-790. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30134137>.

472. Long GV, Atkinson V, Lo S, et al. Combination nivolumab and ipilimumab or nivolumab alone in melanoma brain metastases: a multicentre randomised phase 2 study. *Lancet Oncol* 2018;19:672-681. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29602646>.

473. Tawbi HA, Forsyth PA, Algazi A, et al. Combined nivolumab and ipilimumab in melanoma metastatic to the brain. *N Engl J Med* 2018;379:722-730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30134131>.

474. Margolin K, Ernstoff MS, Hamid O, et al. Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial. *Lancet Oncol* 2012;13:459-465. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22456429>.

475. Goldberg SB, Gettinger SN, Mahajan A, et al. Pembrolizumab for patients with melanoma or non-small-cell lung cancer and untreated brain metastases: early analysis of a non-randomised, open-label, phase 2 trial. *Lancet Oncol* 2016;17:976-983. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27267608>.

476. Kluger HM, Chiang V, Mahajan A, et al. Long-term survival of patients with melanoma with active brain metastases treated with pembrolizumab on a phase II trial. *J Clin Oncol* 2019;37:52-60. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30407895>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

477. Davies MA, Saiag P, Robert C, et al. Dabrafenib plus trametinib in patients with BRAFV600-mutant melanoma brain metastases (COMBI-MB): a multicentre, multicohort, open-label, phase 2 trial. *Lancet Oncol* 2017;18:863-873. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28592387>.

478. McArthur GA, Maio M, Arance A, et al. Vemurafenib in metastatic melanoma patients with brain metastases: an open-label, single-arm, phase 2, multicentre study. *Ann Oncol* 2017;28:634-641. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27993793>.

479. Dummer R, Goldinger SM, Turtzsch CP, et al. Vemurafenib in patients with BRAF(V600) mutation-positive melanoma with symptomatic brain metastases: final results of an open-label pilot study. *Eur J Cancer* 2014;50:611-621. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24295639>.

480. Ascierto PA, McArthur GA, Dreno B, et al. Cobimetinib combined with vemurafenib in advanced BRAF(V600)-mutant melanoma (coBRIM): updated efficacy results from a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2016;17:1248-1260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27480103>.

481. Larkin J, Ascierto PA, Dreno B, et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med* 2014;371:1867-1876. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25265494>.

482. Holbrook K, Lutzky J, Davies MA, et al. Intracranial antitumor activity with encorafenib plus binimetinib in patients with melanoma brain metastases: a case series. *Cancer* 2020;126:523-530. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31658370>.

483. Rizvi NA, Mazieres J, Planchard D, et al. Activity and safety of nivolumab, an anti-PD-1 immune checkpoint inhibitor, for patients with advanced, refractory squamous non-small-cell lung cancer (CheckMate 063): a phase 2, single-arm trial. *Lancet Oncol* 2015;16:257-265. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25704439>.

484. Brahmer J, Reckamp KL, Baas P, et al. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med* 2015;373:123-135. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26028407>.

485. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med* 2015;373:1627-1639. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26412456>.

486. Goldman JW, Crino L, Vokes EE, et al. P2.36: Nivolumab (nivo) in patients (pts) with advanced (adv) NSCLC and central nervous system (CNS) metastases (mets). *J Thorac Oncol* 2016;11:S238-S239. Available at: [https://www.jto.org/article/S1556-0864\(16\)30816-4/fulltext](https://www.jto.org/article/S1556-0864(16)30816-4/fulltext).

487. Gauvain C, Vauleon E, Chouaid C, et al. Intracerebral efficacy and tolerance of nivolumab in non-small-cell lung cancer patients with brain metastases. *Lung Cancer* 2018;116:62-66. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29413052>.

488. Rangachari D, Yamaguchi N, VanderLaan PA, et al. Brain metastases in patients with EGFR-mutated or ALK-rearranged non-small-cell lung cancers. *Lung Cancer* 2015;88:108-111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25682925>.

489. Costa DB, Shaw AT, Ou SH, et al. Clinical experience with crizotinib in patients with advanced ALK-rearranged non-small-cell lung cancer and brain metastases. *J Clin Oncol* 2015;33:1881-1888. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25624436>.

490. Peters S, Camidge DR, Shaw AT, et al. Alectinib versus crizotinib in untreated ALK-positive non-small-cell lung cancers. *N Engl J Med* 2017;377:829-838. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28586279>.

491. Ou SI, Ahn JS, De Petris L, et al. Alectinib in crizotinib-refractory ALK-rearranged non-small-cell lung cancer: a phase II global study. *J Clin Oncol* 2016;34:661-668. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26598747>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

492. Shaw AT, Gandhi L, Gadgeel S, et al. Alectinib in ALK-positive, crizotinib-resistant, non-small-cell lung cancer: a single-group, multicentre, phase 2 trial. *Lancet Oncol* 2016;17:234-242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26708155>.

493. Gandhi L, Ou SI, Shaw AT, et al. Efficacy of alectinib in central nervous system metastases in crizotinib-resistant ALK-positive non-small-cell lung cancer: comparison of RECIST 1.1 and RANO-HGG criteria. *Eur J Cancer* 2017;82:27-33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28646771>.

494. Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib versus crizotinib in ALK-positive non-small-cell lung cancer. *N Engl J Med* 2018;379:2027-2039. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30280657>.

495. Kim DW, Tiseo M, Ahn MJ, et al. Brigatinib in patients with crizotinib-refractory anaplastic lymphoma kinase-positive non-small-cell lung cancer: a randomized, multicenter phase II trial. *J Clin Oncol* 2017;35:2490-2498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28475456>.

496. Kim DW, Mehra R, Tan DSW, et al. Activity and safety of ceritinib in patients with ALK-rearranged non-small-cell lung cancer (ASCEND-1): updated results from the multicentre, open-label, phase 1 trial. *Lancet Oncol* 2016;17:452-463. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26973324>.

497. Bauer TM, Shaw AT, Johnson ML, et al. Brain penetration of lorlatinib: cumulative incidences of CNS and non-CNS progression with lorlatinib in patients with previously treated ALK-positive non-small-cell lung cancer. *Target Oncol* 2020;15:55-65. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32060867>.

498. Ceresoli GL, Cappuzzo F, Gregorc V, et al. Gefitinib in patients with brain metastases from non-small-cell lung cancer: a prospective trial. *Ann Oncol* 2004;15:1042-1047. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15205197>.

499. Wu C, Li YL, Wang ZM, et al. Gefitinib as palliative therapy for lung adenocarcinoma metastatic to the brain. *Lung Cancer* 2007;57:359-364. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17434236>.

500. Arbour KC, Kris MG, Riely GJ, et al. Twice weekly pulse and daily continuous-dose erlotinib as initial treatment for patients with epidermal growth factor receptor-mutant lung cancers and brain metastases. *Cancer* 2018;124:105-109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28940498>.

501. Hoffknecht P, Tufman A, Wehler T, et al. Efficacy of the irreversible ErbB family blocker afatinib in epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI)-pretreated non-small-cell lung cancer patients with brain metastases or leptomeningeal disease. *J Thorac Oncol* 2015;10:156-163. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25247337>.

502. Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. *N Engl J Med* 2018;378:113-125. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29151359>.

503. Reungwetwattana T, Nakagawa K, Cho BC, et al. CNS response to osimertinib versus standard epidermal growth factor receptor tyrosine kinase inhibitors in patients with untreated EGFR-mutated advanced non-small-cell lung cancer. *J Clin Oncol* 2018;JCO2018783118. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30153097>.

504. Mok TS, Wu YL, Ahn MJ, et al. Osimertinib or platinum-pemetrexed in EGFR T790M-positive lung cancer. *N Engl J Med* 2017;376:629-640. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27959700>.

505. Wu YL, Ahn MJ, Garassino MC, et al. CNS efficacy of osimertinib in patients with T790M-positive advanced non-small-cell lung cancer: data from a randomized phase III trial (AURA3). *J Clin Oncol* 2018;36:2702-2709. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30059262>.

506. Yang JC, Ahn MJ, Kim DW, et al. Osimertinib in pretreated T790M-positive advanced non-small-cell lung cancer: AURA study phase II





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

extension component. J Clin Oncol 2017;35:1288-1296. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28221867>.

507. Goss G, Tsai CM, Shepherd FA, et al. Osimertinib for pretreated EGFR Thr790Met-positive advanced non-small-cell lung cancer (AURA2): a multicentre, open-label, single-arm, phase 2 study. Lancet Oncol 2016;17:1643-1652. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27751847>.

508. Goss G, Tsai CM, Shepherd FA, et al. CNS response to osimertinib in patients with T790M-positive advanced NSCLC: pooled data from two phase II trials. Ann Oncol 2018;29:687-693. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29293889>.

509. Hedde JP, Neuhaus T, Schuller H, et al. A phase I/II trial of topotecan and radiation therapy for brain metastases in patients with solid tumors. Int J Radiat Oncol Biol Phys 2007;68:839-844. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17379446>.

510. Neuhaus T, Ko Y, Muller RP, et al. A phase III trial of topotecan and whole brain radiation therapy for patients with CNS-metastases due to lung cancer. Br J Cancer 2009;100:291-297. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19127261>.

511. Petrelli F, Ghidini M, Lonati V, et al. The efficacy of lapatinib and capecitabine in HER-2 positive breast cancer with brain metastases: A systematic review and pooled analysis. Eur J Cancer 2017;84:141-148. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28810186>.

512. Freedman RA, Gelman RS, Anders CK, et al. TBCRC 022: a phase II trial of neratinib and capecitabine for patients with human epidermal growth factor receptor 2-positive breast cancer and brain metastases. J Clin Oncol 2019;37:1081-1089. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30860945>.

513. Saura C, Oliveira M, Feng YH, et al. Neratinib plus capecitabine versus lapatinib plus capecitabine in HER2-positive metastatic breast cancer previously treated with  $\geq 2$  HER2-directed regimens: phase III

NALA trial. J Clin Oncol 2020;JCO2000147. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32678716>.

514. Murthy RK, Loi S, Okines A, et al. Tucatinib, trastuzumab, and capecitabine for HER2-positive metastatic breast cancer. N Engl J Med 2020;382:597-609. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31825569>.

515. Rivera E, Meyers C, Groves M, et al. Phase I study of capecitabine in combination with temozolomide in the treatment of patients with brain metastases from breast carcinoma. Cancer 2006;107:1348-1354. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16909414>.

516. Fabi A, Vidiri A, Ferretti G, et al. Dramatic regression of multiple brain metastases from breast cancer with capecitabine: another arrow at the bow? Cancer Invest 2006;24:466-468. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16777702>.

517. Hikino H, Yamada T, Johbara K, et al. Potential role of chemo-radiation with oral capecitabine in a breast cancer patient with central nervous system relapse. Breast 2006;15:97-99. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16005228>.

518. Siegelmann-Danieli N, Stein M, Bar-Ziv J. Complete response of brain metastases originating in breast cancer to capecitabine therapy. Isr Med Assoc J 2003;5:833-834. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14650117>.

519. Wang ML, Yung WK, Royce ME, et al. Capecitabine for 5-fluorouracil-resistant brain metastases from breast cancer. Am J Clin Oncol 2001;24:421-424. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11474279>.

520. Awada A, Colomer R, Inoue K, et al. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. JAMA Oncol 2016;2:1557-1564. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27078022>.





# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

521. Lassman AB, Abrey LE, Shah GD, et al. Systemic high-dose intravenous methotrexate for central nervous system metastases. *J Neurooncol* 2006;78:255-260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16344918>.

522. Vinolas N, Graus F, Mellado B, et al. Phase II trial of cisplatin and etoposide in brain metastases of solid tumors. *J Neurooncol* 1997;35:145-148. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9266451>.

523. Franciosi V, Cocconi G, Michiara M, et al. Front-line chemotherapy with cisplatin and etoposide for patients with brain metastases from breast carcinoma, nonsmall cell lung carcinoma, or malignant melanoma: a prospective study. *Cancer* 1999;85:1599-1605. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10193952>.

524. Cocconi G, Lottici R, Bisagni G, et al. Combination therapy with platinum and etoposide of brain metastases from breast carcinoma. *Cancer Invest* 1990;8:327-334. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2207757>.

525. Pavlidis N. The diagnostic and therapeutic management of leptomeningeal carcinomatosis. *Ann Oncol* 2004;15 Suppl 4:iv285-291. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15477323>.

526. Waki F, Ando M, Takashima A, et al. Prognostic factors and clinical outcomes in patients with leptomeningeal metastasis from solid tumors. *J Neurooncol* 2009;93:205-212. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19043775>.

527. Clarke JL, Perez HR, Jacks LM, et al. Leptomeningeal metastases in the MRI era. *Neurology* 2010;74:1449-1454. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20439847>.

528. Chamberlain MC. Leptomeningeal metastasis. *Curr Opin Neurol* 2009;22:665-674. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19738466>.

529. Liaw CC, Ng KT, Huang JS, et al. Meningeal carcinomatosis from solid tumors: clinical analysis of 42 cases. *J Formos Med Assoc*

1992;91:299-303. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1354692>.

530. Wasserstrom WR, Glass JP, Posner JB. Diagnosis and treatment of leptomeningeal metastases from solid tumors: experience with 90 patients. *Cancer* 1982;49:759-772. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6895713>.

531. Wolf A, Donahue B, Silverman JS, et al. Stereotactic radiosurgery for focal leptomeningeal disease in patients with brain metastases. *J Neurooncol* 2017;134:139-143. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28536993>.

532. Sandberg DI, Bilsky MH, Souweidane MM, et al. Ommaya reservoirs for the treatment of leptomeningeal metastases. *Neurosurgery* 2000;47:49-54; discussion 54-45. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10917346>.

533. Fleischhack G, Jaehde U, Bode U. Pharmacokinetics following intraventricular administration of chemotherapy in patients with neoplastic meningitis. *Clin Pharmacokinet* 2005;44:1-31. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15634030>.

534. Boogerd W, van den Bent MJ, Koehler PJ, et al. The relevance of intraventricular chemotherapy for leptomeningeal metastasis in breast cancer: a randomised study. *Eur J Cancer* 2004;40:2726-2733. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15571954>.

535. Glantz MJ, Jaeckle KA, Chamberlain MC, et al. A randomized controlled trial comparing intrathecal sustained-release cytarabine (DepoCyt) to intrathecal methotrexate in patients with neoplastic meningitis from solid tumors. *Clin Cancer Res* 1999;5:3394-3402. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10589750>.

536. Grossman SA, Finkelstein DM, Ruckdeschel JC, et al. Randomized prospective comparison of intraventricular methotrexate and thiotepa in patients with previously untreated neoplastic meningitis. Eastern Cooperative Oncology Group. *J Clin Oncol* 1993;11:561-569. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8445432>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

537. Glantz MJ, LaFollette S, Jaeckle KA, et al. Randomized trial of a slow-release versus a standard formulation of cytarabine for the intrathecal treatment of lymphomatous meningitis. *J Clin Oncol* 1999;17:3110-3116. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10506606>.

538. Jaeckle KA, Phuphanich S, Bent MJ, et al. Intrathecal treatment of neoplastic meningitis due to breast cancer with a slow-release formulation of cytarabine. *Br J Cancer* 2001;84:157-163. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11161370>.

539. Gutin PH, Weiss HD, Wiernik PH, Walker MD. Intrathecal N, N', N"-triethylenethiophosphoramidate [thio-TEPA (NSC 6396)] in the treatment of malignant meningeal disease: phase I-II study. *Cancer* 1976;38:1471-1475. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/825215>.

540. Chamberlain MC, Johnston SK, Van Horn A, Glantz MJ. Recurrent lymphomatous meningitis treated with intra-CSF rituximab and liposomal ara-C. *J Neurooncol* 2009;91:271-277. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18820836>.

541. Groves MD, Glantz MJ, Chamberlain MC, et al. A multicenter phase II trial of intrathecal topotecan in patients with meningeal malignancies. *Neuro Oncol* 2008;10:208-215. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18316473>.

542. Chamberlain MC, Tsao-Wei DD, Groshen S. Phase II trial of intracerebrospinal fluid etoposide in the treatment of neoplastic meningitis. *Cancer* 2006;106:2021-2027. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16583432>.

543. Zagouri F, Sergentanis TN, Bartsch R, et al. Intrathecal administration of trastuzumab for the treatment of meningeal carcinomatosis in HER2-positive metastatic breast cancer: a systematic review and pooled analysis. *Breast Cancer Res Treat* 2013;139:13-22. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23588955>.

544. Mrugala MM, Kim B, Sharma A, et al. Phase II study of systemic high-dose methotrexate and intrathecal liposomal cytarabine for treatment

of leptomeningeal carcinomatosis from breast cancer. *Clin Breast Cancer* 2019;19:311-316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31175053>.

545. Bokstein F, Lossos A, Lossos IS, Siegal T. Central nervous system relapse of systemic non-Hodgkin's lymphoma: results of treatment based on high-dose methotrexate combination chemotherapy. *Leuk Lymphoma* 2002;43:587-593. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12002763>.

546. Grommes C, Oxnard GR, Kris MG, et al. "Pulsatile" high-dose weekly erlotinib for CNS metastases from EGFR mutant non-small cell lung cancer. *Neuro Oncol* 2011;13:1364-1369. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21865399>.

547. Yang JCH, Kim SW, Kim DW, et al. Osimertinib in patients with epidermal growth factor receptor mutation-positive non-small-cell lung cancer and leptomeningeal metastases: the BLOOM study. *J Clin Oncol* 2020;38:538-547. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31809241>.

548. Nanjo S, Hata A, Okuda C, et al. Standard-dose osimertinib for refractory leptomeningeal metastases in T790M-positive EGFR-mutant non-small cell lung cancer. *Br J Cancer* 2018;118:32-37. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29190637>.

549. Ahn MJ, Chiu CH, Cheng Y, et al. Osimertinib for patients with leptomeningeal metastases associated with EGFR T790M-positive advanced NSCLC: the AURA leptomeningeal metastases analysis. *J Thorac Oncol* 2020;15:637-648. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31887431>.

550. Lee JS, Melisko ME, Magbanua MJ, et al. Detection of cerebrospinal fluid tumor cells and its clinical relevance in leptomeningeal metastasis of breast cancer. *Breast Cancer Res Treat* 2015;154:339-349. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26520840>.

551. Lin X, Fleisher M, Rosenblum M, et al. Cerebrospinal fluid circulating tumor cells: a novel tool to diagnose leptomeningeal metastases from



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

epithelial tumors. *Neuro Oncol* 2017;19:1248-1254. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28821205>.

552. Nayak L, Fleisher M, Gonzalez-Espinoza R, et al. Rare cell capture technology for the diagnosis of leptomeningeal metastasis in solid tumors. *Neurology* 2013;80:1598-1605; discussion 1603. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23553479>.

553. Ibrahim A, Crockard A, Antonietti P, et al. Does spinal surgery improve the quality of life for those with extradural (spinal) osseous metastases? An international multicenter prospective observational study of 223 patients. Invited submission from the Joint Section Meeting on Disorders of the Spine and Peripheral Nerves, March 2007. *J Neurosurg Spine* 2008;8:271-278. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18312079>.

554. Byrne TN. Spinal cord compression from epidural metastases. *N Engl J Med* 1992;327:614-619. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1296600>.

555. Schmidt MH, Klimo P, Jr., Vrionis FD. Metastatic spinal cord compression. *J Natl Compr Canc Netw* 2005;3:711-719. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16194459>.

556. Helweg-Larsen S, Sorensen PS. Symptoms and signs in metastatic spinal cord compression: a study of progression from first symptom until diagnosis in 153 patients. *Eur J Cancer* 1994;30A:396-398. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8204366>.

557. Levack P, Graham J, Collie D, et al. Don't wait for a sensory level--listen to the symptoms: a prospective audit of the delays in diagnosis of malignant cord compression. *Clin Oncol (R Coll Radiol)* 2002;14:472-480. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12512970>.

558. Sundaresan N, Rothman A, Manhart K, Kelliher K. Surgery for solitary metastases of the spine: rationale and results of treatment. *Spine (Phila Pa 1976)* 2002;27:1802-1806. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12195075>.

559. North RB, LaRocca VR, Schwartz J, et al. Surgical management of spinal metastases: analysis of prognostic factors during a 10-year experience. *J Neurosurg Spine* 2005;2:564-573. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15945430>.

560. Tomita K, Kawahara N, Kobayashi T, et al. Surgical strategy for spinal metastases. *Spine (Phila Pa 1976)* 2001;26:298-306. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11224867>.

561. Tokuhashi Y, Matsuzaki H, Oda H, et al. A revised scoring system for preoperative evaluation of metastatic spine tumor prognosis. *Spine (Phila Pa 1976)* 2005;30:2186-2191. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16205345>.

562. van der Linden YM, Dijkstra SP, Vonk EJ, et al. Prediction of survival in patients with metastases in the spinal column: results based on a randomized trial of radiotherapy. *Cancer* 2005;103:320-328. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15593360>.

563. Bartels RH, Feuth T, van der Maazen R, et al. Development of a model with which to predict the life expectancy of patients with spinal epidural metastasis. *Cancer* 2007;110:2042-2049. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17853394>.

564. Choi D, Crockard A, Bunger C, et al. Review of metastatic spine tumour classification and indications for surgery: the consensus statement of the Global Spine Tumour Study Group. *Eur Spine J* 2010;19:215-222. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20039084>.

565. Eleraky M, Papanastassiou I, Vrionis FD. Management of metastatic spine disease. *Curr Opin Support Palliat Care* 2010;4:182-188. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20671554>.

566. Klimo P, Jr., Thompson CJ, Kestle JR, Schmidt MH. A meta-analysis of surgery versus conventional radiotherapy for the treatment of metastatic spinal epidural disease. *Neuro Oncol* 2005;7:64-76. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15701283>.



# NCCN Guidelines Version 3.2020

## Central Nervous System Cancers

567. Witham TF, Khavkin YA, Gallia GL, et al. Surgery insight: current management of epidural spinal cord compression from metastatic spine disease. *Nat Clin Pract Neurol* 2006;2:87-94; quiz 116. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16932530>.

568. Patchell RA, Tibbs PA, Regine WF, et al. Direct decompressive surgical resection in the treatment of spinal cord compression caused by metastatic cancer: a randomised trial. *Lancet* 2005;366:643-648. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16112300>.

569. Patil CG, Lad SP, Santarelli J, Boakye M. National inpatient complications and outcomes after surgery for spinal metastasis from 1993-2002. *Cancer* 2007;110:625-630. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17577227>.

570. Itshayek E, Yamada J, Bilsky M, et al. Timing of surgery and radiotherapy in the management of metastatic spine disease: a systematic review. *Int J Oncol* 2010;36:533-544. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20126972>.

571. Maranzano E, Bellavita R, Rossi R, et al. Short-course versus split-course radiotherapy in metastatic spinal cord compression: results of a phase III, randomized, multicenter trial. *J Clin Oncol* 2005;23:3358-3365. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15738534>.

572. Gerszten PC, Mendel E, Yamada Y. Radiotherapy and radiosurgery for metastatic spine disease: what are the options, indications, and outcomes? *Spine (Phila Pa 1976)* 2009;34:S78-92. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19829280>.

573. Redmond KJ, Robertson S, Lo SS, et al. Consensus contouring guidelines for postoperative stereotactic body radiation therapy for metastatic solid tumor malignancies to the spine. *Int J Radiat Oncol Biol Phys* 2017;97:64-74. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27843035>.

574. Glicksman RM, Tjong MC, Neves-Junior WFP, et al. Stereotactic ablative radiotherapy for the management of spinal metastases: a review.

*JAMA Oncol* 2020;6:567-577. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31895403>.

575. Sprave T, Verma V, Forster R, et al. Local response and pathologic fractures following stereotactic body radiotherapy versus three-dimensional conformal radiotherapy for spinal metastases - a randomized controlled trial. *BMC Cancer* 2018;18:859. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30170568>.

576. Redmond KJ, Sciubba D, Khan M, et al. A phase 2 study of post-operative stereotactic body radiation therapy (SBRT) for solid tumor spine metastases. *Int J Radiat Oncol Biol Phys* 2020;106:261-268. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31628959>.

577. Ryu S, Pugh SL, Gerszten PC, et al. RTOG 0631 phase 2/3 study of image guided stereotactic radiosurgery for localized (1-3) spine metastases: phase 2 results. *Pract Radiat Oncol* 2014;4:76-81. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24890347>.

578. Cox BW, Spratt DE, Lovelock M, et al. International Spine Radiosurgery Consortium consensus guidelines for target volume definition in spinal stereotactic radiosurgery. *Int J Radiat Oncol Biol Phys* 2012;83:e597-605. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22608954>.

579. Redmond KJ, Lo SS, Soltys SG, et al. Consensus guidelines for postoperative stereotactic body radiation therapy for spinal metastases: results of an international survey. *J Neurosurg Spine* 2017;26:299-306. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27834628>.

580. Mendel E, Bourekas E, Gerszten P, Golan JD. Percutaneous techniques in the treatment of spine tumors: what are the diagnostic and therapeutic indications and outcomes? *Spine (Phila Pa 1976)* 2009;34:S93-100. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19829281>.

581. Sorensen S, Helweg-Larsen S, Mouridsen H, Hansen HH. Effect of high-dose dexamethasone in carcinomatous metastatic spinal cord compression treated with radiotherapy: a randomised trial. *Eur J Cancer*





National  
Comprehensive  
Cancer  
Network®

## NCCN Guidelines Version 3.2020 Central Nervous System Cancers

---

1994;30A:22-27. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8142159>.

582. Vecht CJ, Haaxma-Reiche H, van Putten WL, et al. Initial bolus of conventional versus high-dose dexamethasone in metastatic spinal cord compression. *Neurology* 1989;39:1255-1257. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/2771077>.

583. Heimdal K, Hirschberg H, Slettebo H, et al. High incidence of serious side effects of high-dose dexamethasone treatment in patients with epidural spinal cord compression. *J Neurooncol* 1992;12:141-144.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1560260>.

584. Kwok Y, Regine WF, Patchell RA. Radiation therapy alone for spinal cord compression: time to improve upon a relatively ineffective status quo. *J Clin Oncol* 2005;23:3308-3310. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15738546>.

585. Rades D, Stalpers LJA, Veninga T, et al. Evaluation of five radiation schedules and prognostic factors for metastatic spinal cord compression. *J Clin Oncol* 2005;23:3366-3375. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15908648>.