

# Randomized Trial of Carboplatin Versus Radiotherapy for Stage I Seminoma: Mature Results on Relapse and Contralateral Testis Cancer Rates in MRC TE19/EORTC 30982 Study (ISRCTN27163214)

R. Timothy D. Oliver, Graham M. Mead, Gordon J.S. Rustin, Johnathan K. Joffe, Nina Aass, Robert Coleman, Rhian Gabe, Philip Pollock, and Sally P. Stenning

See accompanying editorial on page 949

From St Bartholomews and the London Hospital; Medical Research Council Clinical Trials Unit, Cancer Group, London; Department of Medical Oncology, Southampton General Hospital, Southampton; Medical Oncology, Mount Vernon Cancer Centre, Northwood; Department of Medical Oncology, Huddersfield Royal Infirmary, Huddersfield; and Weston Park Hospital, Sheffield, United Kingdom; and Norwegian Radium Hospital, Oslo, Norway.

Submitted October 2, 2009; accepted February 23, 2010; published online ahead of print at www.jco.org on January 31, 2011.

Written on behalf of the MRC TE19/EORTC 30982 Study Group.

Supported by United Kingdom Medical Research Council.

Presented at the 44th Annual Meeting of the American Society for Clinical Oncology, May 30-June 3, 2008, Chicago, IL, and at the Clinical Trials Showcase, National Cancer Research Institute Annual Conference, October 5-8, 2008, Birmingham United Kingdom.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Clinical Trials repository link available on JCO.org.

Corresponding author: Sally P. Stenning, MSc, MRC Clinical Trials Unit, 222 Euston Rd, London, NW1 2DA, United Kingdom; e-mail: ss@ctu.mrc.ac.uk.

© 2011 by American Society of Clinical Oncology

0732-183X/11/2908-957/\$20.00

DOI: 10.1200/JCO.2009.26.4655

## ABSTRACT

### Purpose

Initial results of a randomized trial comparing carboplatin with radiotherapy (RT) as adjuvant treatment for stage I seminoma found carboplatin had a noninferior relapse-free rate (RFR) and had reduced contralateral germ cell tumors (GCTs) in the short-term. Updated results with a median follow-up of 6.5 years are now reported.

### Patients and Methods

Random assignment was between RT and one infusion of carboplatin dosed at  $7 \times$  (glomerular filtration rate + 25) on the basis of EDTA ( $n = 357$ ) and 90% of this dose if determined on the basis of creatinine clearance ( $n = 202$ ). The trial was powered to exclude a doubling in RFRs assuming a 96-97% 2-year RFR after radiotherapy (hazard ratio [HR], approximately 2.0).

### Results

Overall, 1,447 patients were randomly assigned in a 3-to-5 ratio (carboplatin,  $n = 573$ ; RT,  $n = 904$ ). RFRs at 5 years were 94.7% for carboplatin and 96.0% for RT (RT-C 90% CI, 0.7% to 3.5%; HR, 1.25; 90% CI, 0.83 to 1.89). One death as a result of seminoma (in RT arm) occurred. Patients receiving at least 99% of the  $7 \times$  AUC dose had a 5-year RFR of 96.1% (95% CI, 93.4% to 97.7%) compared with 92.6% (95% CI, 88.0% to 95.5%) in those who received lower doses (HR, 0.51; 95% CI, 0.24 to 1.07;  $P = .08$ ). There was a clear reduction in the rate of contralateral GCTs (carboplatin,  $n = 2$ ; RT,  $n = 15$ ; HR, 0.22; 95% CI, 0.05 to 0.95;  $P = .03$ ), and elevated pretreatment follicle-stimulating hormone (FSH) levels ( $> 12$  IU/L) was a strong predictor (HR, 8.57; 95% CI, 1.82 to 40.38).

### Conclusion

These updated results confirm the noninferiority of single dose carboplatin (at  $7 \times$  AUC dose) versus RT in terms of RFR and establish a statistically significant reduction in the medium term of risk of second GCT produced by this treatment.

*J Clin Oncol* 29:957-962. © 2011 by American Society of Clinical Oncology

## INTRODUCTION

Stage I seminoma is the commonest presentation of testicular germ cell tumor (GCT) and accounts for approximately 40% of all occurrences.<sup>1</sup> Surveillance studies suggest that 15% to 20% of these patients harbor metastatic disease.<sup>2</sup> Three management approaches have been investigated: surveillance, adjuvant radiotherapy,<sup>3-5</sup> and carboplatin.<sup>6,7</sup> The latter two treatments have been the subject of large, Medical Research Council (MRC) – organized, randomized trials.<sup>8-10</sup> The third and largest of these trials, TE19 (in collaboration with European Organisa-

tion for Research and Treatment of Cancer study EORTC 30982), was conducted between 1996 and 2001; 1,477 patients were randomly assigned to either radiotherapy (RT) or a single injection of adjuvant carboplatin at seven times the area under the curve (AUC) dose (AUC 7). Preliminary results of this study, which were reported in 2005,<sup>10</sup> suggested that carboplatin was noninferior to RT in preventing metastatic relapse and, in addition, may reduce the risk of subsequent contralateral testicular cancer. To confirm these initial findings with extended follow-up, we now present more mature data, together with new analyses investigating a possible carboplatin

dose response effect on relapse rates and risk factors for contralateral cancer.

## PATIENTS AND METHODS

Consenting patients with stage I, histologically confirmed, seminomatous GCT (classical or anaplastic) and normal markers were eligible, as previously described. The primary random assignment of the study was between irradiation (optional subrandom assignment with respect to dose) and one infusion of carboplatin with dose adjusted by renal function to provide seven times the AUC estimate of glomerular filtration rate (GFR). Random assignment was effected by minimization, and stratification was by center, previous inguino/pelvic/scrotal surgery, and intended RT schedule (ie, randomly assigned or elected).

### Treatment Schedules

**Carboplatin.** Using  $7 \times (\text{GFR} + 25)$ , the carboplatin dose in milligrams was given as a single, intravenous (IV) dose.<sup>11</sup> EDTA or a comparable isotope-measurement technique was the recommended method of measuring GFR. When this method was unavailable, a urinary 24-hour creatinine clearance (but not a calculated GFR by using, for example, the Cockcroft formula) was also permitted. In the latter case, the protocol recommended that the dose given should be 10% lower than if the dose had been calculated on the basis of EDTA clearance. This has allowed exploratory analysis of the impact of carboplatin dose on outcome.

**RT.** The RT dose was determined either by an optional random assignment between 30 Gy in 15 fractions and 20 Gy in 10 fractions as part of the TE18 trial<sup>9</sup> or, for those patients who did not enter this additional random assignment, by physician choice. In this latter group, schedules delivering a total dose of 20 Gy, 30 Gy, or an intermediate dose were permissible after discussion with the principal investigators. The recommended RT field was a para-aortic (PA) strip; however a dog leg (DL) field was recommended for patients with previous inguinal surgery, as in our previous trials.

### Gonadal Function Assays

Serial gonadal function was assessed by measurement of follicle-stimulating hormone (FSH), luteinizing hormone (LH), and testosterone levels. Levels were obtained at pretreatment and at 12 and 24 months after treatment.

### Follow-Up Investigations and Management of Relapse


Clinical and tumor marker assessments took place every 3 months in year 1; every 4 months in year 2; every 6 months in year 3; annually thereafter until year 10. Chest x-rays were done at 6, 12, 20, 30, and 36 months, and computed tomography (CT) scans of chest, abdomen, and pelvis were done at 12, 24, and 36 months. On relapse, standard chemotherapy regimens for metastatic seminoma were recommended, but those patients who experienced relapses localized outside previously irradiated areas could be managed with RT at the treating clinician's discretion.

The primary outcome measure was relapse-free rate; death as a result of any cause and second cancer incidence were also recorded, as was morbidity, which was assessed via patient diary cards. The latter was reported in the initial publication and is not repeated here.

### Statistical Considerations

The aim of the trial was to demonstrate the noninferiority of carboplatin in comparison with irradiation, irrespective of dose. It is now known that 20 Gy is equivalent in efficacy to 30 Gy.<sup>9</sup> As this had not been demonstrated at the time the trial was initiated, the trial included an optional random assignment between these doses. To optimize the overall power of the trial for both primary (RT v carboplatin) and secondary comparisons (30 Gy v 20 Gy v carboplatin) patients were randomly assigned between RT and carboplatin in a five-to-three ratio; hence, 30 Gy RT, 20 Gy RT, and carboplatin, when applicable, were randomly assigned in a ratio of 2.5 to 2.5 to 3, respectively.

Assuming a 2-year relapse-free rate after RT of 96% to 97%, the trial was powered to exclude a doubling of the relapse rate at 2 years with 90% power at a one-sided significance level of 5%. This corresponds to a hazard ratio (HR) of



**Fig 1.** CONSORT diagram. SD, standard deviation; HCG, human chorionic gonadotrophin; IQR, interquartile range; RT, radiotherapy; ITT, intent-to-treat; PPA, per-protocol analysis.

approximately 2.0; thus, for this updated analysis to demonstrate noninferiority with respect to relapse rates in the longer term, the upper confidence limit of a two-sided 90% CI for the HR must be less than 2.0.

### Analysis


As the clinical equivalence of 30 Gy and 20 Gy has now been demonstrated,<sup>9</sup> this analysis has focused on the primary comparison of RT versus carboplatin. RFRs were calculated by using the Kaplan-Meier method and were compared by using the log-rank test; HRs and 90% CIs were computed by using Cox proportional hazard regression model (HR > 1 favors RT). CIs for the difference in relapse-free rates at 2 and 5 years were calculated by direct comparison of proportions and by application of the HR and its 90% CI to the relapse-free rate in the control arm. Both intent-to-treat (ITT) analyses and per-protocol analyses (PPA) were carried out, as poor compliance can bias toward equivalence in an ITT analysis. PPA included all patients known to have started their allocated treatment (CONSORT diagram, Fig 1).

## RESULTS

Overall, 1,477 patients were recruited to the trial between 1996 and 2001; 573 were randomly assigned to carboplatin, and 904 were randomly assigned to RT. Eighty-seven percent of the patients had PA-field radiotherapy. Ten patients did not receive their allocated treatment because of revised stage (carboplatin, *n* = 2; RT, *n* = 4) or histology (RT, *n* = 4) before starting treatment; an additional 17 patients (carboplatin, *n* = 9; RT, *n* = 8) withdrew and requested alternative (or no) treatment; and treatment data are missing for one patient in each group. These patients were excluded from the PPA.

Follow-up data for this analysis were obtained directly from the participating sites during 2007. All events up to the date of last clinic visit were included to ensure reliable data on recurrence. Protocol-specified follow-up was for a maximum of 10 years. The median follow-up is now approximately 6.5 years in both treatment groups (interquartile range, 5.2 to 8 years). The number of patients who have a minimum of 5 years documented follow-up has increased since our last report from 333 to 1,148 (79%), whereas 615 patients (40%) have a minimum follow-up of 7 years (which has increased from 18 patients).

The updated results confirm the noninferiority of single-dose carboplatin (Table 1; Fig 2) with only one additional relapse reported (in the R arm at 61 months). The RFRs were 97.3% (C) versus 96.5% (RT) at 2 years and 94.7% for the carboplatin arm versus 96.0% for the RT arm respectively at 5 years (ITT analysis: HR, 1.25; 90% CI, 0.83 to 1.89; *P* = .37; PPA: HR, 1.27; 90% CI, 0.83 to 1.92; *P* = .36). The 90% CI for the absolute difference excludes rates above 3% at 2 years and



**Fig 2.** Relapse-free rates (RFRs) by allocated treatment. Intent-to-treat (ITT) analysis (hazard ratio [HR] > 1 favors radiotherapy [RT]): HR, 1.25; 90% CI, 0.83 to 1.89; *P* = .37. Per-protocol analysis (PPA): HR, 1.27; 90% CI, 0.83 to 1.92; *P* = .36. RFR at 5 years: radiotherapy, 96.0% (95% CI, 94.5% to 97.1%); carboplatin (C), 94.7% (95% CI, 92.5% to 96.3%). RFR absolute difference at 5 years: RT-to-C difference, 1.34%; 90% CI, -0.7% to 3.5%.

above 3.5% at 5 years by both methods. Ten patients have died in the RT arm: one as a result of seminoma, three as a result of other (ie, non-GCT) cancers, and six as a result of other noncancer causes. Six patients have died in the carboplatin arm: none as a result of seminoma; two as a result of other (ie, non-GCT) cancers; and four as a result of other noncancer causes.

Contralateral GCTs have been reported in two patients in the carboplatin arm and in 15 in the RT arm; the contralateral GCT-free rates at 5 years are 99.8% and 98.8%, respectively (Fig 3A), which results in a relative reduction in risk of nearly 80% (HR, 0.22; 95% CI, 0.05 to 0.95; *P* = .03). A strong predictor of contralateral GCT was pretreatment FSH. This was recorded in 1,008 patients, of whom 31% had raised levels (ie, > 12 IU/L). These patients showed an 8.6-fold increase in the rate of contralateral cancer compared with those who had a normal (ie, ≤ 12 IU/L) FSH (HR, 8.57; 95% CI, 1.82 to 40.4; *P* = .001; Fig 3B).


### Exploratory Analyses

Of 559 patients in the carboplatin arm for whom the method of GFR assessment was known, 347 used EDTA or comparable isotopic techniques and had a mean GFR of 128 mL/min, whereas 212 used urinary creatinine clearance and had a mean GFR of 125 mL/min. The patients in the former group were planned to receive a dose of 7 × (GFR + 25); those in the latter group were planned to receive 90% of this dose. This gave an opportunity for an exploratory analysis of RFRs according to the proportion of the AUC 7 dose received (Fig 4). The 5-year RFRs were 92.6% (95% CI, 88.0% to 95.5%) and 96.1% (95% CI, 93.4 to 97.7%), respectively, (HR, 0.51; 95% CI, 0.24 to 1.07; *P* = .08) whether the dose received was less than AUC 7 or at least AUC 7; after adjustment for GFR method, the results were similar (HR, 0.48; 95% CI, 0.16 to 1.42; *P* = .19). Serial FSH measurements (at pretreatment and also at 12 ± 2 months and 24 ± 2 months) were available for 170 patients receiving carboplatin and for 243 patients receiving RT, of whom the majority (*n* =

**Table 1.** Summary of Events

Event	Treatment Arm			
	Carboplatin ( <i>n</i> = 573)		Radiotherapy ( <i>n</i> = 904)	
	No.	%	No.	%
Total relapse	29	5.1	37	4.1
New primary cancers	7	1.2	25	2.8
GCT	2		15	
Other	5		10	
Total deaths	6	1.0	10	1.1
Death as a result of seminoma	0		1	
Death as a result of other cause	6		9	

Abbreviation: GCT, germ cell tumor.




**Fig 3.** (A) Contralateral germ cell tumor (GCT) rates by allocated treatment (truncated y-axis). Second GCT-free rate at 5 years: radiotherapy (RT), 98.8% (95% CI, 97.8% to 99.4%); carboplatin (C), 99.8% (95% CI, 98.6% to 99.9%). Hazard ratio (HR), 0.22; 95% CI, 0.05 to 0.95;  $P = .03$ . (B) Contralateral GCT rate by pretreatment follicle-stimulating hormone (FSH; truncated y-axis):  $\leq 12$  IU/L  $v > 12$  IU/L. Pretreatment FSH: normal ( $\leq 12$  IU/L), 99.8% (95% CI, 98.9% to 99.9%); increased ( $> 12$  IU/L), 98.2% (95% CI, 96.0% to 99.3%); HR, 8.57; 95% CI, 1.82 to 40.4; log-rank  $P = .001$ .

185) received RT to the PA strip without scrotal shielding. Median FSH levels at these time points were 9.4, 10.0, and 9.6 IU/L for patients receiving carboplatin and were 9.9, 11.0, and 9.9 IU/L for the patients receiving PA RT. Within-patient changes from baseline were compared across treatment groups and showed no clear evidence of a differential treatment impact ( $t$  test  $P = .15$  and  $P = .17$  at 12 and 24 months, respectively).

## DISCUSSION

This article provides mature data for patients and their physicians on the relative long-term efficacy of a single infusion of carboplatin as adju-



**Fig 4.** Relapse-free rate (RFRs) by received carboplatin dose less than seven times area under the curve (AUC) compared with seven or more times the AUC. RFR at 5 years: carboplatin dose less than seven times AUC, 92.6% (95% CI, 88.0% to 95.5%); seven or more times AUC, 96.1% (95% CI, 93.4% to 97.7%); hazard ratio (HR), 0.51; 95% CI, 0.24 to 1.07;  $P = .08$  (unadjusted for glomerular filtration rate [GFR] method); HR, 0.48; 95% CI, 0.16 to 1.42;  $P = .19$  (adjusted for GFR method). HR less than 1 favors higher dose.

vant treatment for stage I seminoma. The initial studies of one and two courses of adjuvant carboplatin that motivated this work, including the first to use a single course at AUC 7,<sup>6</sup> are also now mature and provide additional evidence of the long-term safety and efficacy of this approach.

The management of this condition should be a matter of informed choice. The anticipated relapse rate in patients managed expectantly is in the range of 15% to 20%,<sup>2,12</sup> and primary cure rates for the three currently available management options (ie, surveillance, adjuvant RT, and carboplatin) approach 100% in all modern series. As a result of this large, randomized trial of RT and carboplatin, of previous radiation-field and dose trials,<sup>8,9</sup> and of retrospective case series of surveillance,<sup>2</sup> there are now adequate data on efficacy, short-term morbidity, contralateral cancer, and necessary follow-up arrangements to inform this decision.

Adjuvant carboplatin as described in our previous paper is associated with minimal short-term toxicity<sup>10</sup> and has no significant effect on fertility, as judged by serial FSH levels.<sup>10,13</sup> Relapse rates are reduced by between two thirds and three quarters compared with surveillance, and the cause-specific survival is currently 100%. Late relapse is a rare event with, to date, only one occurring beyond 3 years; in an accompanying paper that investigates patterns of relapse,<sup>14</sup> we provide data to suggest that post-treatment follow-up can be limited to 3 years.

An unexpected advantage of carboplatin reported at our initial analysis was a marked reduction in the risk of contralateral cancer. This has been maintained with extended follow-up, as all additional contralateral GCTs reported subsequently were in patients receiving RT. Elevated pretreatment FSH has been proposed as a risk factor for contralateral testis cancer in other studies,<sup>15</sup> and we were able to confirm this finding prospectively. Carcinoma in situ (CIS) of the contralateral testis is associated with testicular atrophy and impaired spermatogenesis, which results in increased FSH levels.<sup>16</sup> The reports of increased sperm counts and decreasing FSH levels after treatment with carboplatin,<sup>13</sup> and of clearance (temporary, in some cases) of CIS in the contralateral testis after bleomycin, etoposide, and cisplatin chemotherapy,<sup>17</sup> support the hypothesis that carboplatin may have a similar impact on CIS. The potential for carboplatin to have delayed,



**Table 2.** 5-Year RFR in Carboplatin Monotherapy Studies Reported in the Literature

Clinical setting	Planned Carboplatin Dose							
	AUC × 10		AUC × ≥ 7*		AUC × < 7†		400 mg/m <sup>2</sup> ‡	
	No.	5-Year RFR (%)	No.	5-Year RFR (%)	No.	5-Year RFR (%)	No.	5-Year RFR (%)
Stage I seminoma	ND		347§	96	212§	93	93	91
Metastatic seminoma	24¶	92	17¶	88	ND		19¶	79
							177#	72
							108**	82

Abbreviations: AUC, area under the curve; RFR, relapse-free rate; ND, not determined.

\*Approximate dose, 1,071 mg.

†Approximately 963 mg.

‡Approximately 800 mg.

§TE19/EORTC 30982 (current data).

||Dieckman et al.<sup>21</sup>

¶Oliver et al.<sup>22</sup>

#Bokemeyer et al.<sup>23</sup>

\*\*Krege et al.<sup>24</sup> (stages IIA and IIB only).

rather than prevented, contralateral GCTs remains, as long-term follow-up of the original phase II study<sup>6</sup> has identified three late contralateral tumors in a series of 73 men observed for 10 to 25 years. Continued follow-up of these series is needed, therefore; however, these data do provide encouragement for patients who have not completed their family and present with elevated FSH and/or an atrophic contralateral testis to consider being included in trials of short-term testis preservation by using partial orchiectomy plus carboplatin.<sup>18-20</sup>

The mean GFR obtained in the two groups of patients assessed by EDTA and creatinine clearance were similar. The previous recommendation for dose adjustment in the latter group was clearly inappropriate; as a consequence, we have data that suggest a carboplatin dose-response effect. As a nonrandomized comparison, partly confounded by center and the GFR method, this must be treated with caution. However, data from a number of studies using carboplatin at varying dose levels in both stage I and metastatic disease (Table 2) support the concept of a steep dose response, though standardization of dosing on the basis of GFR obtained with an isotope technique and more studies in patients with metastatic seminoma are required to be certain.

Retrospective analysis of data from a number of centers has suggested that tumor size greater than 4 cm and infiltration of the rete testis were the main risk factors for relapse on surveillance.<sup>2</sup> Two large, prospective studies with carboplatin that used this and other prognostic factor data have been reported by a Spanish group. Patients were divided into prognostic groups by using histologic risk factors, and carboplatin (two cycles of AUC 7 given 21 days apart) was reserved for high-risk groups. In an initial study,<sup>25</sup> vascular invasion and/or pT stage were used to divide the patients into a low-risk group (41% of patients), who underwent surveillance, whereas the remaining patients receiving carboplatin. Relapse rates were, respectively, 16% and 3%, and 5-year cause-specific survival was 100%. In a second study,<sup>26</sup> this group modified the prognostic criteria by using tumor size (> 4 cm) and/or rete involvement to identify patients for treatment. Sixty-eight percent of patients received carboplatin, and relapse rates were 7% on surveillance and 4% on carboplatin.

The data reported by this group, together with extensive phase II data,<sup>6,27</sup> support the efficacy of single-agent carboplatin used for one or two treatment courses, as described. Concern has been expressed about

potential long-term toxicity; however, in a recent publication documenting long-term follow-up (median, 9 years) in 199 patients, there was no increase in overall mortality nor in death as a result of circulatory disease or the incidence of secondary, non-GCT malignancies.<sup>6</sup>

RT has been the adjuvant management approach of choice for at least 50 years. The Medical Research Council trials have shown that this treatment can be limited to 20 Gy to a PA field for most patients, reducing the risks of infertility. Radiation, however, is associated with more short-term toxicity than carboplatin<sup>10</sup> and is associated with an increased risk of delayed cardiovascular disease and second malignancy.<sup>4</sup> As a result, this treatment is in marked decline in Britain<sup>28</sup> and the United States.<sup>29</sup> The principal alternative to carboplatin is currently surveillance. This approach has been recommended by European Germ Cell Cancer Consensus Group guidelines<sup>30</sup> for low-risk patients for whom compliance is not a concern. However, there are currently no evidence-based protocols to guide follow-up and the frequency of radiologic investigation. Surveillance has the advantage of restricting treatment (usually with RT) to the 15% to 20% of patients who experience relapse. However, because of an incidence of late relapse, follow-up needs to be prolonged, which causes potential associated anxiety and noncompliance. Most clinicians recommend multiple follow-up CT scans that are associated with undesirable radiation exposure.<sup>31</sup> A current Medical Research Council randomized trial (Trial of Imaging and Surveillance in Seminoma Testis) is evaluating the optimal frequency of scans and the utility of magnetic resonance imaging in these patients, thereby avoiding potentially needless irradiation.<sup>32</sup>

To date, when a risk-based approach to management has been taken, histopathologic risk factors have been used to target surveillance or adjuvant chemotherapy.<sup>25,26</sup> Our confirmatory data on pre-treatment FSH as a predictor of contralateral GCT suggest this might also be used to target patients for adjuvant carboplatin.

In conclusion, we now provide mature data on a substantial, prospectively treated, patient cohort that confirm that carboplatin can be regarded as a standard management option for stage I seminoma alongside surveillance and RT, offering a less toxic alternative that at least delays and may substantially reduce the incidence of contralateral GCT. Each management approach has distinct advantages and disadvantages, and it is for individual patients and their families, together with their physicians, to decide which management approach they prefer, in the context of their healthcare system.

#### AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The author(s) indicated no potential conflicts of interest.

#### AUTHOR CONTRIBUTIONS

**Conception and design:** R. Timothy D. Oliver, Graham M. Mead, Sally P. Stenning

**Administrative support:** Philip Pollock

**Provision of study materials or patients:** R. Timothy D. Oliver, Graham M. Mead, Gordon J.S. Rustin, Johnathan K. Joffe, Nina Aass, Robert Coleman

**Collection and assembly of data:** Philip Pollock, Rhian Gabe, Sally P. Stenning

**Data analysis and interpretation:** R. Timothy D. Oliver, Graham M. Mead, Gordon J.S. Rustin, Rhian Gabe, Sally P. Stenning

**Manuscript writing:** R. Timothy D. Oliver, Graham M. Mead, Gordon J.S. Rustin, Johnathan K. Joffe, Nina Aass, Robert Coleman, Rhian Gabe, Sally P. Stenning

**Final approval of manuscript:** R. Timothy D. Oliver, Graham M. Mead, Gordon J.S. Rustin, Johnathan K. Joffe, Nina Aass, Robert Coleman, Philip Pollock, Rhian Gabe, Sally P. Stenning

## REFERENCES

- Powles T, Bhardwa J, Shamash J, et al: The changing presentation of germ cell tumours of the testis between 1983 and 2002. *BJU International* 95:1197-1200, 2005
- Warde P, Specht L, Horwich A, et al: Prognostic factors for relapse in stage I seminoma managed by surveillance: A pooled analysis. *J Clin Oncol* 20:4448-4452, 2002
- Martin JM, Panzarella T, Zwahlen DR, et al: Evidence-based guidelines for following stage I seminoma. *Cancer* 109:2248-2256, 2007
- Zagars GK, Ballo MT, Lee AK, et al: Mortality after cure of testicular seminoma. *J Clin Oncol* 22:640-647, 2004
- de Wit R, Fizazi K: Controversies in the management of clinical stage I testis cancer. *J Clin Oncol* 24:5482-5492, 2006
- Powles T, Robinson D, Shamash J, et al: The long-term risks of adjuvant carboplatin treatment for stage I seminoma of the testis. *Ann Oncol* 19:443-447, 2008
- Oliver T: Conservative management of testicular germ-cell tumors. *Nature Clinical Practice Urology* 4:550-560, 2007
- Fossa SD, Horwich A, Russell JM, et al: Optimal planning target volume for stage I testicular seminoma: A Medical Research Council randomized trial—Medical Research Council Testicular Tumor Working Group. *J Clin Oncol* 17:1146, 1999
- Jones WG, Fossa SD, Mead GM, et al: Randomized trial of 30 versus 20 Gy in the adjuvant treatment of stage I testicular seminoma: A report on Medical Research Council Trial TE18, European Organisation for the Research and Treatment of Cancer Trial 30942 (ISRCTN18525328). *J Clin Oncol* 23:1200-1208, 2005
- Oliver RTD, Mason MD, Mead GM, et al: Radiotherapy versus single-dose carboplatin in adjuvant treatment of stage I seminoma: A randomised trial. *Lancet* 366:293-300, 2005
- Calvert AH, Egorin MJ: Carboplatin dosing formulae: Gender bias and the use of creatinine-based methodologies. *Eur J Cancer* 38:11-16, 2002
- Warde PR, Chung P, Sturgeon J, et al: Should surveillance should be considered the standard of care in stage I seminoma? *J Clin Oncol* 23:16S, 2005 (suppl; abstr 4520)
- Reiter WJ, Kratzik CB, Brodowicz T, et al: Sperm analysis and serum follicle-stimulating hormone levels before and after adjuvant single agent carboplatin therapy for clinical stage I seminoma. *Urology* 52:117-119, 1998
- Mead GM, Fossa S, Oliver R, et al: Relapse patterns in 2466 stage I seminoma patients entered into Medical Research Council (MRC) randomised trials. *J Clin Oncol* 26:255s, 2008 (suppl; abstr 5020)
- Wanderas E, Fossa SD, Heilo A, et al: Serum follicle stimulating hormone: Predictor of cancer in the remaining testis in patients with unilateral testicular cancer. *Br J Urol* 66:315-317, 1990
- Harland SJ, Cook PA, Fossa SD, et al: Intratubular germ cell neoplasia of the contralateral testis in testicular cancer: Defining a high risk group. *Journal of Urology* 160:1353-1357, 1998
- Christensen TB, Daugaard G, Geertsen PF, et al: Effect of chemotherapy on carcinoma in situ of the testis. *Ann Oncol* 9:657-660, 1998
- Oliver RTD, Ong J, Berney D, et al: Testis conserving chemotherapy in germ cell cancer: Its potential to increase understanding of the biology and treatment of carcinoma-in-situ. *APMIS (Acta Pathologica, Microbiologica et Immunologica Scandinavica)* 111:86-92, 2003
- Heidenreich A, Weissbach L, Holtl W, et al: Organ sparing surgery for malignant germ cell tumor of the testis. *J Urol* 166:2161-2165, 2001
- Ng R: The reproductive imperative: A case report highlighting the possibility of using chemotherapy to conserve the testis in patients with testis cancer. *Clin Oncol* 9:334-337, 1997
- Dieckmann K, Bruggeboes B, Pichlmeier U, et al: Adjuvant treatment of clinical stage I seminoma: Is a single course of carboplatin sufficient? *Urology* 55:102-106, 2000
- Oliver T, Powles T, Somasundram U, et al: 22 year phase 1/2 study of single agent carboplatin in metastatic seminoma: could it have been accelerated by 72 hour PET scan response? *J Clin Oncol* 24:642s, 2006 (suppl; abstr 14565)
- Bokemeyer C, Kollmannsberger C, Stenning SP, et al: Metastatic seminoma treated with either single agent carboplatin or cisplatin-based combination chemotherapy: A pooled analysis of two randomised trials. *Br J Cancer* 91:683-687, 2004
- Krege S, Boergermann C, Baschek R, et al: Single agent carboplatin for CS IIA/B testicular seminoma: A phase II study of the German Testicular Cancer Study Group (GTCSG). *Ann Oncol* 17:276-280, 2006
- Aparicio J, Garcia del Muro X, Maroto P, et al: Multicentre study evaluating a dual policy of post-orchietomy surveillance and selective adjuvant single-agent carboplatin for patients with clinical stage I seminoma. *Ann Oncol* 14:867-872, 2003
- Aparicio J, Germa JR, Garcia del Muro X, et al: Risk-adapted management for patients with clinical stage I seminoma: The Second Spanish Germ Cell Cancer Cooperative Group study. *J Clin Oncol* 23:8717-8723, 2005
- Oliver T, Dieckmann K-P, Steiner H, et al: Pooled analysis of phase II reports of 2 versus 1 course of carboplatin as adjuvant for stage I seminoma. *J Clin Oncol* 23:395s, 2005 (suppl; abstr 4572)
- Huddart RA, Joffe JK: Preferred treatment for stage I seminoma: A survey of Canadian radiation oncologists. *Clin Oncol (R Coll Radiol)* 18:693-695, 2006
- Hoffman KE, Chen MH, Punglia RS, et al: Influence of year of diagnosis, patient age, and sociodemographic status on recommending adjuvant radiation treatment for stage I testicular seminoma. *J Clin Oncol* 26:3937-3942, 2008
- Krege S, Beyer J, Souchon R, et al: EGCCCC European Consensus Conference on Diagnosis and Treatment of Germ Cell Cancer: A report of the Second Meeting of the European Germ Cell Cancer Consensus group (EGCCCG)—Part I. *Eur Urol* 53:478-496, 2008
- Sodickson A, Baeyens PF, Andriole KP, et al: Recurrent CT, cumulative radiation exposure, and associated radiation-induced cancer risks from CT of adults. *Radiology* 251:175-184, 2009
- Medical Research Council Clinical Trials Unit: MRC CTU trial protocol: TE24/TRISST (Trial of Imaging and Surveillance in Seminoma Testis). London, United Kingdom, Medical Research Council Clinical Trials Unit, 2008

