

Locally Advanced Non-Small Cell Lung Cancer

Arya Amini MD

Department of Radiation Oncology



Case Presentation

- 52-year-old female, 60-pack-year smoking history
- T2a N2 M0 adenocarcinoma of the left lower lobe
- Presented with intermittent shortness of breath which became worse
- Chest x-ray: opacity in the left lung
- June 6, 2019, CT chest: 3.6 × 2.6 cm mass in the left lung with enlarged left hilar and suprahilar nodes
- June 11, 2019, endobronchial ultrasound (EBUS) and bx: squamous cell carcinoma
- June 13, 2019, PET / CT: hypermetabolic left lower lobe mass with uptake in upper paratracheal nodes
- June 28, 2019, MRI brain: negative

Imaging




6/13/2019

3. CT WB, 6. AC WB




5



Alpha: 60 %



Alpha: 60 %




Alpha: 60 %




Pathology

Lung, left lower lobe, core biopsy:

- Adenocarcinoma, solid type
- TTF-1 positive, p63 and Napsin A negative





High local recurrence in stage III patients after surgery alone

Study	No.	Stage	XRT dose	Survival	LRF w/o RT
Belgium (van Houtte et al. 1980)	202	I-III	60 Gy	43%	11%
LCSG (Weisenburger 1986)	230	II-III	50 Gy	40%	21%
CAMS (Feng et al. 2000)	317	II-III	60 Gy	41%	33%
Lille (Lafitte et al., 1996)	163	I	45-60 Gy	52%	17%
MRC LU11 (Stephens et al 1996)	308	II-III	40 Gy	25%	29%
Austria (Mayer et al. 1997)	155	I-III	50-56 Gy	20%	24%
GETCB (Dautzenberg et al. 1999)	720	I-III	60 Gy	43%	34%
Slovenia (Debevec et al. 1996)	74	III	30 Gy	20%	NA
Italy (Trodella et al. 2002)	104	I	50 Gy	58%	22%

Treatment for IIIA NSCLC

Pattern of Relapse Following Resection

Stage	Nodal Status	% chest	% distant
IA	N0	10	15
IB	N0	10	30
II	N1	12	40
IIIA	N2	15	60

Trimodality Therapy for Stage IIIA Intergroup 0139, “The Albain Study


429 pts w/ resectable biopsy proven N2
+ IIIA NSCLC randomized to:

Chemo/XRT 45 Gy + Surgery +
Consolidation chemo


vs.

Chemo/XRT 61 Gy + Consolidation
Chemo


Trimodality Therapy for Stage IIIA Intergroup 0139



Trimodality Therapy for Stage IIIA Intergroup 0139




Lobectomy – following neoadjuvant chemoradiation is reasonable treatment option, associated with survival >40%




Pneumonectomy – following neoadjuvant chemoradiation is associated with increased mortality

Trimodality Therapy for Stage IIIA Intergroup 0139




- Eradication of N2 disease by neoadjuvant therapy correlated strongly with survival following surgery
- 46% of patients were downstaged to N0
- Chemoradiation increased N2 clearance compared to neoadjuvant chemotherapy alone


Trimodality Therapy for Stage IIIA Intergroup 0139



Trimodality Therapy for Stage IIIA Intergroup 0139




Lobectomy – following neoadjuvant chemoradiation is reasonable treatment option, associated with survival >40%



Pneumonectomy – following neoadjuvant chemoradiation is associated with increased mortality

Trimodality Therapy for Stage IIIA Intergroup 0139



- Eradication of N2 disease by neoadjuvant therapy correlated strongly with survival following surgery
- 46% of patients were downstaged to N0
- Chemoradiation increased N2 clearance compared to neoadjuvant chemotherapy alone



Summary of Data on Radiation and Chemotherapy

- RT dose
 - RTOG 7301 (highest local control at 60 Gy)
- Sequential CRT vs RT alone
 - CALGB 8433 (7-yr OS higher, 13% vs 6%)
- Concurrent chemotherapy better than sequential
 - RTOG 9410 (5-yr OS higher with concurrent; improved LRC)
 - Meta-analysis
- Concurrent CRT +/- induction chemo
 - LAMP: no difference with induction

Perez et al. Cancer 1980 Jun; 45(11):2744-53.
Dillman et al. J Natl Cancer Inst. 1996 Sep; 88(17):1210-15.
Curran et al. J Natl Cancer Inst. 2011 Oct; 103(19):1452-60.
Auperin et al. JCO 2010 May; 28(13):2181-90.
Belani et al. JCO 2005 Sep; 23(55):5883-91.



Data for Chemotherapy Combined with Radiation

- RTOG 9410 (Curran et al, JNCI 2011)
 - Unresected stage III lung, randomized to 3 arms
 - 60 Gy RT -> sequential chemo (cis/vinb) (median OS 14.6 mo)
 - 60 Gy RT + concurrent chemo (cis/vinb) (median OS 17 mo)
 - 69.2 Gy Hyperfrac RT + concurrent chemo (cis/etop) (median OS 15.6 Gy)



Data for Chemotherapy Combined with Radiation

- RTOG 9410 (Curran et al, JNCI 2011)
 - Unresected stage III lung, randomized to 3 arms
 - 60 Gy RT -> sequential chemo (cis/vinb) (median OS 14.6 mo)
 - 60 Gy RT + concurrent chemo (cis/vinb) (median OS 17 mo)
 - 69.2 Gy Hyperfrac RT + concurrent chemo (cis/etop) (median OS 15.6 Gy)



RTOG 9410

Table 3. Patterns of failure*

Component of first failure	No. (%)		
	Arm 1 (n = 195)	Arm 2 (n = 195)	Arm 3 (n = 187)
Primary tumor	65 (33)	56 (29)	47 (25)
Thoracic lymph nodes (infield)	34 (17)	24 (12)	18 (10)
Thoracic lymph nodes (out of field)	4 (2)	8 (4)	3 (2)
Brain metastases	24 (12)	28 (14)	24 (13)
Other metastases	65 (33)	64 (33)	60 (32)
Infield only	59 (30)	49 (25)	38 (20)
Out of field only	67 (34)	73 (37)	69 (37)
Both infield and out of field	22 (11)	20 (10)	16 (9)

* Arm 1 was the sequential arm. Arm 2 used the same chemotherapy regimen as arm 1 with 60 Gy thoracic radiotherapy beginning on day 1. Arm 3 used cisplatin at 50 mg/m² on days 1, 8, 29, and 36 with oral etoposide at 50 mg twice daily for 10 weeks on days 1, 2, 5, and 6 with 69.6 Gy delivered as 1.2-Gy twice-daily fractions beginning on day 1

Arm 1: Sequential
Arm 2: Concurrent
Arm 3: HyperFx Conc


Total In-field failures as site of first failure ~30-40% (lower at 30-35% with concurrent chemoRT)

Gr3+ Esophagitis higher with concurrent chemoRT arms (4 vs 22 vs 45%)

No diff in late toxicity

RTOG 0617

- Randomized, 2x2 factorial phase 3
- Concurrent carbo (AUC 2)/paclitaxel (45 mg/m²) +/- CTX, 2c consolidation (AUC 6/200 mg/m²) +/- CTX
- 60 vs 74 Gy
- Median OS worse for 74 Gy, 20.3 vs 28.7 mo (HR 1.38, p=0.004)
- Median OS no different with CTX



RTOG 0617


- Not just a negative trial
- We learned
 - Higher dose may not be the answer
 - IMRT should be standard
 - Gr ≥ 3 pneumonitis, decrease heart dose,
 - Lung V20 associated with Gr ≥ 3 pneumonitis; V5 did not
 - Heart V40 \rightarrow predicted OS

PACIFIC Trial – The current standard of care for Unresectable Stage III NSCLC

- Purpose: evaluate the role of adjuvant durvalumab (anti-PDL1) after chemoradiation in stage III NSCLC
- Enrolled: stage III, unresectable; excluded grade 2+ pneumonitis from CRT or grade 2+ unresolved
- Treatment:
 - Platinum-based chemotherapy (etoposide, vinorelbine, vinorelbine & taxane [paclitaxel or docetaxel] or pemetrexed)
 - Concurrent RT (54-66 Gy)
 - Durvalumab 10 mg/kg or placebo q2 weeks for 1 year (2:1 randomization)

PACIFIC Trial - Results

- PFS with Durvalumab vs Placebo ($p < 0.001$)
 - Median: 16.8 vs 5.6 mo
 - 12-mo: 55.9% vs 35.3%
 - 18-mo: 44.2% vs 27.0%
- Median time to death or DM
 - 23.2 mo vs 14.6 mo ($p < 0.001$)
- **Analysis of OS was not planned at this interim**



PACIFIC Trial – Results




Figure 2. Subgroup Analysis of Prognostic Factors for Progression-free Survival in the Intention-to-Treat Population.

Progression-free survival was defined according to RECIST, version 1.1, and assessed by means of blinded independent central review. The hazard ratio and 95% confidence interval were not calculated for the complete response because this subgroup had less than 20 events. EGFR denotes epidermal growth factor receptor, and PD-L1 programmed death ligand 1.


Randomization to durvalumab or placebo within 14 days of completion of CRT: 0.39 (95% CI, 0.26-0.58) vs 0.63 (95% CI, 0.49-0.80) when randomized between 14-72 days

PACIFIC Trial – Adverse Events

- Grade 3 or 4 toxicity in 29.9% durvalumab and 26.1% placebo
 - Most common pneumonia
- Grade 5 in 4.4% durvalumab and 5.6% placebo
- Most frequent AEs leading to discontinuation of durvalumab and placebo were pneumonitis (6.3% vs 4.3%) and pneumonia (1.1% vs 1.3%)
- Durvalumab related toxicities (any grade): diarrhea (18.3% vs 18.8%), pneumonitis (12.6% vs 7.7%), rash (12.2% vs 7.3%) and pruritis (12.2% vs 7.7%)

PACIFIC Trial – OS Results


- Updated Results including **OS** (published Dec 2018) comparing durvalumab vs placebo
- 2-yr OS 66.3% vs 55.6%; (HR 0.68, 0.47-0.997; p=0.0025)
- 30.5% vs 26.1% had G3-4 toxicity; 15.4% vs 9.8% discontinued the trial 2/2 toxicity



Antonia et al. NEJM 2018 Dec; 379:2342-50.


PACIFIC Trial – Time to Death or DM

- Time to death longer with durvalumab, 28.3 vs 16.2 mo
- Frequency of newer lesions reduced, 22.5% vs 33.8%
- Reduced new brain metastases, 6.3% vs 11.8%




PACIFIC Trial – Significance of PD-L1

PFS



(C)


OS



Current and Future Points of Discussion

- Role of concurrent ICI with radiation treatment
- Dose of radiation: >60 Gy?, hypofractionation?
- How do we interpret data from ICI -> surgery trials or surgery + TKI when compared to CRT/ICI?
- Duration of ICI? Which ICI?
- ? of maintenance therapy selection after CRT for actionable mutation (EGFR/ALK) patients?

LAURA Trial



PACIFIC-2

- Phase 3 study of concurrent durvalumab and platinum-based chemoradiotherapy in patients with unresectable, stage III NSCLC
- 300 pts with unresectable stage III NSCLC will be randomized (2:1) to receive either durvalumab (intravenous 1500 mg) every 4 weeks (q4w) + cCRT, or placebo q4w + cCRT
- Primary endpoints are PFS and ORR

NRG LU-004

Cohort 1, n=6

MEDI4736 (durvalumab)[†] q4 weeks x 13 doses
+ ACRT 60 Gy in 15 fractions x 3 weeks (weeks 1-3)

Cohort 2, n=6


MEDI4736 (durvalumab)[†] q4 weeks x 13 doses
+ standard RT 60 Gy in 30 fractions x 6 weeks (weeks 1-6)

[†] MEDI4736 (durvalumab) begins 2 weeks (Day -14) before RT (+/- 48 hours); see [Section 5.1](#) for dosing details

EXPANSION COHORTS

After completing one of the Initial Safety Schedules of concurrent RT+MEDI4736 (durvalumab):

- If Cohort 1 only is deemed safe, all patients will be registered to Cohort 3.
- If Cohort 2 only is deemed safe, all patients will be registered to Cohort 4.
- **If both Cohorts 1 and 2 are deemed safe, patients will be randomized to either Cohort 3 or Cohort 4 with 1:1 randomization.**



[†] MEDI4736 (durvalumab) begins 2 weeks (Day -14) before RT (+/- 48 hours); see [Section 5.1](#) for dosing details