

Comparative genetic architectures of schizophrenia in East Asian and European populations

Armand González Escalante
Ana Sánchez Fernández

Index

Introduction


Samples

Results

1. Schizophrenia genetic associations in EAS population
2. Genetic effects across populations
3. Schizophrenia genetic associations from the meta-analysis of EAS and EUR
4. Population diversity and fine-mapping
5. Transferability of genetics across populations

Discussion

Introduction. Schizophrenia



Introduction. The limitations of only using European ancestry samples



- It **hinders** the **discovery of biological clues** about schizophrenia.
- **Associations** that can be **detected in one population** may **not be detected in others**.
- Impossible to determine if **differences** between populations **represent etiologic heterogeneity** on the illness.
- **Prediction accuracy of Polygenic Risk Scores (PRS) decays with increasing genetic divergence** between **the risk allele discovery** (i.e. Europeans) and **target datasets** (i.e. East Asians)

Samples. East Asian

Study	Case	Control	Chip	Design	Population	Raw data	Stage	X chr
IMH-1	856	946	I_1M	CC	Han Chinese	Y	1	Y
IMH-2	766	913	I_OZH	CC	Han Chinese	Y	1	Y
HNK-1	476	2018	I_610	CC	Han Chinese	Y	1	Y
JPN-1	547	540	A_SNP5.0	CC	Japanese	Y	1	Y
BIX-1	1045	2272	A_SNP6.0	CC	Han Chinese	Y	1	N
BIX-2	1021	1001	A_SNP6.0	CC	Han Chinese	Y	1	N
BIX-3	489	679	A_SNP6.0	CC	Han Chinese	Y	1	N
XJU-1	1846	947	I_OZH	CC	Han Chinese	Y	1	Y
UMC-1	2260	2241	I_Psyc	CC	Han Chinese	Y	1	Y
UWA-1	988	1001	I_Psyc	CC	Indonesia	Y	1	Y
BJM-1	1312	1987	I_OZH	CC	Han Chinese	Y	1	Y
TAI-1	1109	1109	I_Psyc	TRIO	Han Chinese	Y	1	Y
TAI-2	590	590	I_Psyc	TRIO	Han Chinese	Y	1	Y
KOR-1	687	492	A_KB	CC	Korean	Y	2	N
SIX-1	192	47	I_Psyc	CC	Han Chinese	Y	2	N
BIX-4	399	478	I_GSA	CC	Han Chinese	Y	2	N
BJM-2	746	1599	I_610	CC	Han Chinese	N	2	N
BJM-3	1595	1447	I_660W	CC	Han Chinese	N	2	N
BJM-4	710	680	I_OZH	CC	Han Chinese	N	2	N
BIX-5	5144	14375	A_SNP6.0, A_CHB1, I_1M	CC	Han Chinese	N	2	N
Total	22,778	35,362						



Cases (n): 22,778
Controls (n): 35,362

Samples. [European](#)

nature

Article | [Published: 22 July 2014](#)

Biological insights from 108 schizophrenia-associated genetic loci

[Schizophrenia Working Group of the Psychiatric Genomics Consortium](#)

[Nature](#) **511**, 421–427(2014) | [Cite this article](#)

35k Accesses | **3171** Citations | **1086** Altmetric | [Metrics](#)

Samples. European

Note	PI	PMID	Site	QC score	Array	Cases	Controls	Male	Tag
------	----	------	------	----------	-------	-------	----------	------	-----

•
•
•

PGC1	Sullivan, PF/Sklar P/Hultman C	23974872	Sweden (sw1)	3	A5.0	215	210	0.527	scz_swe1_eur
PGC1/New CC	Sullivan, PF/Sklar P/Hultman C	23974872	Sweden (sw234)	3	A6.0	1980	2274		scz_s234_eur
New CC	Sullivan, PF/Sklar P/Hultman C	23974872	Sweden (sw5)	3	omni	1764	2581	0.553	scz_swe5_eur
New CC	Sullivan, PF/Sklar P/Hultman C	23974872	Sweden (sw6)	3	omni	975	1145	0.543	scz_swe6_eur
New CC	Walters, J	21850710	Cardiff, UK (CogUK)	9	omni	530	678	0.554	scz_cou3_eur
New CC	Weinberger, D	11381111	NIMH CBDB	5	O25	133	269	0.547	scz_lie2_eur
New CC	Weinberger, D	11381111	NIMH CBDB	5	I550	497	389	0.627	scz_lie5_eur
PGC1	Werge, T	19571808	Denmark	8	I650	471	456	0.583	scz_denm_eur
Total EUR CC						32,405	42,221		
Trios	Kirov, G/Owen M	22083728	Bulgaria	8	A6.0	649	649	0.502	ms.scz_butr_eur
Trios	Levinson, D	22885689	Six countries	4	I650	516	516	0.556	ms.scz_lemu_eur
Trios	Kirov, G/Owen M	NP	Bulgaria	8	omni	70	70	0.595	ms.scz_uktr_eur
Total Trios						1,235	1,235		
East Asia	Iwata, N	20832056	Japan	3	A5.0	492	427	0.507	scz_jpn1_asn
East Asia	Liu, J	NP	Singapore (STCRP)	8	I1M	868	938		scz_tcr1_asn
East Asia	Sham, P	24043878	China	6	I550	476	2018	0.398	scz_hok2_asn
Total Asia CC						1,836	3,383		
Total Discovery						35,476	46,839		
Replication	Stefánsson, H	19571808	Iceland (SGENE+, deCODE)		ILMN*	628	65,312		N/A
Replication	Stefánsson, H	23164818	Non-Icelandic (SGENE+, deCODE)		ILMN*	885	924		N/A


Cases (n): 33,640
Controls (n): 43,456


Samples. Preprocessing

20 samples in total from EAS

- Quality control, phasing and imputation (**same method for EUR samples**) for **16** samples for which they had individual genotype data (raw data = Y)
- **Two** of the samples from these 16 were from a trio design so pseudo-controls were used.
- For the **4** remaining samples only summary statistics were available.


Two-stage study design


Discovery (13 samples)


Replication (7 samples)

Study	Case	Control	Chip	Design	Population	Raw data	Stage	X chr.
IMH-1	856	946	I_1M	CC	Han Chinese	Y	1	Y
IMH-2	766	913	I_OZH	CC	Han Chinese	Y	1	Y
HNK-1	476	2018	I_610	CC	Han Chinese	Y	1	Y
JPN-1	547	540	A_SNP5.0	CC	Japanese	Y	1	Y
BIX-1	1045	2272	A_SNP6.0	CC	Han Chinese	Y	1	N
BIX-2	1021	1001	A_SNP6.0	CC	Han Chinese	Y	1	N
BIX-3	489	679	A_SNP6.0	CC	Han Chinese	Y	1	N
XJU-1	1846	947	I_OZH	CC	Han Chinese	Y	1	Y
UMC-1	2260	2241	I_Psyc	CC	Han Chinese	Y	1	Y
UWA-1	988	1001	I_Psyc	CC	Indonesia	Y	1	Y
BJM-1	1312	1987	I_OZH	CC	Han Chinese	Y	1	Y
TAI-1	1109	1109	I_Psyc	TRIO	Han Chinese	Y	1	Y
TAI-2	590	590	I_Psyc	TRIO	Han Chinese	Y	1	Y
KOR-1	687	492	A_KB	CC	Korean	Y	2	N
SIX-1	192	47	I_Psyc	CC	Han Chinese	Y	2	N
BIX-4	399	478	I_GSA	CC	Han Chinese	Y	2	N
BJM-2	746	1599	I_610	CC	Han Chinese	N	2	N
BJM-3	1595	1447	I_660W	CC	Han Chinese	N	2	N
BJM-4	710	680	I_OZH	CC	Han Chinese	N	2	N
BIX-5	5144	14375	A_SNP6.0, A_CHB1, I_1M	CC	Han Chinese	N	2	N
Total	22,778	35,362						

Schizophrenia genetic associations in EAS population

METANALYSIS


Fixed-effect model with
inverse-variance weighting



ASSUMPTION

There exists a **true effect size** shared by all the studies

Schizophrenia genetic associations in EAS population




Schizophrenia genetic associations in EAS population

- **15 of 21 loci found had higher MAFs in EAS compared to EUR:**


Example: **rs374528934** (the top association) has a MAF of **45%** in EAS but only **0.7%** in EUR . It wasn't detected on the EUR association study.

This locus contains **CACNA2D2** that codes for a **protein** subunit associated to **childhood epilepsy** and to which the anticonvulsant medication **gabapentin** binds, suggesting a future **possible therapeutic investigation**.



Results. Genetic effects are consistent across populations

Linkage Disequilibrium Score Regression (LDSC) was used in order to quantify the **SNP heritability** for schizophrenia in both EUR and EAS populations and was found to be very similar, 0.24 ± 0.02 against 0.23 ± 0.03



Using the same set of variants the **genetic correlation** between schizophrenia at EUR and EAS was found to be very close to 1 ($r = 0.98 \pm 0.03$)

This implies that the **common variant genetic architecture of schizophrenia is highly consistent** across EUR and EAS samples.


Genetic effects are consistent across populations

Gene set enrichment analysis was also performed for EUR and EAS samples. There was **no significant difference** between gene set ranks($P = 0.72$, two-sided Wilcoxon test).


An **evaluation of the heterogeneity of schizophrenia genetic effects within EAS samples** was performed (Cochran's Q). None of the 21 associations showed significant heterogeneity across samples.

Using **PCA**, the samples were grouped into Northeast Asian, Southeast Asian and Indonesian subpopulations. A heterogeneity test (Cochran's Q) was then performed and found **no significant heterogeneity among the three subpopulations**.


a




b



c




d



Schizophrenia genetic associations from the meta-analysis of EAS and EUR

- Fixed effect meta-analysis **EAS + EUR**:
56,418 cases – 78,818 controls
- **208** independent **variants associated** with schizophrenia in **176 loci**
- **53 novel loci**
- 89 variants of the only EUR study remained significant




Trans-ethnicity bayesian fine-mapping

- **Fine-mapping** defines a ‘credible set’ of variants that contains the **causal variant** with certain probability
- **Bayes theorem** to calculate the probability of being causal for each associated variant in a locus


$$P(A|B) = \frac{P(B|A) \cdot P(A)}{P(B)}$$

- **Credible set:** variants for which sum of the probabilities $\geq 95\%$
- **Trans-ethnicity.** Heterogeneity was considered.

Causal variants: ↓ heterogeneity ↑ statistical significance




Trans-ethnicity bayesian fine-mapping




44 out of 59 associations were improved by trans-ancestry fine mapping.

Population diversity and fine-mapping



Transeferability of genetics across populations

- Comparing cause of liability variance across EUR and EAS populations
- Most of them had similar odd ratio but **different allele frequencies**
- Even if different ancestries have the same risk alleles and effect sizes ➡ **predictive power not equivalent**
- Evaluation of PRS in EAS with EAS and EUR as training data



Discussion

- This study compiles the **largest non-EUR psychiatric cohort**
- **Single population** studies → results **biased** toward higher **frequency alleles**. Including global samples increases discovery power
- Remarkable **genetic correlation** between EAS and EUR
- **Novel algorithm** for trans-ethnicity fine-mapping
- **Reduced performance predicting** PRS across populations due to different allele frequencies and LD patterns