

Continuous-time systems 1

March 2, 2015

- 1 Linear differential equations
- 2 Laplace transform
- 3 Solving LDEs with the Laplace transform

Outline

- 1 Linear differential equations
- 2 Laplace transform
- 3 Solving LDEs with the Laplace transform

Linear differential equations: definitions 1/2

Linear differential equations (LDE) are of the following form:

$$L[y(t)] = f(t),$$

where L is some linear operator.

Linear differential equations: definitions 1/2

Linear differential equations (LDE) are of the following form:

$$L[y(t)] = f(t),$$

where L is some linear operator.

The linear operator L is of the following form:

$$L_n(y) = \sum_{i=0}^n A_i(t) \frac{d^{n-i}y}{dt^{n-i}},$$

with given functions $A_{1:n}$.

Linear differential equations: definitions 1/2

Linear differential equations (LDE) are of the following form:

$$L[y(t)] = f(t),$$

where L is some linear operator.

The linear operator L is of the following form:

$$L_n(y) = \sum_{i=0}^n A_i(t) \frac{d^{n-i}y}{dt^{n-i}},$$

with given functions $A_{1:n}$.

The **order of a LDE** is the index of the highest derivative of y .

Linear differential equations: definitions 2/2

$$L_n(y) = \sum_{i=0}^n A_i(t) \frac{d^{n-i}y}{dt^{n-i}} = f(t).$$

- y is a scalar function \rightarrow **ordinary differential equation (ODE)**

Linear differential equations: definitions 2/2

$$L_n(y) = \sum_{i=0}^n A_i(t) \frac{d^{n-i}y}{dt^{n-i}} = f(t).$$

- y is a scalar function \rightarrow **ordinary differential equation (ODE)**
- y is a vector function \rightarrow **partial differential equation (PDE)**

Linear differential equations: definitions 2/2

$$L_n(y) = \sum_{i=0}^n A_i(t) \frac{d^{n-i}y}{dt^{n-i}} = f(t).$$

- y is a scalar function \rightarrow **ordinary differential equation (ODE)**
- y is a vector function \rightarrow **partial differential equation (PDE)**
- $f = 0 \rightarrow$ **homogeneous equation**
 \rightarrow solutions are called **complementary functions**

Linear differential equations: definitions 2/2

$$L_n(y) = \sum_{i=0}^n A_i(t) \frac{d^{n-i} y}{dt^{n-i}} = f(t).$$

- y is a scalar function \rightarrow **ordinary differential equation** (ODE)
- y is a vector function \rightarrow **partial differential equation** (PDE)
- $f = 0 \rightarrow$ **homogeneous equation**
 \rightarrow solutions are called **complementary functions**
- if $A_{0:n}(t)$ are constants (ie. not functions of time), the LDE is said to have **constant coefficients**

Example: radioactive decay 1/2

Let $N(t)$ be the number of radioactive atoms at time t , then:

$$\frac{dN(t)}{dt} = -kN(t),$$

for some constant $k > 0$.

Example: radioactive decay 1/2

Let $N(t)$ be the number of radioactive atoms at time t , then:


$$\frac{dN(t)}{dt} = -kN(t),$$

for some constant $k > 0$.

This is a first order homogeneous LDE with constant coefficients.

Example: radioactive decay 2/2

Decay of Carbon - 14



Solving homogeneous LDEs with constant coefficients 1/3

Solutions of LDEs must be of the form e^{zt} with $z \in \mathbb{C}$.

Solving homogeneous LDEs with constant coefficients 1/3

Solutions of LDEs must be of the form e^{zt} with $z \in \mathbb{C}$.

We assume an LDE with constant coefficients:

$$\sum_{i=0}^n A_i y^{(n-i)} = 0.$$

Solving homogeneous LDEs with constant coefficients 1/3

Solutions of LDEs must be of the form e^{zt} with $z \in \mathbb{C}$.

We assume an LDE with constant coefficients:

$$\sum_{i=0}^n A_i y^{(n-i)} = 0.$$

Replacing $y = e^{zt}$ leads to:

$$\sum_{i=0}^n A_i z^{n-i} e^{zt} = 0$$

Solving homogeneous LDEs with constant coefficients 1/3

Solutions of LDEs must be of the form e^{zt} with $z \in \mathbb{C}$.

We assume an LDE with constant coefficients:

$$\sum_{i=0}^n A_i y^{(n-i)} = 0.$$

Replacing $y = e^{zt}$ leads to:

$$\sum_{i=0}^n A_i z^{n-i} e^{zt} = 0$$

Dividing by e^{zt} yields the n th order **characteristic polynomial**:

$$F(z) = \sum_{i=0}^n A_i z^{n-i} = 0.$$

Solving homogeneous LDEs with constant coefficients 2/3

Characteristic equation:

$$F(z) = \sum_{i=0}^n A_i z^{n-i} = 0.$$

Solving homogeneous LDEs with constant coefficients 2/3

Characteristic equation:

$$F(z) = \sum_{i=0}^n A_i z^{n-i} = 0.$$

- 1 Solving the polynomial $F(z)$ yields n zeros z_1 to z_n .
- 2 Substituting a given zero z_i into e^{zt} gives a solution $e^{z_i t}$.

Solving homogeneous LDEs with constant coefficients 2/3

Characteristic equation:

$$F(z) = \sum_{i=0}^n A_i z^{n-i} = 0.$$

- 1 Solving the polynomial $F(z)$ yields n zeros z_1 to z_n .
- 2 Substituting a given zero z_i into e^{zt} gives a solution $e^{z_i t}$.

Homogeneous LDEs obey the superposition position:

→ any linear combination of solutions $e^{z_1 t}, \dots, e^{z_n t}$ is a solution

Solving homogeneous LDEs with constant coefficients 2/3

Characteristic equation:

$$F(z) = \sum_{i=0}^n A_i z^{n-i} = 0.$$

- 1 Solving the polynomial $F(z)$ yields n zeros z_1 to z_n .
- 2 Substituting a given zero z_i into e^{zt} gives a solution $e^{z_i t}$.

Homogeneous LDEs obey the superposition position:

- any linear combination of solutions $e^{z_1 t}, \dots, e^{z_n t}$ is a solution
- $e^{z_1 t}, \dots, e^{z_n t}$ form a basis of the solution space of the LDE

Solving homogeneous LDEs with constant coefficients 2/3

Characteristic equation:

$$F(z) = \sum_{i=0}^n A_i z^{n-i} = 0.$$

- ① Solving the polynomial $F(z)$ yields n zeros z_1 to z_n .
- ② Substituting a given zero z_i into e^{zt} gives a solution $e^{z_i t}$.

Homogeneous LDEs obey the superposition position:

→ any linear combination of solutions $e^{z_1 t}, \dots, e^{z_n t}$ is a solution

→ $e^{z_1 t}, \dots, e^{z_n t}$ form a basis of the solution space of the LDE

The specific linear combination depends on initial conditions.

Solving homogeneous LDEs with constant coefficients 3/3

Example:

$$y^{(4)}(t) - 2y^{(3)}(t) + 2y^{(2)}(t) - 2y^{(1)}(t) + y(t) = 0.$$

Solving homogeneous LDEs with constant coefficients 3/3

Example:

$$y^{(4)}(t) - 2y^{(3)}(t) + 2y^{(2)}(t) - 2y^{(1)}(t) + y(t) = 0.$$

This is a 4th order homogeneous LDE with constant coefficients.

Solving homogeneous LDEs with constant coefficients 3/3

Example:

$$y^{(4)}(t) - 2y^{(3)}(t) + 2y^{(2)}(t) - 2y^{(1)}(t) + y(t) = 0.$$

This is a 4th order homogeneous LDE with constant coefficients.

The corresponding characteristic equation:

$$F(z) = z^4 - 2z^3 + 2z^2 - 2z + 1 = 0.$$

Solving homogeneous LDEs with constant coefficients 3/3

Example:

$$y^{(4)}(t) - 2y^{(3)}(t) + 2y^{(2)}(t) - 2y^{(1)}(t) + y(t) = 0.$$

This is a 4th order homogeneous LDE with constant coefficients.

The corresponding characteristic equation:

$$F(z) = z^4 - 2z^3 + 2z^2 - 2z + 1 = 0.$$

The zeros of $F(z)$ are ($j = \sqrt{-1}$):

$$z_1 = j, \quad z_2 = -j, \quad z_{3,4} = 1.$$

Solving homogeneous LDEs with constant coefficients 3/3

Example:

$$y^{(4)}(t) - 2y^{(3)}(t) + 2y^{(2)}(t) - 2y^{(1)}(t) + y(t) = 0.$$

This is a 4th order homogeneous LDE with constant coefficients.

The corresponding characteristic equation:

$$F(z) = z^4 - 2z^3 + 2z^2 - 2z + 1 = 0.$$

The zeros of $F(z)$ are ($j = \sqrt{-1}$):

$$z_1 = j, \quad z_2 = -j, \quad z_{3,4} = 1.$$

These zeros correspond to the following basis functions t :

$$e^{jt}, \quad e^{-jt}, \quad e^t, \quad te^t.$$

Outline

- 1 Linear differential equations
- 2 Laplace transform
- 3 Solving LDEs with the Laplace transform

The Laplace transform

The Laplace transform of $f(t)$, for all real numbers $t \geq 0$:

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt.$$

The Laplace transform

The Laplace transform of $f(t)$, for all real numbers $t \geq 0$:

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt.$$

The parameter $s = \sigma + j\omega$ is the complex number frequency.

The Laplace transform

The Laplace transform of $f(t)$, for all real numbers $t \geq 0$:

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt.$$

The parameter $s = \sigma + j\omega$ is the complex number frequency.

The initial value theorem states $f(0^+) = \lim_{s \rightarrow \infty} sF(s)$.

The Laplace transform

The Laplace transform of $f(t)$, for all real numbers $t \geq 0$:

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt.$$

The parameter $s = \sigma + j\omega$ is the complex number frequency.

The initial value theorem states $f(0^+) = \lim_{s \rightarrow \infty} sF(s)$.

The final value theorem states $f(\infty) = \lim_{s \rightarrow 0} sF(s)$,
if all poles of $sF(s)$ are in the left half plane (ie. real part < 0).

Important properties of the Laplace transform

property	time domain	s -domain
linearity	$af(t) + bg(t)$	$aF(s) + bG(s)$

Important properties of the Laplace transform

property	time domain	s -domain
linearity	$af(t) + bg(t)$	$aF(s) + bG(s)$
differentiation	$f^{(1)}(t)$	$sF(s) - f(0)$

Important properties of the Laplace transform

property	time domain	s -domain
linearity	$af(t) + bg(t)$	$aF(s) + bG(s)$
differentiation	$f^{(1)}(t)$	$sF(s) - f(0)$
integration	$\int_0^t f(\tau)d\tau = (u * f)(t)$	$\frac{1}{s}F(s)$

Important properties of the Laplace transform

property	time domain	s -domain
linearity	$af(t) + bg(t)$	$aF(s) + bG(s)$
differentiation	$f^{(1)}(t)$	$sF(s) - f(0)$
integration	$\int_0^t f(\tau)d\tau = (u * f)(t)$	$\frac{1}{s}F(s)$
convolution	$(f * g)(t) = \int_0^t f(\tau)g(t - \tau)d\tau$	$F(s) \cdot G(s)$

Important properties of the Laplace transform

property	time domain	s -domain
linearity	$af(t) + bg(t)$	$aF(s) + bG(s)$
differentiation	$f^{(1)}(t)$	$sF(s) - f(0)$
integration	$\int_0^t f(\tau)d\tau = (u * f)(t)$	$\frac{1}{s}F(s)$
convolution	$(f * g)(t) = \int_0^t f(\tau)g(t - \tau)d\tau$	$F(s) \cdot G(s)$
time scaling	$f(at)$	$\frac{1}{a}F\left(\frac{s}{a}\right)$

Important properties of the Laplace transform

property	time domain	s -domain
linearity	$af(t) + bg(t)$	$aF(s) + bG(s)$
differentiation	$f^{(1)}(t)$	$sF(s) - f(0)$
integration	$\int_0^t f(\tau)d\tau = (u * f)(t)$	$\frac{1}{s}F(s)$
convolution	$(f * g)(t) = \int_0^t f(\tau)g(t - \tau)d\tau$	$F(s) \cdot G(s)$
time scaling	$f(at)$	$\frac{1}{a}F\left(\frac{s}{a}\right)$
time shifting	$f(t - a)u(t - a)$	$e^{-as}F(s)$

Important properties of the Laplace transform

property	time domain	s-domain
linearity	$af(t) + bg(t)$	$aF(s) + bG(s)$
differentiation	$f^{(1)}(t)$	$sF(s) - f(0)$
integration	$\int_0^t f(\tau)d\tau = (u * f)(t)$	$\frac{1}{s}F(s)$
convolution	$(f * g)(t) = \int_0^t f(\tau)g(t - \tau)d\tau$	$F(s) \cdot G(s)$
time scaling	$f(at)$	$\frac{1}{a}F\left(\frac{s}{a}\right)$
time shifting	$f(t - a)u(t - a)$	$e^{-as}F(s)$

with $u(t) = \int_{-\infty}^t \delta(\tau)d\tau$ (Heaviside) and $\delta(t)$ the Dirac delta.

Inverse Laplace transform

The inverse Laplace transform converts s -domain to time domain:

$$f(t) = \mathcal{L}^{-1}\{F(s)\} = \frac{1}{j2\pi} \int_{\gamma-jT}^{\gamma+jT} e^{st} F(s) ds.$$

Inverse Laplace transform

The inverse Laplace transform converts s -domain to time domain:

$$f(t) = \mathcal{L}^{-1}\{F(s)\} = \frac{1}{j2\pi} \int_{\gamma-jT}^{\gamma+jT} e^{st} F(s) ds.$$

Practically, the inverse Laplace transform takes two steps:

- 1 write $F(s)$ in terms of partial fractions
- 2 transform each term in the partial fraction based on tables of s/t -domain pairs (course notes p 4.32-4.33)

Outline

- 1 Linear differential equations
- 2 Laplace transform
- 3 Solving LDEs with the Laplace transform

Solving LDEs with the Laplace transform 1/3

The Laplace transform can be used to solve LDEs with given initial conditions (the previous approach gave us the basis functions).

Solving LDEs with the Laplace transform 1/3

The Laplace transform can be used to solve LDEs with given initial conditions (the previous approach gave us the basis functions).

This is done by using the following property (differentiation):

$$\mathcal{L}\{f^{(1)}\} = sF(s) - f(0),$$

$$\mathcal{L}\{f^{(2)}\} = s^2F(s) - sf(0) - f^{(1)}(0).$$

Solving LDEs with the Laplace transform 1/3

The Laplace transform can be used to solve LDEs with given initial conditions (the previous approach gave us the basis functions).

This is done by using the following property (differentiation):

$$\mathcal{L}\{f^{(1)}\} = sF(s) - f(0),$$

$$\mathcal{L}\{f^{(2)}\} = s^2F(s) - sf(0) - f^{(1)}(0).$$

Via induction, the Laplace transform of the n th order derivative:

$$\mathcal{L}\{f^{(n)}\} = s^n F(s) - \sum_{i=1}^n s^{n-i} f^{(n-i)}(0)$$

Solving LDEs with the Laplace transform 2/3

$$\mathcal{L}\{f^{(n)}\} = s^n F(s) - \sum_{i=1}^n s^{n-i} f^{(n-i)}(0)$$

Solving LDEs with the Laplace transform 2/3

$$\mathcal{L}\{f^{(n)}\} = s^n F(s) - \sum_{i=1}^n s^{n-i} f^{(n-i)}(0)$$

We want to solve the following LDE:

$$\sum_{i=0}^n A_i y^{(n-i)}(t) = f(t),$$

$$y^{(i)}(0) = c_i \quad \forall i = 0 \dots n.$$

Solving LDEs with the Laplace transform 2/3

$$\mathcal{L}\{f^{(n)}\} = s^n F(s) - \sum_{i=1}^n s^{n-i} f^{(n-i)}(0)$$

We want to solve the following LDE:

$$\sum_{i=0}^n A_i y^{(n-i)}(t) = f(t),$$

$$y^{(i)}(0) = c_i \quad \forall i = 0 \dots n.$$

Via the linearity of the Laplace transform:

$$\sum_{i=0}^n A_i \mathcal{L}\{y^{(n-i)}(t)\} = \mathcal{L}\{f(t)\}$$

Solving LDEs with the Laplace transform 3/3

$$\sum_{i=0}^n A_i \mathcal{L}\{y^{(n-i)}(t)\} = \mathcal{L}\{f(t)\} \quad (1)$$

$$\mathcal{L}\{f^{(n)}\} = s^n F(s) - \sum_{i=1}^n s^{n-i} f^{(n-i)}(0) \quad (2)$$

Solving LDEs with the Laplace transform 3/3

$$\sum_{i=0}^n A_i \mathcal{L}\{y^{(n-i)}(t)\} = \mathcal{L}\{f(t)\} \quad (1)$$

$$\mathcal{L}\{f^{(n)}\} = s^n F(s) - \sum_{i=1}^n s^{n-i} f^{(n-i)}(0) \quad (2)$$

Expanding Eq. (2) into (1) yields:

$$Y(s) \sum_{i=0}^n A_i s^i - \sum_{i=1}^n \sum_{j=1}^i A_i s^{i-j} y^{j-1}(0) = F(s)$$

Solving LDEs with the Laplace transform 3/3

$$\sum_{i=0}^n A_i \mathcal{L}\{y^{(n-i)}(t)\} = \mathcal{L}\{f(t)\} \quad (1)$$

$$\mathcal{L}\{f^{(n)}\} = s^n F(s) - \sum_{i=1}^n s^{n-i} f^{(n-i)}(0) \quad (2)$$

Expanding Eq. (2) into (1) yields:

$$Y(s) \sum_{i=0}^n A_i s^i - \sum_{i=1}^n \sum_{j=1}^i A_i s^{i-j} y^{j-1}(0) = F(s)$$

The solution in the time domain is obtained via the inverse Laplace transform: $y(t) = \mathcal{L}^{-1}\{Y(s)\}$.