

Some statistical insights into PINNs

CRM Spring School, 2025



Work team

2 / 38



Nathan Doumèche



Gérard Biau



Claire Boyer

Statistical model: $Y = f^*(\mathbf{X}) + \varepsilon$

Goal: estimate f^* using

- ▶ Supervised learning: an i.i.d. training sample $(\mathbf{X}_i, Y_i)_{1 \leqslant i \leqslant n}$
- ▶ Physical modeling: a prior knowledge

$$\mathcal{D}_k(f^*, \cdot) \simeq 0, \quad 1 \leqslant k \leqslant M,$$


+ boundary/initial conditions for f^*

- ▶ Neural networks

- ▶ Physics-Informed Neural Networks (PINNs)



PINNs in a nutshell

4 / 38



Example: Blood flow in an aneurysm



5 / 38



Modeling the blood flow

6 / 38

[Arzani et al., 2021]



Goal: estimate the blood flow $f = (f_x, f_y, P)$

Navier-Stokes equations:


- ▶ $\mathcal{D}_1(f \cdot) = f_x \partial_x f_x + f_y \partial_y f_x - \partial_{x,x}^2 f_x - \partial_{y,y}^2 f_x + \partial_x P$
- ▶ $\mathcal{D}_2(f, \cdot) = f_x \partial_x f_y + f_y \partial_y f_y - \partial_{x,x}^2 f_y - \partial_{y,y}^2 f_y + \partial_y P$
- ▶ $\mathcal{D}_3(f, \cdot) = \partial_x f_x + \partial_y f_y$

(Incomplete) boundary conditions:

- ▶ $(f_x, f_y) = 0$ on the boundaries of the vessel
- ▶ Unknown inflow and outflow


Example: Heat transfer in uranium bundles

7 / 38



Modeling the heat transfer

8 / 38



Goal: estimate the temperature T on the bundles

Navier-Stokes and diffusion equations on $f = (f_x, f_y, P, T)$:

- ▶ $\mathcal{D}_1(f, \cdot) = f_x \partial_x f_x + f_y \partial_y f_x - \partial_{x,x}^2 f_x - \partial_{y,y}^2 f_x + \partial_x P$
- ▶ $\mathcal{D}_2(f, \cdot) = f_x \partial_x f_y + f_y \partial_y f_y - \partial_{x,x}^2 f_y - \partial_{y,y}^2 f_y + \partial_y P$
- ▶ $\mathcal{D}_3(f, \cdot) = \partial_x f_x + \partial_y f_y$
- ▶ $\mathcal{D}_4(f, \cdot) = f_x \partial_x T + f_y \partial_y T - \partial_{x,x}^2 T - \partial_{y,y}^2 T$

(Incomplete) boundary conditions:

- ▶ $(f_x, f_y) = 0$ and $T = 0$ on the physical boundaries
- ▶ Inflow with $f_x = 1$, $f_y = 0$, and $T = 0$
- ▶ Outflow with $\partial_x T = 0$

The PDE solver case

9 / 38


Specificity: no data Y_i and exact modeling

Example: the nonlinear Schrödinger PDE

[Raissi et al., 2019]

$$i\partial_t f + 0.5\partial_{x,x}^2 f + |f|^2 u = 0$$

Periodic boundary conditions and initial condition: $f(x, 0) = 2 / \cosh(x)$



Hybrid modeling problems:

- ▶ Improve **imperfect/incomplete** physical models with data
- ▶ Conversely, provide **interpretability** and **extrapolation** in ML

PDE solvers:

- ▶ Rely on **complex triangulations** of the domain
- ▶ Prone to the **curse of the dimension**

PINNs:

- ▶ A modern and efficient ML tool for both problems
- ▶ Natural **implementation** in the deep learning framework

Our objective


To better understand the **capabilities** and **limitations** of PINNs

1. Hybrid modeling
2. Consistency of the risk
3. Strong convergence
4. Numerical illustrations

1. Hybrid modeling
2. Consistency of the risk
3. Strong convergence
4. Numerical illustrations

Geometry of the problem


13 / 38



- ▶ $\Omega \subseteq \mathbb{R}^{d_1}$: the **bounded set** on which the problem is posed
- ▶ $f^* : \Omega \rightarrow \mathbb{R}^{d_2}$: the **unknown** target function
- ▶ Differential operators $\mathcal{D}_k(f^*, \cdot) \simeq 0$ on Ω , $1 \leq k \leq M$
- ▶ $\partial\Omega$: the boundary of $\Omega \Rightarrow$ often **not C^1** but **Lipschitz**
- ▶ Dirichlet conditions: $f^*(x) \simeq h(x)$ on $E \subseteq \partial\Omega$
- ▶ Possible extensions to other types of boundary/initial conditions

A general framework: 3 samplings

14 / 38




- ▶ Training sample $(\mathbf{X}_1, Y_1), \dots, (\mathbf{X}_n, Y_n) \in \Omega \times \mathbb{R}^{d_2}$ (**unknown** distribution)
- ▶ Boundary/initial sample $\mathbf{X}_1^{(e)}, \dots, \mathbf{X}_{n_e}^{(e)} \in E \subseteq \partial\Omega$ (**chosen** distribution)
- ▶ Collocation points $\mathbf{X}_1^{(r)}, \dots, \mathbf{X}_{n_r}^{(r)} \in \Omega$ (**uniform** distribution)

Empirical risk function

$$R_{n,n_e,n_r}(f_\theta) = \underbrace{\frac{1}{n} \sum_{i=1}^n \|f_\theta(\mathbf{X}_i) - Y_i\|_2^2}_{\text{data-fidelity}} + \underbrace{\frac{\lambda_e}{n_e} \sum_{j=1}^{n_e} \|f_\theta(\mathbf{X}_j^{(e)}) - h(\mathbf{X}_j^{(e)})\|_2^2}_{\text{boundary conditions}}$$
$$+ \underbrace{\frac{\lambda_r}{n_r} \sum_{k=1}^M \sum_{\ell=1}^{n_r} \mathcal{D}_k(f_\theta, \mathbf{X}_\ell^{(r)})^2}_{\text{PDEs}}$$

- ▶ $\text{NN}_H(D)$: the set of neural networks with H hidden layers of width D
- ▶ $\text{NN}_H = \cup_D \text{NN}_H(D)$
- ▶ θ : parameter of the neural network
- ▶ tanh: activation function
- ▶ $f_\theta \in C^\infty(\bar{\Omega}, \mathbb{R}^{d_2})$



Minimizing sequence

We denote by $(\hat{\theta}(p, n_e, n_r, D))_{p \in \mathbb{N}}$ any minimizing sequence, i.e.,

$$\lim_{p \rightarrow \infty} R_{n, n_e, n_r}(f_{\hat{\theta}(p, n_e, n_r, D)}) = \inf_{f_\theta \in \text{NN}_H(D)} R_{n, n_e, n_r}(f_\theta).$$

- ▶ The training of PINNs relies on the backpropagation algorithm

Hybrid modeling

- ▶ Statistical properties of PINNs
- ▶ Impact of the physical model
- ▶ Tuning of the PINN hyperparameters

PDE solver

- ▶ Reconstruction of the solution f^* of a PDE system
- ▶ Curse of the dimension

Proposition

Let $\Omega \subseteq \mathbb{R}^{d_1}$ be a bounded Lipschitz domain and $K \in \mathbb{N}$. Then, for any function $f \in C^\infty(\bar{\Omega}, \mathbb{R}^{d_2})$, there exists a sequence $(f_p)_{p \in \mathbb{N}} \in \text{NN}_H$ such that $\lim_{p \rightarrow \infty} \|f - f_p\|_{C^K(\Omega)} = 0$.

- ▶ Valid for bounded Lipschitz domains + $C^K(\Omega)$ norm
- ▶ Generalization of De Ryck et al. (2021)
- ▶ In line with practical applications, where $D \gg H$
- ▶ Key property to solve PDE systems

1. Hybrid modeling
2. Consistency of the risk
3. Strong convergence
4. Numerical illustrations

Theoretical risk

$$\begin{aligned}\mathcal{R}_n(f) &= \frac{1}{n} \sum_{i=1}^n \|f(\mathbf{X}_i) - Y_i\|_2^2 + \lambda_e \mathbb{E} \|f(\mathbf{X}^{(e)}) - h(\mathbf{X}^{(e)})\|_2^2 \\ &\quad + \frac{\lambda_r}{|\Omega|} \sum_{k=1}^M \int_{\Omega} \mathcal{D}_k(f, \mathbf{x})^2 d\mathbf{x}\end{aligned}$$


A natural requirement: Risk-consistency

$$\lim_{n_e, n_r \rightarrow \infty} \lim_{p \rightarrow \infty} \mathcal{R}_n(f_{\hat{\theta}(p, n_e, n_r, D)}) \stackrel{?}{=} \inf_{u \in \text{NN}_H(D)} \mathcal{R}_n(u)$$

- Warning: possible overfitting

Overfitting: hybrid modeling

- ▶ **Observations:** $Y_i = f^*(\mathbf{X}_i) + \varepsilon_i$
- ▶ **Goal:** estimate the trajectory f^* on $\Omega =]0, 1[$
- ▶ **Model (dynamics with friction):** $\mathcal{D}(f, \mathbf{x}) = f''(\mathbf{x}) + f'(\mathbf{x})$




- ▶ **Overfitting:** $R_{n,n_r} = 0$ but $\mathcal{R}_n = \infty$

Overfitting: PDE solver


22 / 38

- ▶ Heat equation: $\mathcal{D}(f, \mathbf{x}) = \partial_t f(\mathbf{x}) - \partial_{x,x}^2 f(\mathbf{x}) + \text{boundary/initial conditions}$
- ▶ Goal: reconstruct the solution f^* on $\Omega =]-1, 1[\times]0, 1[$

--- Initial and boundary conditions



--- Initial and boundary conditions



- ▶ Overfitting: $R_{n_e, n_r} = 0$ but $\mathcal{R} = \infty$

Proposition

There exists a constant $C_{K,H} > 0$ such that

$$\|f_\theta\|_{C^K(\mathbb{R}^{d_1})} \leq C_{K,H}(D+1)^{HK+1}(1 + \|\theta\|_2)^{HK}\|\theta\|_2.$$

Ridge PINNs

$$R_{n,n_e,n_r}^{(\text{ridge})}(f_\theta) = R_{n,n_e,n_r}(f_\theta) + \lambda_{(\text{ridge})}\|\theta\|_2^2$$

We denote by $(\hat{\theta}_{(p,n_e,n_r,D)}^{(\text{ridge})})_{p \in \mathbb{N}}$ a minimizing sequence of this risk.

- Implemented in standard DL libraries via weight decay

Example: the Navier-Stokes equations on $f = (f_x, f_y, P)$:

- ▶ $\mathcal{D}_1(f, \cdot) = f_x \partial_x f_x + f_y \partial_y f_x - \partial_{x,x}^2 f_x - \partial_{y,y}^2 f_x + \partial_x P$
- ▶ $\mathcal{D}_1(f, \cdot) = \mathcal{P}(f_x, \partial_x f_x, \partial_{x,x}^2 f_x, \partial_y f_x, \partial_{y,y}^2 f_x, f_y, \partial_x P)$
- ▶ $\mathcal{P}(Z_1, Z_2, Z_3, Z_4, Z_5, Z_6, Z_7) = Z_1 Z_2 + Z_6 Z_4 - Z_3 - Z_5 + Z_7$
- ▶ The coefficient in front of the monomial $Z_1 Z_2$ is 1 $\in C^\infty(\bar{\Omega}, \mathbb{R}^{d_2})$
- ▶ Warning: $\deg \mathcal{D}_1 = 3$ but $\deg \mathcal{P} = 2$

Polynomial operator

An operator $\mathcal{D}(u, \cdot)$ is polynomial if it can be expressed as a polynomial in u and its derivatives, with smooth functions as coefficients.

- ✓ Linear PDEs (e.g., advection, heat, and Maxwell)
- ✓ Some nonlinear PDEs (e.g., Blasius, Burger, and Navier-Stokes)

Assumptions:

- ▶ The condition function h is Lipschitz
- ▶ $\mathcal{D}_1, \dots, \mathcal{D}_M$ are polynomial operators

Theorem

With a ridge hyperparameter of the form

$$\lambda_{(\text{ridge})} = \min(n_e, n_r)^{-\kappa}, \quad \kappa = \frac{1}{12 + 4H(1 + (2 + H) \max_k \deg(\mathcal{D}_k))},$$

one has, almost surely,

$$\lim_{n_e, n_r \rightarrow \infty} \lim_{p \rightarrow \infty} \mathcal{R}_n(f_{\hat{\theta}(\text{ridge})}(p, n_e, n_r, D)) = \inf_{u \in \text{NN}_H(D)} \mathcal{R}_n(u)$$

and

$$\lim_{D \rightarrow \infty} \lim_{n_e, n_r \rightarrow \infty} \lim_{p \rightarrow \infty} \mathcal{R}_n(f_{\hat{\theta}(\text{ridge})}(p, n_e, n_r, D)) = \inf_{u \in C^\infty(\bar{\Omega}, \mathbb{R}^{d_2})} \mathcal{R}_n(u).$$

- ▶ Ridge regularization prevents overfitting of PINNs
- ▶ The decay rate of $\lambda_{(\text{ridge})} = \min(n_e, n_r)^{-\kappa}$ does not depend on the dimension d_1 of Ω
- ▶ $\lambda_{(\text{ridge})}$ can be tuned by monitoring the overfitting gap

1. Hybrid modeling
2. Consistency of the risk
3. Strong convergence
4. Numerical illustrations

- ✓ Ridge PINNs are risk-consistent

Question

Is this sufficient to have $\lim_{D, n_e, n_r, p \rightarrow \infty} f_{\hat{\theta}(\text{ridge})}(p, n_e, n_r, D) = f^*$ in $L^2(\Omega)$?

Answer: No

Let $\Omega =]0, 1[^2$, $h(x, 0) = 1$, $h(0, t) = 1$, and $\mathcal{D}(f, \cdot) = \partial_x f + \partial_t f$. Then, for any $(X_i, Y_i)_{1 \leq i \leq n}$, there exists $(f_p)_{p \in \mathbb{N}} \in \text{NN}_H(2n)$ such that

$$\lim_{p \rightarrow \infty} \mathcal{R}_n(f_p) = 0,$$

but $\lim_{p \rightarrow \infty} f_p = 1$ in $L^2(\Omega)$ (independently of f^*).

- ✗ KO if imperfect modeling
- ✓ Possible solution: Sobolev regularization



Weak derivatives

A function $v \in L^2(\Omega, \mathbb{R}^{d_2})$ is the α -th weak derivative of $u \in L^2(\Omega, \mathbb{R}^{d_2})$ if, for any $\varphi \in C^\infty(\bar{\Omega}, \mathbb{R}^{d_2})$ with compact support in Ω , one has

$$\int_{\Omega} \langle v, \varphi \rangle = (-1)^{|\alpha|} \int_{\Omega} \langle u, \partial^{\alpha} \varphi \rangle.$$

Notation: $v = \partial^{\alpha} u$.

Sobolev spaces

$H^m(\Omega, \mathbb{R}^{d_2})$ is the space of all functions $u \in L^2(\Omega, \mathbb{R}^{d_2})$ such that $\partial^{\alpha} u$ exist for all $|\alpha| \leq m$. This space is naturally endowed with the norm

$$\|u\|_{H^m(\Omega)}^2 = \frac{1}{|\Omega|} \sum_{|\alpha| \leq m} \int_{\Omega} \|\partial^{\alpha} u\|_2^2.$$

- ▶ $C^m(\bar{\Omega}, \mathbb{R}^{d_2}) \subseteq H^m(\Omega, \mathbb{R}^{d_2})$
- ▶ Standard derivatives \leftrightarrow weak derivatives

Sobolev-regularized risks

- Empirical risk:

$$R_{n,n_e,n_r}^{(\text{reg})}(f_\theta) = R_{n,n_e,n_r}(f_\theta) + \lambda_{(\text{ridge})} \|\theta\|_2^2 + \frac{\lambda_t}{n_r} \sum_{\ell=1}^{n_r} \sum_{|\alpha| \leq m+1} \|\partial^\alpha f_\theta(\mathbf{X}_\ell^{(r)})\|_2^2$$

- Minimizing sequence: $(\hat{\theta}^{(\text{reg})}(p, n_e, n_r, D))_{p \in \mathbb{N}}$
- Theoretical risk:

$$\mathcal{R}_n^{(\text{reg})}(f) = \mathcal{R}_n(f) + \lambda_{(\text{sob})} \|f\|_{H^{m+1}(\Omega)}^2$$

- The Sobolev regularization is straightforward to implement in the PINN framework with $\mathcal{D}_\alpha(f, \cdot) = \partial^\alpha f$
- Computational scalability via the backpropagation algorithm
- Coercivity of the risk

Theorem (Linear PDE systems)

Assume that there exists a unique solution $f^* \in H^{m+1}(\Omega, \mathbb{R}^{d_2})$ to the PDE system, where $m \geq \max_k \deg(\mathcal{D}_k)$. Thus, taking

$$\lambda_{(\text{ridge})} = \min(n_e, n_r)^{-\kappa}, \quad \kappa = \frac{1}{12 + 4H(1 + (2 + H)(m + 2))},$$

one has, almost surely,

$$\lim_{\lambda_t \rightarrow 0} \lim_{D \rightarrow \infty} \lim_{n_e, n_r \rightarrow \infty} \lim_{p \rightarrow \infty} \|f_{\hat{\theta}^{(\text{reg})}(p, n_e, n_r, D, \lambda_t)} - f^*\|_{H^m(\Omega)} = 0.$$

- ▶ The parameters m and $\lambda_{(\text{ridge})}$ do not depend on d_1
- ▶ The convergence is in $H^m(\Omega)$ for the penalty $\|u\|_{H^{m+1}(\Omega)}^2$
- ▶ Tools: Lax Milgram + functional analysis (weak topology)

Strong convergence: hybrid modeling

Physics inconsistency

For any $f \in H^{m+1}(\Omega, \mathbb{R}^{d_2})$, the **physics inconsistency** of f is defined by

$$\text{PI}(f) = \lambda_e \mathbb{E} \|u(\mathbf{X}^{(e)}) - h(\mathbf{X}^{(e)})\|_2^2 + \frac{\lambda_r}{|\Omega|} \sum_{k=1}^M \int_{\Omega} \mathcal{D}_k(u, \mathbf{x})^2 d\mathbf{x}.$$

Theorem (Linear PDE systems)

Assume that $f^* \in H^{m+1}(\Omega, \mathbb{R}^{d_2})$ for some $m \geq \max(\lfloor d_1/2 \rfloor, K)$. Let $\lambda_{e/r} = \log(n)/\sqrt{n}$ and $\lambda_{(\text{sob})} = 1/\sqrt{n}$. Then

$$\lim_{D \rightarrow \infty} \lim_{n_e, n_r \rightarrow \infty} \lim_{p \rightarrow \infty} \mathbb{E} \int_{\Omega} \|f_{\hat{\theta}^{(\text{reg})}(p, n_e, n_r, D)}^{(n)} - f^*\|_2^2 d\mu_{\mathbf{x}} \lesssim \frac{\log^2(n)}{n^{1/2}}$$

$$\text{and } \lim_{D \rightarrow \infty} \lim_{n_e, n_r \rightarrow \infty} \lim_{p \rightarrow \infty} \mathbb{E}(\text{PI}(f_{\hat{\theta}^{(\text{reg})}(p, n_e, n_r, D)}^{(n)})) \leq \text{PI}(f^*) + \underset{n \rightarrow \infty}{o}(1).$$

- Conclusion: statistical accuracy + physical consistency

1. Hybrid modeling
2. Consistency of the risk
3. Strong convergence
4. Numerical illustrations


Setting

Regression model:


- ▶ $Y = f^*(\mathbf{X}) + \mathcal{N}(0, 10^{-2})$
- ▶ $f^*(x, t) = \exp(t - x) + 0.1 \cos(2\pi x)$ on $\Omega =]0, 1[^2$
- ▶ $((x_i, t_i), Y_i)_{1 \leq i \leq n}$ for $0 < t_i < 0.5$

Advection model:

- ▶ $\mathcal{D}(f, \cdot) = \partial_x f + \partial_t f$
- ▶ $h(x, 0) = \exp(-x)$ and $h(0, t) = \exp(t)$
- ▶ $f_{\text{model}}(x, t) = \exp(t - x)$



f_{model}




f^*

Monitoring the risks


35 / 38

- ▶ Stability of the empirical risk $R_{n,n_e,n_r}^{(\text{reg})} \Rightarrow p \simeq \infty$
- ▶ Overfitting gap $\text{OG}_{n,n_e,n_r} = |R_{n,n_e,n_r}^{(\text{ridge})} - \mathcal{R}_n|$
⇒ choose the lowest possible $\lambda_{(\text{ridge})}$
- ▶ Illustration with $n = 10$





Result for $n = 10^3$

36 / 38



$f_{\text{model}}, f^*, \text{ and regularized PINN estimator}$

- ▶ Convergence on $\text{supp}(\mu_x) =]0, 1[\times]0, 0.5[$
- ▶ The regularized PINN follows the advection model (constant on the characteristics $x = t + \text{cst}$)
- ▶ Flattening effect of the Sobolev regularization on $\Omega \setminus \text{supp}(\mu_x)$



As predicted by the theory:

- ▶ The convergence rate is less than -0.5
- ▶ The regularized PINN is more accurate than f_{model} for $n > 10$
- ▶ The physics inconsistency is bounded by the modeling error $\text{PI}(f^*)$

- ▶ Statistical and PDE models can successfully complement each other
- ▶ Risk-consistency and strong convergence are guaranteed, with an implementable regularization
- ▶ PINNs are an interesting approach to scientific computing

Thank you!

- ☞ Doumèche, Biau, Boyer - Bernoulli 2025 [arXiv] [pdf] [supp]
- ☞ Code available [here](#)