

Analysis output: all-atom contacts and geometry for DIAPH1_762-1215H.pdb

Summary statistics

	Clashscore, all atoms:	5.91		91 st percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein	Poor rotamers	30	7.28%	Goal: <1%
	Ramachandran outliers	15	3.32%	Goal: <0.05%
	Ramachandran favored	413	91.37%	Goal: >98%
	MolProbity score [^]	2.49		48 th percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	0	0.00%	Goal: 0
	Bad backbone bonds:	9 / 3752	0.24%	Goal: 0%
	Bad backbone angles:	3 / 5025	0.06%	Goal: <0.1%

In the two column results, the left column gives the raw count, right column gives the percentage.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 0.26 points.

Multi-criterion visualizations



View (435 Kb)

1 of 2 6/13/14, 9:35 AM

^{* 100&}lt;sup>th</sup> percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

Single-criterion visualizations

- Clash list (1.7 Kb): View
- Ramachandran plot kinemage (432 Kb): View in KiNG | Download
- Ramachandran plot PDF (1.7 Mb): View
- Cβ deviation scatter plot (37 Kb): View in KiNG | Download



About MolProbity | Website for the Richardson Lab | Using ecloud x-H | Internal reference 4.1-537

2 of 2