



# Analysis output: all-atom contacts and geometry for HGF\_305-470H.pdb

## Summary statistics

All-Atom Contacts	Clashscore, all atoms:	50.06		4 <sup>th</sup> percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	0	0.00%	Goal: <1%
	Ramachandran outliers	5	3.05%	Goal: <0.05%
	Ramachandran favored	143	87.20%	Goal: >98%
	MolProbity score^	2.79		32 <sup>nd</sup> percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	4	2.63%	Goal: 0
	Bad backbone bonds:	13 / 1392	0.93%	Goal: 0%
	Bad backbone angles:	23 / 1894	1.21%	Goal: <0.1%

In the two column results, the left column gives the raw count, right column gives the percentage.

\* 100<sup>th</sup> percentile is the best among structures of comparable resolution; 0<sup>th</sup> percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

<sup>^</sup> MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 1.97 points.

## Multi-criterion visualizations



[View](#) (169 Kb)

## Single-criterion visualizations

- **Clash list** (2.8 Kb): [View](#)
- **Ramachandran plot kinemage** (415 Kb): [View in KiNG](#) | [Download](#)

- **Ramachandran plot PDF** (1.7 Mb): [View](#)
- **C $\beta$  deviation scatter plot** (19 Kb): [View in KiNG](#) | [Download](#)

[Continue >](#)

---

[About MolProbity](#) | [Website for the Richardson Lab](#) | [Using ecloud x-H](#) | [Internal reference 4.1-537](#)