



Analysis output: all-atom contacts and geometry for GPSM2_594-648.pdb_4_1H.pdb

Summary statistics

All-Atom Contacts	Clashscore, all atoms:	0.58		99 th percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	0	0.00%	Goal: <1%
	Ramachandran outliers	0	0.00%	Goal: <0.05%
	Ramachandran favored	98	92.45%	Goal: >98%
	MolProbity score [^]	1.16		99 th percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	5	4.81%	Goal: 0
	Bad backbone bonds:	0 / 894	0.00%	Goal: 0%
	Bad backbone angles:	4 / 1206	0.33%	Goal: <0.1%

In the two column results, the left column gives the raw count, right column gives the percentage.

* 100th percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

Multi-criterion visualizations



[View \(106 Kb\)](#)

Single-criterion visualizations

- **Clash list** (217 bytes): [View](#)
- **Ramachandran plot kinemage** (411 Kb): [View in KiNG](#) | [Download](#)
- **Ramachandran plot PDF** (1.7 Mb): [View](#)
- **C β deviation scatter plot** (17 Kb): [View in KiNG](#) | [Download](#)

Continue >

About [MolProbity](#) | Website for [the Richardson Lab](#) | Using ecloud x-H | Internal reference 4.1-537