

Analysis output: all-atom contacts and geometry for MYO7A_993_1686H.pdb

Summary statistics

Contacts	Clashscore, all atoms:	29.16		16 th percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	13	2.16%	Goal: <1%
	Ramachandran outliers	8	1.16%	Goal: <0.05%
	Ramachandran favored	651	94.08%	Goal: >98%
	MolProbity score [^]	2.60		41 st percentile [*] (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	6	0.92%	Goal: 0
	Bad backbone bonds:	42 / 5640	0.74%	Goal: 0%
	Bad backbone angles:	79 / 7642	1.03%	Goal: <0.1%

In the two column results, the left column gives the raw count, right column gives the percentage.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* fix 1 bad rotamers and improve your clashscore by 0.73 points.

Multi-criterion visualizations

1 of 2

^{* 100&}lt;sup>th</sup> percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.



View (671 Kb)

Single-criterion visualizations

- Clash list (5.2 Kb): View
- Ramachandran plot kinemage (446 Kb): View in KiNG | Download
- Ramachandran plot PDF (1.7 Mb): View
- Cβ deviation scatter plot (49 Kb): View in KiNG | Download



About MolProbity | Website for the Richardson Lab | Using ecloud x-H | Internal reference 4.1-537

2 of 2