



Analysis output: all-atom contacts and geometry for MCM2_195- 821H.pdb

Summary statistics

All-Atom Contacts	Clashscore, all atoms:	59.16		2 nd percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	2	0.37%	Goal: <0.3%
	Favored rotamers	529	97.60%	Goal: >98%
	Ramachandran outliers	24	3.84%	Goal: <0.05%
	Ramachandran favored	540	86.40%	Goal: >98%
	MolProbity score^	2.88		28 th percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	15	2.54%	Goal: 0
	Bad bonds:	46 / 5011	0.92%	Goal: 0%
	Bad angles:	90 / 6778	1.33%	Goal: <0.1%
Peptide Omegas	Cis Prolines:	1 / 29	3.45%	Expected: ≤1 per chain, or ≤5%
	Cis nonProlines:	21 / 597	3.52%	Goal: <0.05%
	Twisted Peptides:	1 / 626	0.16%	Goal: 0

In the two column results, the left column gives the raw count, right column gives the percentage.

* 100th percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 0.50 points.

Multi-criterion visualizations



[View \(713 Kb\)](#)

Single-criterion visualizations

- **Clash list** (25 Kb): [View](#)
- **Ramachandran plot kinemage** (443 Kb): [View in KiNG](#) | [Download](#)
- **Ramachandran plot PDF** (1.7 Mb): [View](#)
- **C β deviation scatter plot** (43 Kb): [View in KiNG](#) | [Download](#)

[Continue >](#)