



# Analysis output: all-atom contacts and geometry for EPS8\_533-589H.pdb

## Summary statistics

All-Atom Contacts	Clashscore, all atoms:	3.17	97 <sup>th</sup> percentile* (N=1784, all resolutions)	
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	4	7.69%	Goal: <0.3%
	Favored rotamers	48	92.31%	Goal: >98%
	Ramachandran outliers	0	0.00%	Goal: <0.05%
	Ramachandran favored	53	96.36%	Goal: >98%
	MolProbity score <sup>^</sup>	2.02	74 <sup>th</sup> percentile* (N=27675, 0Å - 99Å)	
	Cβ deviations >0.25Å	0	0.00%	Goal: 0
	Bad bonds:	0 / 480	0.00%	Goal: 0%
	Bad angles:	0 / 646	0.00%	Goal: <0.1%
Peptide Omegas	Cis Prolines:	0 / 1	0.00%	Expected: ≤1 per chain, or ≤5%

In the two column results, the left column gives the raw count, right column gives the percentage.

\* 100<sup>th</sup> percentile is the best among structures of comparable resolution; 0<sup>th</sup> percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

<sup>^</sup> MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* fix 1 bad rotamers.

## Multi-criterion visualizations



[View \(63 Kb\)](#)

## Single-criterion visualizations

- **Clash list** (301 bytes): [View](#)
- **Ramachandran plot kinemage** (408 Kb): [View in KiNG](#) | [Download](#)
- **Ramachandran plot PDF** (1.7 Mb): [View](#)
- **C $\beta$  deviation scatter plot** (14 Kb): [View in KiNG](#) | [Download](#)

Continue >