

Analysis output: all-atom contacts and geometry for KCNE1_1-129H.pdb

Summary statistics

	Clashscore, all atoms:	1.94		99 th percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	12	10.43%	Goal: <0.3%
	Favored rotamers	83	72.17%	Goal: >98%
	Ramachandran outliers	2	1.57%	Goal: <0.05%
	Ramachandran favored	119	93.70%	Goal: >98%
	MolProbity score [^]	2.15		67 th percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	0	0.00%	Goal: 0
	Bad bonds:	5 / 1059	0.47%	Goal: 0%
	Bad angles:	1 / 1436	0.07%	Goal: <0.1%
Peptide Omegas	Cis Prolines:	0 / 7	0.00%	Expected: ≤1 per chain, or ≤5%

In the two column results, the left column gives the raw count, right column gives the percentage.

Multi-criterion visualizations



View (138 Kb)

Single-criterion visualizations

- Clash list (343 bytes): View
- Ramachandran plot kinemage (412 Kb): View in KiNG | Download
- Ramachandran plot PDF (1.7 Mb): View
- **C**β **deviation scatter plot** (17 Kb): View in KiNG | Download

^{* 100&}lt;sup>th</sup> percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.



About MolProbity | Website for the Richardson Lab | Using ecloud x-H | Internal reference 4.3