

Analysis output: all-atom contacts and geometry for FOXI1_121-212_FFX1H.pdb

Summary statistics

All-Atom Contacts	Clashscore, all atoms:	0		100 th percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	8	9.88%	Goal: <0.3%
	Favored rotamers	69	85.19%	Goal: >98%
	Ramachandran outliers	2	2.22%	Goal: <0.05%
	Ramachandran favored	70	77.78%	Goal: >98%
	MolProbity score [^]	2.02		75 th percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	0	0.00%	Goal: 0
	Bad bonds:	0 / 773	0.00%	Goal: 0%
	Bad angles:	8 / 1049	0.76%	Goal: <0.1%
Peptide Omegas	Cis Prolines:	0 / 7	0.00%	Expected: ≤1 per chain, or ≤5%
	Twisted Peptides:	1 / 91	1.10%	Goal: 0

In the two column results, the left column gives the raw count, right column gives the percentage.

Multi-criterion visualizations



View (100 Kb)

^{* 100&}lt;sup>th</sup> percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

Single-criterion visualizations

- Clash list (175 bytes): View
- Ramachandran plot kinemage (410 Kb): View in KiNG | Download
- Ramachandran plot PDF (1.7 Mb): View
- **C**β **deviation scatter plot** (16 Kb): View in KiNG | Download

Continue >

About MolProbity | Website for the Richardson Lab | Using ecloud x-H | Internal reference 4.3