



Analysis output: all-atom contacts and geometry for EYA1_322-592H.pdb

Summary statistics

All-Atom Contacts	Clashscore, all atoms:	21.91		28 th percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	5	2.16%	Goal: <0.3%
	Favored rotamers	211	91.34%	Goal: >98%
	Ramachandran outliers	0	0.00%	Goal: <0.05%
	Ramachandran favored	252	93.68%	Goal: >98%
	MolProbity score [^]	2.51		47 th percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	0	0.00%	Goal: 0
	Bad bonds:	15 / 2200	0.68%	Goal: 0%
	Bad angles:	5 / 2984	0.17%	Goal: <0.1%
Peptide Omegas	Cis Prolines:	0 / 7	0.00%	Expected: ≤1 per chain, or ≤5%
	Cis nonProlines:	1 / 263	0.38%	Goal: <0.05%

In the two column results, the left column gives the raw count, right column gives the percentage.

* 100th percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 0.94 points.

Multi-criterion visualizations



Multi-criterion chart

[View](#) (295 Kb)

Single-criterion visualizations

- **Clash list** (4.1 Kb): [View](#)
- **Ramachandran plot kinemage** (420 Kb): [View in KiNG](#) | [Download](#)
- **Ramachandran plot PDF** (1.7 Mb): [View](#)
- **C β deviation scatter plot** (25 Kb): [View in KiNG](#) | [Download](#)

[Continue >](#)