

## Analysis output: all-atom contacts and geometry for MYO15A\_1222-1838.pdb\_7\_1H.pdk

## **Summary statistics**

	Clashscore, all atoms:	1.51		99 <sup>th</sup> percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	0	0.00%	Goal: <1%
	Ramachandran outliers	14	2.28%	Goal: <0.05%
	Ramachandran favored	556	90.41%	Goal: >98%
	MolProbity score <sup>^</sup>	1.43		97 <sup>th</sup> percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	15	2.55%	Goal: 0
	Bad backbone bonds:	0 / 5063	0.00%	Goal: 0%
	Bad backbone angles:	41 / 6849	0.60%	Goal: <0.1%

In the two column results, the left column gives the raw count, right column gives the percentage.

## **Multi-criterion visualizations**



View (584 Kb)

## Single-criterion visualizations

- Clash list (763 bytes): View
- Ramachandran plot kinemage (442 Kb): View in KiNG | Download

1 of 2 6/18/14, 2:08 PM

<sup>\* 100&</sup>lt;sup>th</sup> percentile is the best among structures of comparable resolution; 0<sup>th</sup> percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

<sup>^</sup> MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

- Ramachandran plot PDF (1.7 Mb): View
- Cβ deviation scatter plot (51 Kb): View in KiNG | Download



About MolProbity | Website for the Richardson Lab | Using ecloud x-H | Internal reference 4.1-537

2 of 2