

Analysis output: all-atom contacts and geometry for ESPN_9-336H.pdb

Summary statistics

	Clashscore, all atoms:	67.8		1st percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	3	1.20%	Goal: <1%
	Ramachandran outliers	16	4.91%	Goal: <0.05%
	Ramachandran favored	266	81.60%	Goal: >98%
	MolProbity score [^]	3.08		20 th percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	5	1.68%	Goal: 0
	Bad backbone bonds:	62 / 2518	2.46%	Goal: 0%
	Bad backbone angles:	81 / 3433	2.36%	Goal: <0.1%

In the two column results, the left column gives the raw count, right column gives the percentage.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 1.66 points.

Multi-criterion visualizations



View (337 Kb)

1 of 2 6/13/14, 10:06 AM

^{* 100&}lt;sup>th</sup> percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

Single-criterion visualizations

- Clash list (7.7 Kb): View
- Ramachandran plot kinemage (425 Kb): View in KiNG | Download
- Ramachandran plot PDF (1.7 Mb): View
- Cβ deviation scatter plot (27 Kb): View in KiNG | Download



About MolProbity | Website for the Richardson Lab | Using ecloud x-H | Internal reference 4.1-537

2 of 2