



# Analysis output: all-atom contacts and geometry for OSBPL2\_64- 444\_FFX1H.pdb

## Summary statistics

All-Atom Contacts	Clashscore, all atoms:	7.54		84 <sup>th</sup> percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	7	2.08%	Goal: <0.3%
	Favored rotamers	293	86.94%	Goal: >98%
	Ramachandran outliers	15	3.99%	Goal: <0.05%
	Ramachandran favored	303	80.59%	Goal: >98%
	MolProbity score <sup>^</sup>	2.38		54 <sup>th</sup> percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	10	2.81%	Goal: 0
	Bad bonds:	0 / 3154	0.00%	Goal: 0%
	Bad angles:	86 / 4262	2.02%	Goal: <0.1%
Peptide Omegas	Cis Prolines:	1 / 20	5.00%	Expected: ≤1 per chain, or ≤5%
	Cis nonProlines:	1 / 357	0.28%	Goal: <0.05%
	Twisted Peptides:	8 / 377	2.12%	Goal: 0

In the two column results, the left column gives the raw count, right column gives the percentage.

\* 100<sup>th</sup> percentile is the best among structures of comparable resolution; 0<sup>th</sup> percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

<sup>^</sup> MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 0.17 points.

## Multi-criterion visualizations



## Multi-criterion chart

[View](#) (415 Kb)

## Single-criterion visualizations

- **Clash list** (2.1 Kb): [View](#)
- **Ramachandran plot kinemage** (428 Kb): [View in KiNG](#) | [Download](#)
- **Ramachandran plot PDF** (1.7 Mb): [View](#)
- **C $\beta$  deviation scatter plot** (33 Kb): [View in KiNG](#) | [Download](#)

[Continue >](#)