



# Analysis output: all-atom contacts and geometry for TECTA\_775-1063H.pdb

## Summary statistics

All-Atom Contacts	Clashscore, all atoms:	82.87	0 <sup>th</sup> percentile* (N=1784, all resolutions)	
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	0	0.00%	Goal: <0.3%
	Favored rotamers	250	96.53%	Goal: >98%
	Ramachandran outliers	3	1.05%	Goal: <0.05%
	Ramachandran favored	275	95.82%	Goal: >98%
	MolProbity score <sup>^</sup>	2.67		38 <sup>th</sup> percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	10	3.68%	Goal: 0
Peptide Omegas	Cis Prolines:	1 / 7	14.29%	Expected: ≤1 per chain, or ≤5%

In the two column results, the left column gives the raw count, right column gives the percentage.

\* 100<sup>th</sup> percentile is the best among structures of comparable resolution; 0<sup>th</sup> percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

<sup>^</sup> MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 0.90 points.

## Multi-criterion visualizations



[View](#) (282 Kb)

## Single-criterion visualizations

- **Clash list** (4.6 Kb): [View](#)
- **Ramachandran plot kinemage** (422 Kb): [View in KiNG](#) | [Download](#)
- **Ramachandran plot PDF** (1.7 Mb): [View](#)

- **C $\beta$  deviation scatter plot** (26 Kb): [View in KiNG](#) | [Download](#)

Continue >

About [MolProbity](#) | Website for [the Richardson Lab](#) | Using ecloud x-H | Internal reference 4.2