



Analysis output: all-atom contacts and geometry for COL11A1_1596-1805H.pdb

Summary statistics

All-Atom Contacts	Clashscore, all atoms:	61.31		2 nd percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	1	0.54%	Goal: <0.3%
	Favored rotamers	175	94.09%	Goal: >98%
	Ramachandran outliers	2	0.96%	Goal: <0.05%
	Ramachandran favored	194	93.27%	Goal: >98%
	MolProbity score^	2.70		36 th percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	2	1.04%	Goal: 0
	Bad bonds:	10 / 1714	0.58%	Goal: 0%
	Bad angles:	28 / 2320	1.21%	Goal: <0.1%
Peptide Omegas	Cis Prolines:	1 / 11	9.09%	Expected: ≤1 per chain, or ≤5%
	Cis nonProlines:	3 / 198	1.52%	Goal: <0.05%
	Twisted Peptides:	3 / 209	1.44%	Goal: 0

In the two column results, the left column gives the raw count, right column gives the percentage.

* 100th percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 3.08 points.

Multi-criterion visualizations



[View \(241 Kb\)](#)

Single-criterion visualizations

- **Clash list** (4.6 Kb): [View](#)
- **Ramachandran plot kinemage** (417 Kb): [View in KiNG](#) | [Download](#)
- **Ramachandran plot PDF** (1.7 Mb): [View](#)
- **C β deviation scatter plot** (23 Kb): [View in KiNG](#) | [Download](#)

Continue >