

Analysis output: all-atom contacts and geometry for MYO6_771-913H.pdb

Summary statistics

	Clashscore, all atoms:	41.44		7 th percentile* (N=1784, all resolutions)
	Clashscore is the number of serious steric overlaps (> 0.4 Å) per 1000 atoms.			
Protein Geometry	Poor rotamers	2	1.49%	Goal: <0.3%
	Favored rotamers	128	95.52%	Goal: >98%
	Ramachandran outliers	2	1.42%	Goal: <0.05%
	Ramachandran favored	138	97.87%	Goal: >98%
	MolProbity score [^]	2.26		61 st percentile* (N=27675, 0Å - 99Å)
	Cβ deviations >0.25Å	3	2.14%	Goal: 0
	Bad bonds:	10 / 1188	0.84%	Goal: 0%
	Bad angles:	17 / 1587	1.07%	Goal: <0.1%
Peptide Omegas	Cis Prolines:	0/3	0.00%	Expected: ≤1 per chain, or ≤5%
	Twisted Peptides:	1 / 142	0.70%	Goal: 0

In the two column results, the left column gives the raw count, right column gives the percentage.

By adding H to this model and allowing Asn/Gln/His flips, we could *automatically* improve your clashscore by 0.41 points.

Multi-criterion visualizations



View (163 Kb)

Single-criterion visualizations

^{* 100&}lt;sup>th</sup> percentile is the best among structures of comparable resolution; 0th percentile is the worst. For clashscore the comparative set of structures was selected in 2004, for MolProbity score in 2006.

[^] MolProbity score combines the clashscore, rotamer, and Ramachandran evaluations into a single score, normalized to be on the same scale as X-ray resolution.

- Clash list (4.4 Kb): View
- Ramachandran plot kinemage (413 Kb): View in KiNG | Download
 Ramachandran plot PDF (1.7 Mb): View
- **C**β deviation scatter plot (18 Kb): View in KiNG | Download

Continue >

About MolProbity | Website for the Richardson Lab | Using ecloud x-H | Internal reference 4.3