

100A

Xiaowu Dai

Basics

Population

Region

Coin

Markov

Reasoning

STATS 100A: BASICS & EXAMPLES

Xiaowu Dai

Department of Statistics and Data Science
University of California, Los Angeles

Some pictures are taken from the internet.
Credits belong to original authors.





Sample space

100A

Xiaowu Dai

Basics

Population

Region

Coin


Markov

Reasoning

Experiment \rightarrow **outcome** \rightarrow **number**

Example 1: Roll a die

Sample Point



Sample space Ω : The set of all the outcomes (or sample points, elements).

Randomly sample an outcome from the sample space.





Event

100A

Xiaowu Dai

Basics

Population

Region


Coin

Markov

Reasoning

Experiment \rightarrow **outcome** \rightarrow **number**

Example 1: Roll a die



Sample space Ω : The set of all the outcomes.

Event A :

- (1) A **statement** about the outcome, e.g., bigger than 4.
- (2) A **subset** of sample space, e.g., $\{5, 6\}$.





Counting equally likely possibilities

100A

Xiaowu Dai

Basics

Population

Region


Coin

Markov

Reasoning

Experiment \rightarrow **outcome** \rightarrow **number**

Example 1: Roll a die



Assume the die is fair so that all the outcomes are **equally likely**.

Probability: defined on event:

$$P(A) = \frac{|A|}{|\Omega|} = \frac{2}{6} = \frac{1}{3}.$$

$|A|$ counts the size of A , i.e., the number of elements in A .





Random variable

100A

Xiaowu Dai

Basics

Population

Region


Coin

Markov

Reasoning

Experiment \rightarrow **outcome** \rightarrow **number**

Example 1: Roll a die



Random variable: Let X be the number:

$$P(X > 4) = \frac{1}{3}.$$

An event is a **math statement** about the random variable.

We can either use events or use random variables.

In Parts 2 and 3, we will focus on random variables.





Conditional probability

100A

Xiaowu Dai

Basics

Population

Region


Coin

Markov

Reasoning

Experiment \rightarrow **outcome** \rightarrow **number**

Example 1: Roll a die



Conditional probability: Let B be the event that the number is 6. Given that A happens, what is the probability of B ?

$$P(B|A) = \frac{1}{2}.$$

As if we randomly sample a number from A .

As if A is the sample space.





Conditional probability

100A

Xiaowu Dai

Basics

Population

Region


Coin

Markov

Reasoning

Experiment \rightarrow **outcome** \rightarrow **number**

Example 1: Roll a die



Random variable

$$P(X = 6 | X > 4) = \frac{1}{2}.$$





Relations

100A

Xiaowu Dai

Basics

Population


Region

Coin

Markov

Reasoning

Example 1: Roll a die



Complement

Statement: Not A

Subset: $A^c = \{1, 2, 3, 4\}$.





Relations

100A

Xiaowu Dai

Basics

Population

Region

Coin

Markov


Reasoning

Example 1: Roll a die

$$A = \{1,2,3\}$$

$$B = \{3,4,5\}$$

$$A \cup B = \{1,2,3,4,5\}$$



Venn diagram

Union

Statement: A or B .

Subset: $A \cup B$.





Relations

100A

Xiaowu Dai

Basics

Population

Region

Coin

Markov


Reasoning

Example 1: Roll a die

$$A = \{1, 2, 3, 4\}$$

$$B = \{3, 4, 5, 6\}$$

$$A \cap B = \{3, 4\}$$



Intersection

Statement: A and B .

Subset: $A \cap B$.





Sample space is population

100A

Xiaowu Dai

Basics

Population

Region

Coin

Markov

Reasoning

Experiment \rightarrow outcome \rightarrow number

Example 2: Sample a random person from a population of 100 people, 50 males and 50 females. 30 males are taller than 6 ft, 10 females are taller than 6 ft.

The sample space Ω is the population.

	male	female
taller than 6 ft	30	10
shorter than 6 ft		
	50	50





Events as sub-populations

100A

Xiaowu Dai

Basics

Population

Region

Coin

Markov

Reasoning

Experiment \rightarrow **outcome** \rightarrow **number**

Example 2: Let A be the event that the person is male. Let B be the event that the person is taller than 6 feet (or simply the person is tall). A is the sub-population of males, and B is the sup-population of tall people.

	male	female
taller than 6 ft	30	10
shorter than 6 ft		
	50	50





Probability is population proportion

100A

Xiaowu Dai

Basics

Population

Region

Coin

Markov

Reasoning

Experiment \rightarrow **outcome** \rightarrow **number**

Example 2: A male, B tall.

	male	female
taller than 6 ft	30	10
shorter than 6 ft		
	50	50

$$P(A) = \frac{|A|}{|\Omega|} = \frac{50}{100} = 50\%.$$

$$P(B) = \frac{|B|}{|\Omega|} = \frac{30 + 10}{100} = 40\%.$$

Probability = population proportion.





Conditional probability is proportion of sub-population

100A

Xiaowu Dai

Basics

Population

Region

Coin

Markov

Reasoning

Experiment → **outcome** → **number**

Example 2: A male, B tall.

	male	female
taller than 6 ft	30	10
shorter than 6 ft		
	50	50

$$P(A|B) = \frac{|A \cap B|}{|B|} = \frac{30}{40} = 75\%.$$

Among tall people, what is the proportion of males?

$$P(B|A) = \frac{|A \cap B|}{|A|} = \frac{30}{50} = 60\%.$$

Among males, what is the proportion of tall people?

Conditional probability = proportion within sub-population.





Random variable as a function of outcome

100A

Xiaowu Dai

Basics

Population

Region

Coin

Markov

Reasoning

Link between event and random variable.

Example 2: A male, B tall.

Let $\omega \in \Omega$ be a person. Let $X(\omega)$ be the gender of ω , so that $X(\omega) = 1$ if ω is male, and $X(\omega) = 0$ if ω is female. Let $Y(\omega)$ be the height of ω . Then

$$A = \{\omega : X(\omega) = 1\}, \quad B = \{\omega : Y(\omega) > 6\}.$$

$$P(A) = P(\{\omega : X(\omega) = 1\}) = P(X = 1).$$

$$P(B) = P(\{\omega : Y(\omega) > 6\}) = P(Y > 6).$$

$$P(B|A) = P(Y > 6|X = 1), \quad P(A|B) = P(X = 1|Y > 6).$$





Axiom 0

100A

Xiaowu Dai

Basics

Population

Region


Coin

Markov

Reasoning

Equally likely scenario

A real population of people, under purely random sampling
or imagined population of equally likely possibilities



$$P(A) = \frac{|A|}{|\Omega|}.$$

Axiom 0.

Can always translate a problem into equally likely setting.





Conditional probability

100A

Xiaowu Dai

Basics

Population


Region

Coin

Markov

Reasoning

Equally likely scenario



$$P(A|B) = \frac{|A \cap B|}{|B|} = \frac{|A \cap B|/|\Omega|}{|B|/|\Omega|} = \frac{P(A \cap B)}{P(B)}.$$

Physical: sample from B . B defines condition.

Mental: know that B happened, as if sample from B .

Axiom 4 or definition of conditional probability.