

## Week 1

### Rings and examples of rings

Consider a field  $F$  and a vector space  $V$  over  $F$ . Then  $\mathcal{L}(V)$  itself is not a vector space: If  $S, T \in \mathcal{L}(V)$  then it is not always the case that  $T \circ S = S \circ T$ , and it is not the case that for every  $T \in \mathcal{L}(V)$  there is a  $T^{-1}$  such that  $T \circ T^{-1} = I$ . Then  $\mathcal{L}(V)$  is a ring:

**Definition 1** (Ring). A ring  $(R, +, \cdot)$  is a set  $R$  with two operations, denoted  $+$  and  $\cdot$  such that for all  $a, b, c$  in  $R$ :

- (i)  $(R, +)$  is an abelian group.
- (ii)  $(ab)c = a(bc)$ .
- (iii)  $a(b + c) = ab + ac$  and  $(b + c)a = ba + ca$ .

If  $ab = ba$  for all  $a, b \in R$  then  $R$  is a commutative ring. If  $1_R a = a = a 1_R$  for all  $a \in R$  then  $R$  is a ring with identity.

**Example.**  $(\mathbb{Z}, +, \cdot)$  is a commutative ring with identity.

**Example.**  $(\mathcal{L}(V), +, \circ)$  is a ring with identity

A nonzero element  $a \in R$  is a left (right) zero divisor if there is a nonzero element  $b \in R$  such that  $ab = 0$  ( $ba = 0$ ). A zero divisor is both a left and right zero divisor.

An element  $a$  in a ring  $R$  with identity is said to be left (right) invertible if there exists a  $c \in R$  such that  $ca = 1_R$  ( $ac = 1_R$ ). The element  $c$  a left (right) inverse of  $a$ . An element  $a \in R$  that is both left and right invertible is invertible or a unit.

The set of units in a ring  $R$  with identity forms a group under multiplication.

There are rings with identity and inverses that do not commute:

**Example.**  $\mathbb{H}$ , the quaternions, is a non-commutative ring with identity and inverses, also known as division ring or skew field.

Consider  $\mathcal{F}(X, R)$ , where  $X \neq \emptyset$  and  $R$  is a ring. This is the set of functions from  $X$  to  $R$ . Then for  $f, g$  we can define addition as  $(f + g)(x) = f(x) + g(x)$  and multiplication as  $(fg)(x) = f(x)g(x)$ . Then  $\mathcal{F}(X, R)$  is a ring.

**Definition 2** (Integral domain). A commutative ring  $R$  with identity  $1_R \neq 0$  and no zero divisors is called an integral domain. A ring  $D$  with identity  $1_D \neq 0$  in which every nonzero element is a unit is called a division ring. A field is a commutative division ring.

### Ring homomorphisms

Let  $R, S$  be rings and  $\phi : R \rightarrow S$  a function such that for all  $a, b \in R$ ,  $\phi(a + b) = \phi(a) + \phi(b)$  and  $\phi(ab) = \phi(a)\phi(b)$ .