

An Improved Bound for Equitable Proper Labellings


Julien Bensmail^a, Clara Marcille^b

a: I3S/INRIA, Université Côte d'Azur, France

b: LaBRI, Université de Bordeaux, France

IWOCA, July 2, 2024

Magic labellings




Magic labellings

2	7	6
9	5	1
4	3	8


Magic labellings

2	7	6
9	5	1
4	3	8




Magic labellings

2	7	6
9	5	1
4	3	8




Magic labellings

2	7	6
9	5	1
4	3	8




Magic labellings

2	7	6
9	5	1
4	3	8





Magic labellings

2	7	6
9	5	1
4	3	8




Antimagic labellings




Antimagic labellings

9	7	6
2	5	1
8	4	3




Antimagic labellings

9	7	6
2	5	1
8	4	3



Antimagic labellings

9	7	6
2	5	1
8	4	3




Definition

A *labelling* is said to be (locally) *distinguishing* if any pair of (adjacent) vertices have different *resulting sum*.

Antimagic labellings

9	7	6
2	5	1
8	4	3



Definition

A *labelling* is said to be (locally) *distinguishing* if any pair of (adjacent) vertices have different *resulting sum*.

We will get back to all this terminology later.

Labellings

Definition

A k -labelling of a graph G is a function $\ell : E(G) \rightarrow \{1, \dots, k\}$.

Labellings

Definition

A k -labelling of a graph G is a function $\ell : E(G) \rightarrow \{1, \dots, k\}$.

Definition

We call *resulting sum* (relative to a labelling ℓ) of a vertex u the sum of labels on edges incident to u . We denote it $\sigma_\ell(u)$.

Labellings

Definition

A k -labelling of a graph G is a function $\ell : E(G) \rightarrow \{1, \dots, k\}$.

Definition

We call *resulting sum* (relative to a labelling ℓ) of a vertex u the sum of labels on edges incident to u . We denote it $\sigma_\ell(u)$.

Definition (related to Irregularity Strength)

We say a labelling ℓ is *distinguishing* if for every two vertices u and v of G , $\sigma_\ell(u) \neq \sigma_\ell(v)$.

Labellings

Definition

A k -labelling of a graph G is a function $\ell : E(G) \rightarrow \{1, \dots, k\}$.

Definition

We call *resulting sum* (relative to a labelling ℓ) of a vertex u the sum of labels on edges incident to u . We denote it $\sigma_\ell(u)$.

Definition (Local)

We say a labelling ℓ is *distinguishing* if for every two adjacent vertices u and v of G , $\sigma_\ell(u) \neq \sigma_\ell(v)$.

Equitable Labellings

Definition

A k -labelling is said to be *equitable* if for all $1 \leq i, j \leq k$, the number of edges with label i and the number of edges with label j differ by at most 1.

Equitable Labellings

Definition

A k -labelling is said to be *equitable* if for all $1 \leq i, j \leq k$, the number of edges with label i and the number of edges with label j differ by at most 1.

In particular, a locally Antimagic labelling can be defined as a special case of equitable $|E(G)|$ -labelling!

Equitable Labellings

Definition

A k -labelling is said to be *equitable* if for all $1 \leq i, j \leq k$, the number of edges with label i and the number of edges with label j differ by at most 1.

In particular, a locally Antimagic labelling can be defined as a special case of equitable $|E(G)|$ -labelling!

Definition

We denote $\overline{\chi}_{\Sigma}(G)$ the smallest integer k such that G admits an equitable k -labelling.

Equitable Labellings

Definition

A k -labelling is said to be *equitable* if for all $1 \leq i, j \leq k$, the number of edges with label i and the number of edges with label j differ by at most 1.

In particular, a locally Antimagic labelling can be defined as a special case of equitable $|E(G)|$ -labelling!

Definition


We denote $\overline{\chi}_{\Sigma}(G)$ the smallest integer k such that G admits an equitable k -labelling.

Theorem (consequence of Lyngsie and Zhong, 2018)

If G is a "nice" graph, then $\overline{\chi}_{\Sigma}(G) \leq |E(G)|$.


Examples

An equitable labelling which happens to be antimagic:




Examples

An equitable labelling which happens to be antimagic:



An equitable 3-labelling:



Conjecture and contribution

Theorem (consequence of Lyngsie and Zhong, 2018)

If G is a nice graph, then $\overline{\chi}_{\Sigma}(G) \leq |E(G)|$.

Conjecture and contribution

Theorem (consequence of Lyngsie and Zhong, 2018)

If G is a nice graph, then $\overline{\chi_{\Sigma}}(G) \leq |E(G)|$.

Conjecture (Bensmail, Fioravantes, Mc Inerney, Nisse, 2021)

If G is a nice graph different from K_4 , then $\overline{\chi_{\Sigma}}(G) \leq 3$.

Conjecture and contribution

Theorem (consequence of Lyngsie and Zhong, 2018)

If G is a nice graph, then $\overline{\chi_{\Sigma}}(G) \leq |E(G)|$.

Conjecture (Bensmail, Fioravantes, Mc Inerney, Nisse, 2021)

If G is a nice graph different from K_4 , then $\overline{\chi_{\Sigma}}(G) \leq 3$.

Theorem (Bensmail, M., 2024+)


If G is a nice graph, then $\overline{\chi_{\Sigma}}(G) \leq \left\lfloor \frac{|E(G)|}{2} \right\rfloor + 2$.

Preliminary work

We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \left\lfloor \frac{|E(G)|}{2} \right\rfloor$).


Preliminary work

We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \left\lfloor \frac{|E(G)|}{2} \right\rfloor$).



Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \left\lfloor \frac{|E(G)|}{2} \right\rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 6, 6)$.

Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \left\lfloor \frac{|E(G)|}{2} \right\rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 6, 6)$.

Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \left\lfloor \frac{|E(G)|}{2} \right\rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 6, 6)$.

Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \left\lfloor \frac{|E(G)|}{2} \right\rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 6, 6)$.

Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \left\lfloor \frac{|E(G)|}{2} \right\rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 5, 5, 6, 6)$.

Preliminary work

We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \left\lfloor \frac{|E(G)|}{2} \right\rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 5, 5, 6, 6)$.

Question

How do we carry the fact that some edges received a label?

Weight function


Solution

We will keep the information by updating a *weight function*.

Weight function

Solution


We will keep the information by updating a *weight function*.



Weight function

Solution


We will keep the information by updating a *weight function*.



Weight function

Solution

We will keep the information by updating a *weight function*.



We now consider the weighted graph (G, c) where c is the weight function.

Ideas of the proof

- Proceed by induction on the number of vertices.

Ideas of the proof

- Proceed by induction on the number of vertices.
- Build a partial labelling of the graph, and extend it.

Ideas of the proof

- Proceed by induction on the number of vertices.
- Build a partial labelling of the graph, and extend it.

Problem

Once all the edges incident to u have been labelled, there is no way to change $\sigma_\ell(u)$.

Ideas of the proof

- Proceed by induction on the number of vertices.
- Build a partial labelling of the graph, and extend it.
- Ensure that the vertex at hand will have a resulting sum smaller than all vertices treated later in the induction.

Ideas of the proof

- Proceed by induction on the number of vertices.
- Build a partial labelling of the graph, and extend it.
- Ensure that the vertex at hand will have a resulting sum smaller than all vertices treated later in the induction.

Problem


Maybe $G - u$ has a component isomorphic to K_2 .

Ideas of the proof

- Proceed by induction on the number of vertices.
- Build a partial labelling of the graph, and extend it.
 - Ensure that a vertex will have a resulting sum smaller than the vertex treated later in the induction.
- Handle exceptions on the way.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



Here, the minimum possible resulting sum for this vertex is $1 + 6 + 7 = 14$. For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



Here, the minimum possible resulting sum for this vertex is $3 + 6 = 9$.
For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.


Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



Handling exceptions


Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



We choose a vertex of highest degree amongst vertices of lowest potential.

Handling exceptions


Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



We choose a vertex of highest degree amongst vertices of lowest potential.

Handling exceptions


Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



How to assign each label?

Handling exceptions


Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



How to assign each label?

Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Problem

A component isomorphic to K_2 with constant weight cannot be labelled.


Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Handling exceptions

Consider this weighted graph with $L = (2, 3, 3, 4, 4)$:



Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Problem

Two adjacent vertices of lowest potential can be in conflict.

Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Problem

Two adjacent vertices of lowest potential can be in conflict.


Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



Handling exceptions

Consider this weighted graph with $L = (2, 3, 3, 4, 4)$:



Conclusion

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.
- The question of a constant bound remains completely open.

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.
- The question of a constant bound remains completely open.
- The tools we used do not generalize nicely to multiplicity higher than 2.

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.
- The question of a constant bound remains completely open.
- The tools we used do not generalize nicely to multiplicity higher than 2.
- More work can be done to characterize the graphs that cannot be done with a sequence of $|E(G) + 1|$ labels and multiplicity 2.

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.
- The question of a constant bound remains completely open.
- The tools we used do not generalize nicely to multiplicity higher than 2.
- More work can be done to characterize the graphs that cannot be done with a sequence of $|E(G) + 1|$ labels and multiplicity 2.


Thank you for your attention!

The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$


The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$




The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$




The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$




The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$




The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$




The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$




The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$



The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$



The extra labels

They are some configurations where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$

