


# Combined PET and MRI Radiomics with Breast Cancer Outcomes

Shih-ying Huang, Benjamin L. Franc, Roy Harnish, Timothy Copeland, Vignesh Arasu  
Ella F. Jones, Nola M. Hylton, and Youngho Seo, *Senior Member, IEEE*

## Why Radiomics?

- Improve cancer treatment with more precise and personalized disease management
- Radiomics from 3D imaging modalities has been shown useful for discovering imaging biomarkers<sup>1</sup>




## Radiomics Workflow

### 1 Data Collection

- PET and dynamic-contrast-enhanced MR (DCE-MRI) images from patients diagnosed with invasive breast cancer in 2003 - 2005
- All images were acquired *prior* to any surgery/treatment
- Gathered clinical data from the medical records and cancer registry including:
  - tumor histology, grade, T/N/overall stages
  - estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (HER2) status
  - recurrence site, status, and duration until recurrence/death since diagnosis

### 2 Tumor Segmentation

- Manually segmented tumors from the SUV-converted PET images using MeVisLab<sup>®</sup>
- Segmented tumors from the DCE-MR images by the signal-enhancement ratio (SER) map





### 3 Radiomic Feature Extraction

- 8 First-order statistics
- 9 Shape and size features
- 25 3D Gray level co-occurrence matrix (GLCM) texture features
- Images were re-sampled to isotropic voxel size
- GLCM matrices were created with fixed bin width
- Computed with in-house Python-TK<sup>2</sup> software and validated with pyradiomics open-source software<sup>3</sup>


### 4 Clustering and Statistical Analysis

- Performed unsupervised clustering using consensus clustering implemented in R<sup>4</sup>
- $\chi^2$  test for significant association between the tumor clusters and tumor profiling variables
- Computed correlation coefficients for any inference between each radiomic feature to the outcome
- Ordered variables: Spearman's rank correlation coefficients
- Un-ordered variables: multiple-regression correlation coefficients

## What Did We Find?



## Patient Demographics



## Conclusion

- Extracted and validated PET and MR image radiomics
- Unsupervised clustering show radiomic features have potential for predicting tumor profiling
- Texture features of PET and MR images, in addition to first-order image statistics, could be key imaging biomarkers in building a predictive model for breast cancer diagnosis and prognosis

## Thank You

This research is supported by Department of Defense Grant W81XWH-17-1-0033, Precision Imaging of Cancer and Therapy Program (PICT) in Departments of Radiation Oncology, and Radiology and Biomedical Imaging, UCSF, and National Cancer Institute Grant R01 CA154561

### References

1. Lambin, P., Rios-Velazquez, E., Leijenaar, R., Carvalho, S., Van Stiphout, R. G. P. M., Granton, P., Zegers, C. M. L., Gillies, R., Boellard, R., Dekker, A., Aerts, H. J. W. L., "Radiomics: Extracting more information from medical images using advanced feature analysis", *European Journal of Cancer* 2012
2. "The Insight Segmentation and Registration Toolkit" ([www.itk.org](http://www.itk.org))
3. Joost JM van Griethuysen, Andriy Fedorov, Chintan Parmar, Ahmed Hosny, Nicole Aucoin, Vivek Narayan, Regina GH Beets-Tan, Jean-Christophe Fillon-Robin, Steve Pieper, Hugo JW Aerts, "Computational Radiomics System to Decode the Radiographic Phenotype"; *Accepted Cancer Research* 2017
4. Wilkerson, M.D., Hayes, D.N. (2010). ConsensusClusterPlus: a class discovery tool with confidence assessments and item tracking. *Bioinformatics*, 2010 Jun 15;26(12):1572-3.