

Stratified homotopy theory

Clark Barwick

Spring 2020

Contents

o	Introduction	1
	Notes on sobriety	2
1	Stratified topology	6
	References	6

o Introduction

Our work here stems from a general principle: if X is a (nice) geometric object (a topological space, a variety, *etc.*), then a suitable class A of sheaves on X determines and is determined by a certain ‘homotopy type’ X_A attached to X . The larger the class A , the finer we can expect the homotopy type X_A to be.

In this introduction, our aim is to examine this principle in a four relatively familiar situations:

- when X is a topological space, and A is the family of all sheaves;
- when X is a topological space, and A is the family of constant sheaves;
- when X is a topological space, and A is the family of locally constant sheaves;
- when X is a scheme, and A is the family of locally constant étale sheaves.

In these circumstances, the object X_A is a homotopy type (or piece thereof) in more or less the usual sense.

In the body of this text, we will study two further situations:

- when X is a stratified topological space, and A is the family of constructible sheaves;
- when X is a scheme, and A is the family of constructible sheaves.

In these situations, the object X_A will be a *stratified homotopy type*. We will see that this is a dramatically different object, but it comes with much finer information about the structure of X .

Notes on sobriety

The most elementary illustration of the principle above is the consideration of topological spaces X that can be recovered from their entire categories of sheaves. These are called the *sober* topological spaces.

o.1 Definition. Let X be a topological space. A closed subset $Z \subseteq X$ is *irreducible* if and only if it is nonempty and if, for any closed subsets $Z_1, Z_2 \subseteq X$ such that $Z = Z_1 \cup Z_2$, either $Z = Z_1$ or $Z = Z_2$. Dually, an open subset $U \subseteq X$ is *irreducibly open* if and only if it is proper and if, for any open subsets $U_1, U_2 \subseteq X$ such that $U_1 \cap U_2 \subseteq U$, either $U_1 \subseteq U$ or $U_2 \subseteq U$.

o.2 Example. If X is a topological space with a point $x \in X$, then the closure \bar{x} is irreducible. In this case, one says that x is a *generic point* of X . The union of all the open subsets that do not contain x is irreducibly open.

o.3 Definition. Let X be a topological space, and let $I(X)$ be the set of irreducible closed subsets of X . Then we say that X is *sober* if the map $X \rightarrow I(X)$ given by $x \mapsto \bar{x}$ is a bijection; that is, X is sober if and only if every irreducible closed subset of X has a unique generic point.

o.4. Sober topological spaces are always Kolmogoroff (*i.e.*, any distinct points are topologically distinguishable). Indeed, Kolmogoroff topological spaces are exactly those with the property that the map $x \mapsto \bar{x}$ is an injection.

o.5 Example. Any Hausdorff space is sober.

o.6 Example. The Zariski topology of any scheme is sober. In particular, not all sober topological spaces are T_1 .

o.7 Example. The set N with the cofinite topology is T_1 but not sober.

o.8 Construction. Let X be a topological space, and let $I(X)$ be the set of irreducible closed subsets thereof. Write γ for the map $x \mapsto \bar{x}$. We topologise $I(X)$ with the finest topology such that γ_X is continuous; in other words, a subset $U \subseteq I(X)$ is open if and only if $\gamma_X^{-1}U$ is open in X . Equivalently, U is open if and only if there exists an open subset $V \subseteq X$ such that $U = \{F \in I(X) : F \cap V \neq \emptyset\}$.

Thus γ induces a bijection between the opens of $I(X)$ and the opens of X , and $I(X)$ is sober.

The assignment $X \mapsto I(X)$ along with the natural transformation γ together define a left adjoint to the forgetful functor from sober topological spaces to all topological spaces. Thus one may *sober up* any topological space X .

Now let us demonstrate that a sober space is completely controlled by its lattice of open sets.

o.9 Notation. Let X be a topological space. Then we write $\Omega(X)$ for the poset of open subsets of X ; this is a complete lattice. If Y is also a topological space, then we write $\text{Map}^{cl}(\Omega(Y), \Omega(X))$ for the set of *complete lattice homomorphisms* – monotonic maps $\Omega(Y) \rightarrow \Omega(X)$ that preserve finite intersections and arbitrary unions.

0.10 Proposition. *Let X and Y be sober topological spaces; Then the assignment $f \mapsto f^{-1}$ is a bijection*

$$\text{Map}(X, Y) \simeq \text{Map}^{cl}(\Omega(Y), \Omega(X)) .$$

Proof. Let $F: \Omega(Y) \rightarrow \Omega(X)$ be a monotonic map that preserves finite intersections and arbitrary unions. Now let us define a map $f: X \rightarrow Y$: let $x \in X$. For each of X and Y , the points are in bijection with the irreducible closed subsets, which are in turn in bijection with the irreducibly open subsets. Now the map F induces a map from the irreducibly open subsets of X to those of Y : this map carries an irreducible open $U \subseteq X$ to the largest open $V \subseteq Y$ such that $F(V) \subseteq U$. Following these bijections, we obtain a unique map f with the property that $f^{-1} = F$.

To be explicit, let $V \in \Omega(Y)$ be the largest open subset such that $x \notin F(V)$. This subset is irreducibly open, and so its complement is an irreducible closed. The value $f(x)$ is the unique generic point of $Y \setminus V$. \square

This provides a fully faithful embedding of sober topological spaces into complete lattices. We now show that this provides a similar fully faithful embedding of sober topological spaces into topoi.

0.11 Notation. Let X be a topological space. We shall write $\tilde{X}_{\leq 0}$ for the topos of sheaves of sets on X . (This notation may seem a little strange, but soon we will have to contend with sheaves of *spaces* on X , and we are reserving this notation for that notion.)

For any two topoi X and Y , we write $\text{Fun}_*(X, Y)$ for the category whose objects are geometric morphisms $f_*: X \rightarrow Y$ and whose morphisms are natural transformations $f_* \rightarrow g_*$.

0.12. If X is a topological space, then we can recover $\Omega(X)$ from $\tilde{X}_{\leq 0}$ as the poset of subobjects of the terminal object $1_{\tilde{X}_{\leq 0}}$. Now we can make the following little observation: if X and Y are topological spaces, then the left adjoint of a geometric morphism $f_*: \tilde{X}_{\leq 0} \rightarrow \tilde{Y}_{\leq 0}$ restricts to a complete lattice homomorphism $\Omega(X) \rightarrow \Omega(Y)$. This is an inverse to the obvious functor $\text{Map}^{cl}(\Omega(Y), \Omega(X)) \rightarrow \text{Fun}_*(\tilde{X}_{\leq 0}, \tilde{Y}_{\leq 0})$.

In particular, please observe that the geometric morphisms $\tilde{X}_{\leq 0} \rightarrow \tilde{Y}_{\leq 0}$ form a discrete category.

0.13 Corollary. *Let X and Y be sober topological spaces. Then the assignment $f \mapsto f_*$ is an equivalence of categories*

$$\text{Map}(X, Y) \simeq \text{Fun}_*(\tilde{X}_{\leq 0}, \tilde{Y}_{\leq 0}) .$$

Connectedness and constant sheaves

Let us begin by understanding the nature of connectedness.

0.14 Notation. We well begin by contemplating sheaves of *sets* on topological spaces. For any topological space X , let $\tilde{X}_{\leq 0}$ be the category of sheaves of sets on X . (Eventually, we shall have to consider sheaves of *spaces* on X , and we are saving the notation \tilde{X} for that.)

0.15 Definition. The *constant sheaf* at a set S on a topological space X is the sheafification of the constant presheaf $U \mapsto S$.

0.16. The formation of the constant sheaf defines a left exact left adjoint

$$\Gamma_X^* : \mathbf{Set} \rightarrow \widetilde{X}_{\leq 0}.$$

Its right adjoint $\Gamma_{X,*}$ is the formation of global sections $F \mapsto F(X)$.

0.17. For any topological space X , any set S , and any point $x \in X$, the stalk of the constant sheaf $\Gamma_X^*(S)$ at X is canonically isomorphic to S .

Indeed, $x^* \Gamma_X^*$ is a left exact left adjoint from \mathbf{Set} to itself; such a functor is isomorphic in a unique fashion to the identity.

0.18 Example. Constant sheaves are not constant as presheaves. Indeed, let X be the discrete space $\{0, 1\}$, and let S be any set. Then the constant sheaf at S on X certainly has the property that its values on $\{0\}$ and $\{1\}$ are each the set S , but now the sheaf condition requires that the global sections are given by

$$\Gamma_{X,*} \Gamma_X^*(S) = \Gamma_X^*(S)\{0, 1\} \cong \Gamma_X^*(S)\{0\} \times \Gamma_X^*(S)\{1\} \cong S \times S.$$

If S has at least two elements, then it follows that $\Gamma_X^*(S)$ is not constant as a presheaf.

The previous example does illustrate a general principle:

0.19 Lemma. Let X be a discrete topological space, and let S be a set. Then $\Gamma_{X,*} \Gamma_X^*(S) \cong \text{Map}(X, S)$.

If we want to understand various constant sheaves, we can do so by coming to grips with the various functor $\Gamma_{X,*} \Gamma_X^* : \mathbf{Set} \rightarrow \mathbf{Set}$ as X varies. The first thing we can notice is that this functor is a left exact accessible functor.

0.20 Definition. A *proobject* of a category C with all finite limits is a left exact accessible functor $C \rightarrow \mathbf{Set}$. The category $\text{Pro}(C)$ of proobjects is the *opposite* of the full subcategory of $\text{Fun}(C, \mathbf{Set})$ spanned by the left exact accessible functors.

0.21 Example. If C is a category with all finite limits, then the Yoneda embedding provides a fully faithful functor $\mathcal{Y} : C \hookrightarrow \text{Pro}(C)$. Explicitly, if X is an object of a category C , then the proobject it defines is $\text{Map}(X, -)$.

0.22 Example. As a matter of terminology, we call the opposite A^{op} of a filtered category an *inverse category*; we call a diagram indexed by an inverse category an *inverse system*; and we call a limit of an inverse system an *inverse limit*.

If $X : A^{op} \rightarrow C$ is an inverse system in a category C with all finite limits, then the limit $\lim_{\alpha \in A^{op}} X_\alpha$ formed in $\text{Fun}(C, \text{Set})^{op}$ is a proobject; this is the proobject $Y \mapsto \text{colim}_{\alpha \in A} \text{Map}(X_\alpha, Y)$. Furthermore, *every* proobject of C can be formed in such a manner.

Now if X and Y are two proobjects that are exhibited as limits of inverse systems in this sense, then one has

$$\text{Map}_{\text{Pro}(C)}(X, Y) \cong \lim_{\beta \in B^{op}} \text{colim}_{\alpha \in A} \text{Map}_C(X, Y).$$

0.23 Example. Let X be a topological space. We obtain a proset $\pi_0^{top}(X) := \Gamma_{X,*} \Gamma_X^*$. This defines a functor from topological spaces to prosets.

More generally, this is a functor from topoi to prosets: this carries a topos \mathbf{X} to the composite

$$\pi_0(\mathbf{X}) := \Gamma_{\mathbf{X},*} \Gamma_{\mathbf{X}}^* : \mathbf{Set} \rightarrow \mathbf{X} \rightarrow \mathbf{Set} .$$

Our claim is that the proset $\pi_0^{top}(X)$ is closely related to – and even identifiable with – the set $\pi_0(X)$. For a relatively nice class of topological spaces, $\pi_0(X)$ has a simple universal property.

0.24. Let us consider the category \mathbf{Top}^{ng} of *numerically generated* topological spaces – these are topological spaces X with the property that a subset $U \subseteq X$ is open if and only if, for any continuous map $\gamma : \mathbf{R}^n \rightarrow X$, the set $\gamma^{-1}U$ is open.

Of course any discrete space is numerically generated, so the assignment $S \mapsto S^{disc}$ is a functor $\mathbf{Set} \rightarrow \mathbf{Top}^{ng}$. This functor has a left adjoint, $\pi_0 : \mathbf{Top}^{ng}$; in other words, for any numerically generated topological space X , the continuous maps $X \rightarrow S^{disc}$ are in bijective correspondence with the maps $\pi_0(X) \rightarrow S$.

For any *numerically generated* topological space X

0.25 Definition. A morphism of topoi $p_* : \mathbf{X} \rightarrow \mathbf{Y}$ is *étale* if and only if the left adjoint p^* admits a further left adjoint $p_!$ that identifies \mathbf{X} with the overcategory $\mathbf{Y}_{/p_!(1_X)}$.

0.26 Construction. Let \mathbf{X} and \mathbf{Y} be topoi, and let $f_* : \mathbf{X} \rightarrow \mathbf{Y}$ be a geometric morphism. Let $U \in \mathbf{Y}$ be an object, and let $\alpha : 1_{\mathbf{X}} \rightarrow f^*U$ be a morphism of \mathbf{X} . Define a functor $F^* : \mathbf{Y}_{/U} \rightarrow \mathbf{X}$ that carries an object $[V \rightarrow U]$ of $\mathbf{Y}_{/U}$ to the object $1_{\mathbf{X}} \times_{f^*U} f^*V$; this functor admits a right adjoint F_* , which is a morphism of topoi. If we write $p_* : \mathbf{Y}_{/U} \rightarrow \mathbf{Y}$ for the canonical étale morphism, then we have an isomorphism $f^* \cong F^* \circ p^*$, and it is a tedious but routine check to confirm that this defines a functor

$$\mathrm{Map}_{\mathbf{X}}(1_{\mathbf{X}}, f^*U) \rightarrow \mathrm{Fun}_{*,/\mathbf{X}}(\mathbf{X}, \mathbf{Y}_{/U}) .$$

Monodromy representations

0.27 Definition. Let X be a topological space.

A *locally constant sheaf* of sets L is a sheaf for which there exists an open cover $\{U_\alpha\}_{\alpha \in A}$ such that for any $\alpha \in A$, the sheaf $L|_{U_\alpha}$ is constant on U_α . We will call such an open cover a *trivialising* open cover of X for L .

A *local system* is a locally constant sheaf for which there is a finite trivialising open cover.

0.28. Let X be a topological space, and let L be a locally constant sheaf on X . Any refinement of a trivialising open cover of X for L is again trivialising. Note also that if X is quasicompact, then any locally constant sheaf is a local system.

0.29 Example. Consider the interval $I := [0, 1]$. Let L be a local system on I . A finite trivialising open cover can be refined to a finite trivialising open cover $\{U_1, \dots, U_n\}$ in which each U_i is an interval.

Now let U_i and U_j be two of these intervals; their intersection is either empty or else an interval. Assume that their intersection is in fact nonempty; then their union is again an interval. Let S be a set such that $L|_{U_i}$ is constant at S . Since U_i and U_j intersect, it follows that $L|_{U_j}$ is constant at S as well. The sheaf condition gives an equaliser

$$L(U_i \cup U_j) \rightarrow S \times S \rightrightarrows S \times S \times S \times S,$$

where the top arrow is $(s, t) \mapsto (s, s, t, t)$ and the bottom arrow is $(s, t) \mapsto (s, t, s, t)$ which now implies that $L(U_i \cup U_j) \cong S$ in a way that is compatible with the restrictions to U_i and U_j . Thus we obtain a morphism η from the constant sheaf at S on $U_i \cup U_j$ to the restriction $L|_{(U_i \cup U_j)}$ that restricts to an isomorphism on U_i and U_j . Thus it follows that η is an isomorphism.

Thus any finite trivialising open cover consisting of $n > 1$ intervals can be replaced by a finite trivialising open cover consisting of $n - 1$ intervals. By induction, it follows that I itself is a trivialising open cover, whence L is a constant sheaf.

We thus conclude that any local system on the interval I is in fact constant!

In the course of this discussion, we encountered some facts that will be useful to us more generally.

0.30 Lemma. *Let X be a topological space, and let F be a sheaf of sets on X . Then if U and V are open sets such that $F|_U$ and $F|_V$ are constant, then*

1 Stratified topology

Posets and stratifications

1.1 Construction. Let P be a poset.

A *sieve* is a subset $Z \subseteq P$

Dually, a *cosieve*

The *Alexandroff topology* on P

1.2 Example. Let us consider the poset $[1] := \{0 < 1\}$.

1.3 Example. More generally, let us consider the linearly ordered poset $[n]$.

1.4 Example. Let P and Q be posets, and let us consider the product poset $P \times Q$.

1.5 Example. For any set S , we can contemplate the trivial poset structure.

1.6 Construction. For any Kolmogoroff topological space X , the *specialisation poset*

1.7 Proposition. *The formation of the Alexandroff topology of a finite poset and the formation of the specialisation poset of a finite Kolmogoroff topological space are mutually inverse equivalences of categories.*

1.8 Notation. Let P be a finite poset P . We shall always regard P as endowed with its Alexandroff topology. We write \tilde{P} for the ∞ -category of sheaves (of spaces) on P .

1.9 Construction. Let P be a finite poset. The stalk of a sheaf F on P at a point $p \in P$

1.10 Proposition. *The construction above defines an equivalence of ∞ -categories*

$$\tilde{P} \simeq \text{Fun}(P, \mathbf{S}).$$